

552056

Mr. Rob Adams
Adams Technology Systems
5145 Forest Run Trace - Suite B
Alpharetta, GA 30022-4504

Dear Mr. Adams:

DEC 16 1998

Subject: Amendment - Add Vegetables
CP Diazinon Lawn & Garden WB Ready-To-Use
EPA Registration Number 67572-1
Your submission dated August 9, 1998

The amendment referred to above, submitted in connection with a registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling change below before you release the product for shipment bearing the amended labeling:
 - a) For Vegetables, in the definition of PHI under the Pre-Harvest Interval table add "to wait" between "days" and "between".
3. Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide/Rodenticide Branch
Registration Division (7505C)

12727

ACCEPTED
with COMMENTS
in EPA Letter Dated

DEC 16 1998

LABELING FOR THE 1 GALLON READY-TO-USE PRODUCT

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

67572-1

- *Ready-To-Use
- *Convenient, Pre-Measured, No Guess Work
- *No Mixing, No Mess, No Touch

CP DIAZINON LAWN & GARDEN WB READY-TO-USE

- A Water-Based Ready-To-Use Spray that contains Diazinon Insecticide.
- Controls: Ants, Aphids, Chinch Bugs, Cutworms, Flea Beetles, Grubs, Mites, Sod Webworms and many others.
- Protects: Lawns, Fruits and Nuts, Roses, Flowers, Trees, Shrubs.
- Easy to Apply to Lawn or Garden
- This Product must Not Be Used on Golf Courses and Sod Farms.
- For Outdoor Use on the Home Lawn and the Home Garden Only.

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate

0.058%

INERT INGREDIENTS:

99.942%

TOTAL

100.000%

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel for Precautionary Statements

Manufactured by:
CONTRACT PACKAGING, Inc.
PO BOX 1508 COVINGTON GA 30209-1508

EPA EST #67572-GA-1

EPA REG #67572-1

Net Contents: 1 gallon

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY OR ILLEGAL RESIDUES

This is a water-based ready-to-use product which must be shaken before application.

This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. This product works fast and application should be repeated only as directed to maintain control.


NOTES: Do not use inside the home. Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since injury may occur. Do not apply to food crops grown in greenhouses.

PRECAUTIONS: Wear a long-sleeved shirt and long-legged pants when spraying. Spray with the wind to your back. Do not spray on windy days. If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

HOW TO APPLY

Step #1: Determine size of spraying job. For lawns, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Application should be repeated only as directed to maintain control.

Step #2: Spray as directed in rate chart. Thorough coverage is important. Provide enough spray coverage to reach the top of mature trees or limit use to small trees, dwarf trees and container grown trees. See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.



LAWN PEST CONTROL

PRECAUTIONS: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustine grass.

To control pests listed below, apply recommended rates:

PESTS	RATE	REMARKS
Lawn Chinch Bugs	1 gallon per 50 sq. ft.	Thoroughly wet grass a few hours before applying.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	1 gallon per 50 sq. ft.	Thoroughly wet grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1 gallon per 50 sq. ft.	NOTE: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevil	1 gallon per 50 sq. ft.	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Millipedes	1 gallon per 50 sq. ft.	Thoroughly wet grass a few hours before applying
Rhodesgrass Scales	1 gallon per 50 sq. ft.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	1 gallon per 50 sq. ft.	Apply any time between late July and early October. Water grass thoroughly after application.

Spray Pests Listed Below When They First Appear. Repeat as Necessary.

INSECT CONTROL ON ORNAMENTAL TREES, SHRUBS AND FLOWERS		
TREES & SHRUBS	INSECTS	TIMING/SPECIAL DIRECTIONS
Arborvitae, Azalea, Birch, Boxwood, Camellia, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew	Aphids, Apple and Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Oblique banded Leafrollers, Obscure Root Weevils, Omnivorous Leafhoppers, Pear Slugs, Scale Crawlers (Cottony cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies.	Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray. Mimosa Webworms: apply when adults first appear. Make two additional applications at 4-5 week intervals.

INSECT CONTROL ON ORNAMENTAL TREES, SHRUBS AND FLOWERS		
ROSES & FLOWERS	INSECTS	TIMING/SPECIAL DIRECTIONS
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses	Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

FRUITS AND NUTS

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

CROP PRE-HARVEST INTERVAL or the Number of Days to Wait Between the Last Application and Harvest	
Almonds	Dormant Spray Only
Apples	21 days
Apricots	21 days
Blackberries	7 days
Blueberries	7 days
Boysenberries	7 days
Cherries	21 days
Dewberries	7 days
Loganberries	7 days
Nectarines	21 days
Peaches	21 days
Pears	21 days
Plums	21 days
Prunes	21 days
Raspberries	7 days
Strawberries	5 days
Walnuts	45 days


NOTE: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.



7227

FRUITS AND NUTS		
CROP	PEST	TIMING/SPECIAL DIRECTIONS
Almonds (CA only)	San Jose Scale, Parlatoria Scale	Apply only as a dormant spray. Do not apply more than 1 gallon per 83.3 sq. ft. of home orchard. This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.
Apples	Aphids, Apple maggots, Codling moths, Eyespotted budmoths, Fruit Tree Leafrollers, Leafhoppers, Mealybugs, Scale Crawlers (San Jose Forbes), Spider Mites	Apply as full cover spray. Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. NOTE: Resistant apple maggots may not be controlled by diazinon or similar type insecticides.
Apricots	Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray. Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Blueberries	Aphids, Blueberry maggots, Cherry fruitworms, Cranberry fruitworms, Thrips, Two-spotted spider mites	Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 1 gallon per 125 sq. ft. of blueberry patch. Do not make more than 2 applications per season. Allow a minimum of 7 days between the last application and harvest.
Caneberries (CA, OR, WA only): Blackberries Boysenberries Dewberries Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	Apply as a full cover spray. Do not apply more than 1 gallon per 100 sq. ft. of caneberry patch. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.
Cherries	Black Cherry Aphids, Cherry Fruit flies, Cherry Fruitworms (Maggots), Fruitree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	Apply as a full cover spray. Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Pears	Aphids, Apple maggots, Codling moths, Mealybugs, Scale crawlers (San Jose, Forbes), Spider Mites, Tentiform Leafminers Pear Psylla	Apply as a full cover spray Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. NOTE: Resistant apple maggots will not be controlled by diazinon or similar type insecticides.

FRUITS AND NUTS		
CROP	PEST	TIMING/SPECIAL DIRECTIONS
Plums & Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	Apply as a full cover spray Do not apply more than 1 gallon per 125 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	Apply as a full cover spray Do not apply more than 1 gallon per 166.6 sq. ft. of strawberry patch. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	Apply as a full cover spray Do not apply more than 1 gallon per 83.3 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.



9827

VEGETABLES

For foliar applications, CP Diazinon Lawn and Garden Ready-To-Use should be applied as recommended in the table below to assure thorough coverage of the foliage.


Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (*PHI) for Foliar Applications	
CROP	* PRE-HARVEST INTERVAL
Beans (succulent varieties only: Lima, Pole & Snap)	7 days
Broccoli	7 days
Cabbage	21 days
Cantaloupes	3 days
Carrots	14 days
Casabas	3 days
Cauliflower	7 days
Chinese Broccoli	10 days
Chinese Cabbage	10 days
Collards	10 days
Crenshaws	3 days
Cucumbers	7 days
Endive (Escarole)	14 days
Honeydew Melons	3 days
Kale	10 days
Lettuce	14 days
Muskmelons	3 days
Onions (Bulb and Green)	14 days
Parsnips	14 days
¹ Peas	7 days
Peppers	5 days
Persian Melons	3 days
Potatoes (Irish)	35 days
Radishes	14 days ¹
Rutabagas	14 days

CROP	* PRE-HARVEST INTERVAL
Spinach	14 days
Squash, Summer	7 days
Squash, Winter	3 days
Sweet Corn	7 days
Tomatoes	1 day
Turnips (Roots & Tops)	14 days
Watermelons	3 days

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas, or dried black-eyed peas.



11 2 27

VEGETABLES

For application to the foliage, thoroughly spray all plant parts.

CROP	PEST	RATE	TIMING / SPECIAL DIRECTIONS
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
<p>NOTE 1: Do not apply this product to dried bean varieties such as pinto beans or dried limas. NOTE 2: To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.</p>			
Broccoli and Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Broccoli, Chinese Cabbage (Bok Choy and Napa), Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cucumbers	Aphids, Thrips, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

CROP	PEST	RATE	TIMING / SPECIAL DIRECTIONS
Melons (cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Aphids, Thrips, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (Bulb and Green)	Onion Thrips	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
<p>NOTE 1: Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. NOTE 2: To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.</p>			
Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Radishes	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Rutabagas	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids, Thrips, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.

CROP	PEST	RATE	TIMING / SPECIAL DIRECTIONS
Squash, Winter	Aphids, Thrips, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last applications.
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (Drosophila Spp.)	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 gal covers 1000 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

STORAGE AND DISPOSAL

STORAGE: Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68° or above) and allow to thaw. Shake well before using.

DISPOSAL: Do not reuse container. Wrap and place in trash.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.


IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.



15727

ACCEPTED
with COMMENTS
in EPA Letter Dated

DEC 16 1998

LABELING FOR THE 24 oz. READY-TO-USE PRODUCT

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
67572-1

- *Ready-To-Use
- *Convenient, Pre-Measured, No Guess Work
- *No Mixing, No Mess, No Touch

CP DIAZINON LAWN & GARDEN WB READY-TO-USE

- A Water-Based Ready-To-Use Spray that contains Diazinon Insecticide.
- Controls: Ants, Aphids, Chinch Bugs, Cutworms, Flea Beetles, Grubs, Mites, Sod Webworms and many others.
- Protects: Lawns, Fruits and Nuts, Roses, Flowers, Trees, Shrubs.
- Easy to Apply to Lawn or Garden
- This Product must Not Be Used on Golf Courses and Sod Farms.
- For Outdoor Use on the Home Lawn and the Home Garden Only.

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)	
phosphorothioate	0.058%
INERT INGREDIENTS:	99.942%
	TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel for Precautionary Statements

Manufactured by:
CONTRACT PACKAGING, Inc.
 PO BOX 1508 COVINGTON GA 30209-1508

EPA EST #67572-GA-1

EPA REG #67572-1

Net Contents 24 fl. oz.

LAWN PEST CONTROL

PRECAUTIONS: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustine grass.

To control pests listed below, apply recommended rates:

PESTS	RATE	REMARKS
Lawn Chinch Bugs	24 oz. per 9.38 sq. ft.	Thoroughly wet grass a few hours before applying.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	24 oz. per 9.38 sq. ft.	Thoroughly wet grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	24 oz. per 9.38 sq. ft.	NOTE: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevil	24 oz. per 9.38 sq. ft.	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Millipedes	24 oz. per 9.38 sq. ft.	Thoroughly wet grass a few hours before applying
Rhodesgrass Scales	24 oz. per 9.38 sq. ft.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	24 oz. per 9.38 sq. ft.	Apply any time between late July and early October. Water grass thoroughly after application.

Spray Pests Listed Below When They First Appear. Repeat as Necessary.

INSECT CONTROL ON ORNAMENTAL TREES, SHRUBS AND FLOWERS		
TREES & SHRUBS	INSECTS	TIMING/SPECIAL DIRECTIONS
Arborvitae, Azalea, Birch, Boxwood, Camellia, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew	Aphids, Apple and Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Oblique banded Leafrollers, Obscure Root Weevils, Omnivorous Leaf tiers, Pear Slugs, Scale Crawlers (Cottony cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies.	Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray. Mimosa Webworms: apply when adults first appear. Make two additional applications at 4-5 week intervals.

INSECT CONTROL ON ORNAMENTAL TREES, SHRUBS AND FLOWERS		
ROSES & FLOWERS	INSECTS	TIMING/SPECIAL DIRECTIONS
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses	Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.


FRUITS AND NUTS

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.


CROP PRE-HARVEST INTERVAL or the Number of Days to Wait Between the Last Application and Harvest	
Almonds	Dormant Spray Only
Apples	21 days
Apricots	21 days
Blackberries	7 days
Blueberries	7 days
Boysenberries	7 days
Cherries	21 days
Dewberries	7 days
Loganberries	7 days
Nectarines	21 days
Peaches	21 days
Pears	21 days
Plums	21 days
Prunes	21 days
Raspberries	7 days
Strawberries	5 days
Walnuts	45 days

NOTE: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.



FRUITS AND NUTS		
CROP	PEST	TIMING/SPECIAL DIRECTIONS
Almonds (CA only)	San Jose Scale, Parlatoria Scale	Apply only as a dormant spray. Do not apply more than 24 oz. per 15.6 sq. ft. of home orchard. This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.
Apples	Aphids, Apple maggots, Codling moths, Eyespotted budmoths, Fruit Tree Leafrollers, Leafhoppers, Mealybugs, Scale Crawlers (San Jose Forbes), Spider Mites	Apply as full cover spray. Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. NOTE: Resistant apple maggots may not be controlled by diazinon or similar type insecticides.
Apricots	Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray. Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Blueberries	Aphids, Blueberry maggots, Cherry fruitworms, Cranberry fruitworms, Thrips, Two-spotted spider mites	Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 24 oz. per 24.4 sq. ft. of blueberry patch. Do not make more than 2 applications per season. Allow a minimum of 7 days between the last application and harvest.
Caneberries (CA, OR, WA only): Blackberries Boysenberries Dewberries Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	Apply as a full cover spray. Do not apply more than 24 oz. per 18.7 sq. ft. of caneberry patch. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.
Cherries	Black Cherry Aphids, Cherry Fruit flies, Cherry Fruitworms (Maggots), Fruitree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	Apply as a full cover spray. Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Pears	Aphids, Apple maggots, Codling moths, Mealybugs, Scale crawlers (San Jose, Forbes), Spider Mites, Tentiform Leafminers Pear Psylla	Apply as a full cover spray Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. NOTE: Resistant apple maggots will not be controlled by diazinon or similar type insecticides.

FRUITS AND NUTS		
CROP	PEST	TIMING/SPECIAL DIRECTIONS
Plums & Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	Apply as a full cover spray Do not apply more than 24 oz. per 24.4 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	Apply as a full cover spray Do not apply more than 24 oz. per 31.2 sq. ft. of strawberry patch. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	Apply as a full cover spray Do not apply more than 24 oz. per 15.6 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.



22 7 27

VEGETABLES

For foliar applications, CP Diazinon Lawn and Garden Ready-To-Use should be applied as recommended in the table below to assure thorough coverage of the foliage.


Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (*PHI) for Foliar Applications	
CROP	* PRE-HARVEST INTERVAL
Beans (succulent varieties only: Lima, Pole & Snap)	7 days
Broccoli	7 days
Cabbage	21 days
Cantaloupes	3 days
Carrots	14 days
Casabas	3 days
Cauliflower	7 days
Chinese Broccoli	10 days
Chinese Cabbage	10 days
Collards	10 days
Crenshaws	3 days
Cucumbers	7 days
Endive (Escarole)	14 days
Honeydew Melons	3 days
Kale	10 days
Lettuce	14 days
Muskmelons	3 days
Onions (Bulb and Green)	14 days
Parsnips	14 days
Peas	7 days
Peppers	5 days
Persian Melons	3 days
Potatoes (Irish)	35 days
Radishes	14 days
Rutabagas	14 days

CROP	* PRE-HARVEST INTERVAL
Spinach	14 days
Squash, Summer	7 days
Squash, Winter	3 days
Sweet Corn	7 days
Tomatoes	1 day
Tumips (Roots & Tops)	14 days
Watermelons	3 days

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas, or dried black-eyed peas.



VEGETABLES

For application to the foliage, thoroughly spray all plant parts.

CROP	PEST	RATE	TIMING / SPECIAL DIRECTIONS
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
<p>NOTE 1: Do not apply this product to dried bean varieties such as pinto beans or dried limas. NOTE 2: To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.</p>			
Broccoli and Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Broccoli, Chinese Cabbage (Bok Choy and Napa), Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cucumbers	Aphids, Thrips, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant Leafminers will not be controlled by diazinon or similar type insecticides.

05 7 27

CROP	PEST	RATE	TIMING / SPECIAL DIRECTIONS
Melons (cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Aphids, Thrips, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (Bulb and Green)	Onion Thrips	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
NOTE 1: Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. NOTE 2: To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.			
Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Radishes	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Rutabagas	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids, Thrips, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.

26 27

CROP	PEST	RATE	TIMING / SPECIAL DIRECTIONS
Squash, Winter	Aphids, Thrips, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last applications.
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (Drosophila Spp.)	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	24 oz covers 187.5 square feet of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

STORAGE AND DISPOSAL

STORAGE: Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68° or above) and allow to thaw. Shake well before using.

DISPOSAL: Do not reuse container. Wrap and place in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

