

CP DIAZINON LAWN & GARDEN WB READY-TO-USE

Ready-to-Use
Convenient, Pre-Measured, No Guess Work
No Mixing, No Mess, No Touch

A Water-Based Ready-to-Use Spray that Contains Diazinon Insecticide

Controls: Ants, Aphids, Chinch Bugs, Cutworms, Flea Beetles, Grubs, Mites, Sod Webworms and Many Others [More]

Protects: Lawns, Fruits and Nuts, Roses, Flowers, Trees and Shrubs
Easy to Apply to Lawn or Garden

This Product Must Not be Used on Golf Courses and Sod Farms

For Outdoor Use Only on the Home Lawn and the Home Garden

ACTIVE INGREDIENT:

| | |
|-----------------------------------|----------------|
| Diazinon (CAS No. 333-41-5) | 0.058% |
| OTHER INGREDIENTS | <u>99.942%</u> |
| Total | 100.000% |

Keep Out of Reach of Children
CAUTION
See back panel for additional precautionary statements.

Manufactured by:
CONTRACT PACKAGING, INC.
P.O. Box 1508
Covington, GA 30015-1508

ACCEPTED

NOV - 9 2000
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67572-1

EPA Reg. No. 67572-1

EPA Est. 67572-GA-1

Net Contents: 1 gallon [32 fl. oz.] [24 fl. oz.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets. If product freezes, place container in a warm room (68°F or above) and allow to thaw. Shake well before using. **Disposal:** Do not reuse empty container; wrap and put in trash.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL WILL RESULT IN POOR INSECT CONTROL, CROP INJURY OR ILLEGAL RESIDUES

This is a water-based ready-to-use product that must be shaken before application.

This product kills many outdoor insects that damage lawns, roses, flowers, trees, shrubs, fruits and nuts, and insects that annoy pets and people. Contains diazinon insecticide to provide positive control of many "hard-to-kill" sucking and chewing insects. This product works fast. Repeat application only as directed to maintain control.

Notes: Do not use inside the home. Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since injury will occur. Do not apply to food crops grown in greenhouses.

Precautions: Wear a long-sleeved shirt and long-legged pants when spraying. Spray with the wind to your back. Do not spray on windy days. If clothing becomes wet from spray during use, remove clothing after spraying, thoroughly wash affected body areas with soap and water and launder clothing before wearing again.

How to Apply

Step #1: Determine size of spraying job. For lawns, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear.

Step #2: Spray as directed in rate chart. Thorough coverage is important. Provide enough spray coverage to reach the top of mature trees or limit use to small trees, dwarf trees and container-grown trees. See chart for complete list of use sites, crops and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Lawn Pest Control

Precautions: Birds, especially waterfowl (i.e., ducks and geese), feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustine grass. To control pests listed below, apply at recommended rates.

| Pests | Rate | Remarks |
|---|---|---|
| Lawn Chinch Bugs | 1 gallon per 50 sq. ft. | Thoroughly wet grass a few hours before applying. |
| Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs) | { 1 gallon container} [or 32 oz. per 13 sq. ft. {32 oz. container} or 24 oz. per 9.38 sq. ft. {24 oz. container}] | Thoroughly wet grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and thoroughly water before treatment, but delay application until grass is dry. |
| Brown Dog Ticks, Chiggers, Fleas | | Note: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house. |
| Hyperodes Weevil | | Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application. |
| Millipedes | | Thoroughly wet grass a few hours before applying. |
| Rhodesgrass Scales | | Apply when crawlers first emerge. |
| White Grubs of Japanese Beetle, European Chafer, Southern Chafer | | Apply any time between late July and early October. Thoroughly Water grass after application. |

Insect Control on Ornamental Trees, Shrubs and Flowers

Spray pests below when they first appear. Repeat as necessary.

| Trees and Shrubs | Insects | Timing/Special Directions |
|--|--|--|
| Arborvitae, Azalea, Birch, Boxwood, Camellia, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pint, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew | Aphids, Apple and Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European, Red, Privet, Two-spotted Spider), Oak Loopers, Oblique-banded Leafrollers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (Cottony Cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Ten Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa) Whiteflies | Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Penetrate dense foliage with spray. Mimosa Webworms: Apply when adults first appear. Make two additional applications at 4-5 week intervals. |
| Roses and Flowers | Insects | Timing/Special Directions |
| Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses | Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies | Apply as a thorough foliar spray. Spray underside of leaves and penetrate dense foliage. |

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites and crawler stages of scale insects are first noted. Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

| Crop Pre-Harvest Interval (or the number of Days to Wait Between the Last Application and Harvest) | |
|---|--------------------|
| Almonds | Dormant Spray Only |
| Apples | 21 days |
| Apricots | 21 days |
| Blackberries | 7 days |
| Blueberries | 7 days |
| Boysenberries | 7 days |
| Cherries | 21 days |
| Dewberries | 7 days |
| Loganberries | 7 days |
| Nectarines | 21 days |
| Peaches | 21 days |
| Pears | 21 days |
| Plums | 21 days |
| Prunes | 21 days |
| Raspberries | 7 days |
| Strawberries | 5 days |
| Walnuts | 45 days |

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

| Crop | Pests | Timing/Special Directions |
|--|--|---|
| Almonds (CA only) | San Jose Scale, Parlatoria Scale | Apply only as a dormant spray. Do not apply more than 1 gal. per 83.3 sq. ft. [32 oz. per 21 sq. ft.] [24 oz. per 15.6 sq. ft.] of home orchard. This product may be used on almonds only in CA. Do not use this product on almonds grown in other states. |
| Apples | Aphids, Apple Maggots, Codling Moths, Eyespotted Budmoths, Fruit Tree Leafrollers, Leafhoppers, Mealybugs, Scale Crawlers (San Jose, Forbes), Spider Mites | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 38 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots may not be controlled by diazinon or similar type insecticides. |
| Apricots | Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 33 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Blueberries | Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites | Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 33 sq. ft.] [24 oz. per 24.4 sq. ft.] of blueberry patch. Do not make more than 2 applications per season. Allow a minimum of 7 days between the last application and harvest. |
| Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries | Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips | Apply as a full cover spray. Do not apply more than 1 gal. per 100 sq. ft. [32 oz. per 25 sq. ft.] [24 oz. per 18.7 sq. ft.] of caneberry patch. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states. |
| Cherries | Black Cherry Aphids, Cherry Fruit Flies, Cherry Fruitworms (Maggots), Fruittree Leafrollers, Mites (Cherry, Rust), Scale Crawlers (San Jose) | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 25 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest. |
| Nectarines | Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 33 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Peaches | Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 33 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Pears | Aphids, Apple Maggots, Codling Moths, Mealybugs, Scale Crawlers (San Jose, Forbes), Spider Mites, Tentiform Leafminers, Pear Psylla | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 33 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 application per season a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots will not be controlled by diazinon or similar type insecticides. |
| Plums and Prunes | Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red) | Apply as a full cover spray. Do not apply more than 1 gal. per 125 sq. ft. [32 oz. per 33 sq. ft.] [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest. |
| Strawberries | Aphids, Two-spotted Spider Mites, Strawberry Leafrollers | Apply as a full cover spray. Do not apply more than 1 gal. per 166.6 sq. ft. [32 oz. per 42 sq. ft.] [24 oz. per 31.2 sq. ft.] of strawberry patch. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest. |
| Walnuts (CA only) | Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers | Apply as a full cover spray. Do not apply more than 1 gal. per 83.3 sq. ft. [32 oz. per 22 sq. ft.] [24 oz. per 15.6 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states. |

Vegetables

For foliar applications, CP Diazinon Lawn and Garden Ready-to-Use should be applied as recommended in the tables below to assure thorough coverage of the foliage. Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

| Pre-Harvest Intervals (PHI*) for Foliar Applications | |
|---|---------|
| Beans (succulent varieties only: Lima, Pole and Snap) | 7 days |
| Broccoli | 7 days |
| Cabbage | 21 days |
| Cantaloupes | 3 days |
| Carrots | 14 days |
| Casabas | 3 days |
| Cauliflower | 7 days |
| Chinese Broccoli | 10 days |
| Chinese Cabbage | 10 days |
| Collards | 10 days |
| Crenshaws | 3 days |
| Cucumbers | 7 days |
| Endive (Escarole) | 14 days |
| Honeydew Melons | 3 days |
| Kale | 10 days |
| Lettuce | 14 days |
| Muskmelons | 3 days |
| Onions (Bulb and Green) | 14 days |
| Parsnips | 14 days |
| Peas ¹ | 7 days |
| Peppers | 5 days |
| Persian Melons | 3 days |
| Potatoes (Irish) | 35 days |
| Radishes | 14 days |
| Rutabagas | 14 day |
| Spinach | 14 days |
| Squash, Summer | 7 days |
| Squash, Winter | 3 days |
| Sweet Corn | 7 days |
| Tomatoes | 1 days |
| Turnips (Roots and Tops) | 14 days |
| Watermelons | 3 days |

*PHI = Pre-Harvest Interval, or the number of days to wait between the last application and harvest.

¹Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas.

For application to the foliage, thoroughly spray all plant parts.

| Crop | Pests | Rate | Timing/Special Directions |
|---|---|--|--|
| Beans (Lima, Pole and Snap) [succulent varieties only] | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | 1 gallon covers 1000 sq. ft. of garden [24 oz. covers] 187.5 sq. ft. of garden [32 oz. covers] 250 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest. |
| | Notes: 1) Do not apply this product to dried bean varieties such as pinto beans or dried limas; 2) To protect bees, do not apply to bean foliage treatments if the beans or weeds in the treatment area are in bloom. | | |
| Broccoli and Cabbage | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 gallon covers 1000 sq. ft. of garden [24 oz. covers] 187.5 sq. ft. of garden [32 oz. covers] 250 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |

| | Pests | Rate | Timing/Special Directions |
|---|---|---|---|
| Carrots | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 gallon covers 1000 sq. ft. of garden [24 oz. covers 187.5 sq. ft. of garden] [32 oz. covers 250 sq. ft. of garden] | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Cauliflower | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Chinese Broccoli, Chinese Cabbage (Bok Choy and Napa), Collards | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Cucumbers | Aphids, Thrips, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. |
| Endive (Escarole) | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Kale | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Lettuce | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians and Hybrids of these) and Watermelons | Aphids, Thrips, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest. |
| Onions (Bulb and Green) | Onion Thrips | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. |
| Parsnips | Aphids, Flea Beetles, Leafhoppers Dipterous Leafminers, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Peas (succulent only) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest. |
| Notes: 1) Do not apply product to dried pea varieties such as split peas or dried black-eyed peas; 2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom. | | | |

| Crop | Pests | Rate | Timing/Special Directions |
|-------------------------|---|--|---|
| Peppers | Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers | 1 gallon covers 1000 sq. ft. of garden [24 oz. covers | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest. |
| Potatoes | Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers | 187.5 sq. ft. of garden] [32 oz. covers 250 sq. ft. of garden] | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Radishes | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | | Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Rutabagas | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Spinach | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Squash, Summer | Aphids, Thrips, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. |
| Squash, Winter | Aphids, Thrips, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest. |
| Sweet Corn | Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application. |
| Tomatoes | Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.) | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest. |
| Turnips and Turnip Tops | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Thoroughly wash thoroughly with soap and water after use handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

First Aid

| | |
|--|---|
| If Swallowed | <ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. |
| If in Eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If Inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| If on Skin or Clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment advice.</p> | |
| <p>Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.</p> | |

Environmental Hazards

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl (i.e. ducks and geese), feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

To obtain additional information or in case of an emergency call 770-787-1977.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}