



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Contract Packaging, Inc.
c/o Ms. Kim Davis
RegWest Company
30856 Rocky Road
Greeley, CO 80631-9375

DEC - 6 2001

Dear Ms. Davis:

Subject: Amendment - Comply with 2000 Diazinon MOA;
Delete Almonds and Pole Beans;
Update First Aid per PR Notice 2001-1; and
Update CSF/replacement diazinon source
CP Diazinon Lawn & Garden WB Ready-to-Use
EPA Registration Number 67572-1
Your submissions dated June 14 and 21, 2001

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. Under the heading DIRECTIONS FOR USE, slightly revise the reformulation prohibition to read "Reformulation or repackaging of this product is prohibited."
 - b. Under the heading NOTES, revise "Do not apply to food crops grown in greenhouses." to read "Do not apply this product in greenhouses."
 - c. Under the heading VEGETABLES, revise "Do not apply to food crops grown in greenhouses." to read "Do not apply this product in greenhouses."
 - d. In the Vegetables PHI table, revise the entry for "Casabas" to "Casaba Melons" and the entry for "Crenshaws" to "Crenshaw Melons".
 - e. In the Hazards to Humans and Domestic Animals text, revise "Do not use inside the home." to read "Do not use inside the home or in greenhouses."

f. In the If Swallowed advice of the First Aid section, add as the fourth item:

- Do not give anything by mouth to an unconscious or convulsing person.

3. Submit three (3) copies of your final printed labeling before you release the product for shipment.

4. Comply with all applicable provisions of the 2000 Memorandum of Agreement Regarding Pesticide Products Containing Diazinon (i.e., the Diazinon MOA). In accordance with the Diazinon MOA you have requested the following revisions to the conditions of registration for this diazinon product with outdoor non-agricultural (i.e., home and garden) uses:

- a. You have requested under section 6(f) of FIFRA that this registration be voluntarily canceled on a date not later than December 31, 2004.
- b. You may not manufacture, label, or release for shipment this product after June 30, 2003.
- c. You may not sell or distribute this product after August 31, 2003.
- d. The registration for this product shall expire and be canceled on December 31, 2004.
- e. You will send to every person who directly purchases this product (i.e., to every seller) a notification of Product Recovery Program to be conducted by Syngenta and Makhteshim after December 31, 2004.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The Basic Confidential Statement of Formula (CSF) dated June 14, 2001 which identifies a replacement diazinon source Manufacturing Use Product is acceptable and has been added to this product's registration file.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

3 7 10

CP DIAZINON LAWN & GARDEN WB READY-TO-USE

Ready-to-Use

Convenient, Pre-Measured, No Guess Work

No Mixing, No Mess, No Touch

A Water-Based Ready-to-Use Spray that Contains Diazinon Insecticide

Controls: Ants, Aphids, Chinch Bugs, Cutworms, Flea Beetles, Grubs, Mites, Sod Webworms and Many Others [More] as Listed in Directions

Protects: Lawns, Fruits and Nuts, Roses, Flowers, Trees and Shrubs
Easy to Apply to Lawn or Garden

This Product Must Not be Used on Golf Courses and Sod Farms

For Outdoor Use Only on the Home Lawn and the Home Garden

Active Ingredient:

Diazinon (CAS No. 333-41-5)	0.058%
Other Ingredients	99.942%
Total	100.000%

Keep Out of Reach of Children
CAUTION
See back panel for additional precautionary statements.

Manufactured by:
CONTRACT PACKAGING, INC.
P.O. Box 1508
Covington, GA 30015-1508

ACCEPTED
with COMMENTS
In EPA Letter Dated:
DEC - 6 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67572-1

EPA Reg. No. 67572-1

EPA Est. 67572-GA-1

Net Contents: 1 quart (32 fl. oz.) [24 fl. oz.]

4 8 10

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Reformulation or repackaging this product is prohibited.

Storage and Disposal

Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets. If product freezes, place container in a warm room (68°F or above) and allow to thaw. Shake well before using. **Disposal:** Do not reuse empty container; wrap and put in trash.

Failure to follow the directions for use and precautions on this label will result in poor insect control, crop injury or illegal residues.

This is a water-based ready-to-use product that must be shaken before application.

This product kills many outdoor insects that damage lawns, roses, flowers, trees, shrubs, fruits and nuts, and insects that annoy pets and people. Contains diazinon insecticide to provide positive control of many "hard-to-kill" sucking and chewing insects. This product works fast. Repeat application only as directed to maintain control.

Notes: Do not use inside the home. Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since injury will occur. Do not apply to food crops grown in greenhouses.

Precautions: Wear a long-sleeved shirt and long-legged pants when spraying. Spray with the wind to your back. Do not spray on windy days. If clothing becomes wet from spray during use, remove clothing after spraying, thoroughly wash affected body areas with soap and water and launder clothing before wearing again.

How to Apply

Step #1: Determine size of spraying job. For lawns, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear.

Step #2: Spray as directed in rate chart. Thorough coverage is important. Provide enough spray coverage to reach the top of mature trees or limit use to small trees, dwarf trees and container-grown trees. See chart for complete list of use sites, crops and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Lawn Pest Control

Precautions: Birds, especially waterfowl (i.e., ducks and geese), feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustine grass. To control pests listed below, apply at recommended rates.

Pests	Rate	Remarks
Lawn Chinch Bugs	32 oz. per 13 sq. ft.	Thoroughly wet grass a few hours before applying.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (<i>Collembola</i>), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	{32 oz. container} [or 24 oz. per 9.38 sq. ft. {24 oz. container}]	Thoroughly wet grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and thoroughly water before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas		Note: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevil		Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Millipedes		Thoroughly wet grass a few hours before applying.
Rhodesgrass Scales		Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer		Apply any time between late July and early October. Thoroughly water grass after application.

Insect Control on Ornamental Trees, Shrubs and Flowers

Spray pests below when they first appear. Repeat as necessary.

Trees and Shrubs	Insects	Timing/Special Directions
Arbervitae, Azalea, Birch, Boxwood, Camellia, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pint, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew	Aphids, Apple and Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arbervitae, Azalea, Birch, Boxwood, Holly, Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European, Red, Privet, Two-spotted Spider), Oak Loopers, Oblique-banded Leafrollers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (Cottony Cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Ten Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa) Whiteflies	Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Penetrate dense foliage with spray. Mimosa Webworms: Apply when adults first appear. Make two additional applications at 4-5 week intervals.
Roses and Flowers	Insects	Timing/Special Directions
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses	Aphids, Dipterous Leafminers, Euonymous Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies	Apply as a thorough foliar spray. Spray underside of leaves and penetrate dense foliage.

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites and crawler stages of scale insects are first noted. Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop Pre-Harvest Interval (or the number of Days to Wait Between the Last Application and Harvest)	
Apples	21 days
Apricots	21 days
Blackberries	7 days
Blueberries	7 days
Boysenberries	7 days
Cherries	21 days
Dewberries	7 days
Loganberries	7 days
Nectarines	21 days
Peaches	21 days
Pears	21 days
Plums	21 days
Prunes	21 days
Raspberries	7 days
Strawberries	5 days
Walnuts	45 days

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

Crop	Pests	Timing/Special Directions
Apples	Aphids, Apple Maggots, Codling Moths, Eyespotted Budmoths, Fruit Tree Leafrollers, Leafhoppers, Mealybugs, Scale Crawlers (San Jose, Forbes), Spider Mites	Apply as a full cover spray. Do not apply more than 32 oz. per 38 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots may not be controlled by diazinon or similar type insecticides.
Apricots	Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray. Do not apply more than 32 oz. per 33 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Blueberries	Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites	Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 32 oz. per 33 sq. ft. [24 oz. per 24.4 sq. ft.] of blueberry patch. Do not make more than 2 applications per season. Allow a minimum of 7 days between the last application and harvest.
Caneberries (CA, OR, WA only): blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	Apply as a full cover spray. Do not apply more than 32 oz. per 25 sq. ft. [24 oz. per 18.7 sq. ft.] of caneberry patch. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.
Cherries	Black Cherry Aphids, Cherry Fruit Flies, Cherry Fruitworms (Maggots), Fruittree Leafrollers, Mites (Cherry, Rust), Scale Crawlers (San Jose)	Apply as a full cover spray. Do not apply more than 32 oz. per 25 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray. Do not apply more than 32 oz. per 33 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	Apply as a full cover spray. Do not apply more than 32 oz. per 33 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Pears	Aphids, Apple Maggots, Codling Moths, Mealybugs, Scale Crawlers (San Jose, Forbes), Spider Mites, Tentiform Leafminers, Pear Psylla	Apply as a full cover spray. Do not apply more than 32 oz. per 33 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 application per season a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots will not be controlled by diazinon or similar type insecticides.
Plums and Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	Apply as a full cover spray. Do not apply more than 32 oz. per 33 sq. ft. [24 oz. per 24.4 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	Apply as a full cover spray. Do not apply more than 32 oz. per 42 sq. ft. [24 oz. per 31.2 sq. ft.] of strawberry patch. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	Apply as a full cover spray. Do not apply more than 32 oz. per 22 sq. ft. [24 oz. per 15.6 sq. ft.] of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.

Vegetables

For foliar applications, CP Diazinon Lawn and Garden Ready-to-Use should be applied as recommended in the tables below to assure thorough coverage of the foliage. Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications	
Beans ¹ (succulent varieties only: Lima and Snap)	7 days
Broccoli	7 days
Cabbage	21 days
Cantaloupes	3 days
Carrots	14 days
Casabas	3 days
Cauliflower	7 days
Chinese Broccoli	10 days
Chinese Cabbage	10 days
Collards	10 days
Crenshaws	3 days
Cucumbers	7 days
Endive (Escarole)	14 days
Honeydew Melons	3 days
Kale	10 days
Lettuce	14 days
Muskmelons	3 days
Onions (Bulb and Green)	14 days
Parsnips	14 days
Peas ¹	7 days
Peppers	5 days
Persian Melons	3 days
Potatoes (Irish)	35 days
Radishes	14 days
Rutabagas	14 day
Spinach	14 days
Squash, Summer	7 days
Squash, Winter	3 days
Sweet Corn	7 days
Tomatoes	1 days
Turnips (Roots and Tops)	14 days
Watermelons	3 days

PHI = Pre-Harvest Interval, or the number of days to wait between the last application and harvest.

¹Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas.

For application to the foliage, thoroughly spray all plant parts.

Crop	Pests	Rate	Timing/Special Directions
Beans (Lima and Snap) [succulent varieties only]	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	24 oz. covers 187.5 sq. ft. of garden [32 oz. covers 250 sq. ft. of garden]	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Notes: 1) Do not apply this product to dried bean varieties such as pinto beans or dried limas; 2) To protect bees, do not apply to bean foliage treatments if the beans or weeds in the treatment area are in bloom.		
Broccoli and Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	24 oz. covers 187.5 sq. ft. of garden [32 oz. covers 250 sq. ft. of garden]	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

	Pests	Rate	Timing/Special Directions
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	24 oz. covers 187.5 sq. ft. of garden [32 oz. covers 250 sq. ft. of garden]	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Broccoli, Chinese Cabbage (Bok Choy and Napa), Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cucumbers	Aphids, Thrips, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians and Hybrids of these) and Watermelons	Aphids, Thrips, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (Bulb and Green)	Onion Thrips		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips	Aphids, Flea Beetles, Leafhoppers Dipterous Leafminers, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites		Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
	Notes: 1) Do not apply product to dried pea varieties such as split peas or dried black-eyed peas; 2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.		
Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	24 oz. covers 187.5 sq. ft. of garden [32 oz. covers 250 sq. ft. of garden]	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.

Crop	Pests	Rate	Timing/Special Directions
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	24 oz. covers 187.5 sq. ft. of garden [32 oz. covers 250 sq. ft. of garden]	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Radishes	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers		Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Rutabagas	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids, Thrips, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Squash, Winter	Aphids, Thrips, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application.
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.)		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers		Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Thoroughly wash with soap and water after use. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

First Aid

If Swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

First Aid (continued)

If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. [You may also contact 1-800-xxx-xxxx for emergency medical treatment advice.]</p>	
<p>Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.</p>	

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Environmental Hazards

his product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl (i.e. ducks and geese), feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

To obtain additional information or in case of an emergency call 770-787-1977.

[] Denotes alternate and/or optional language
 { } Denotes language that does not appear on the market label