


67572-2

6-6-2003

1/15

Please read instructions on reverse before completing form.

Form Approved OMB No. 2070-0060

United States <b>Environmental Protection Agency</b> 1200 Pennsylvania Avenue, N.W. Washington, DC 20460-0001		<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
<b>Application for Pesticide – Section I</b>			
1. Company/Product Number <b>EPA Reg. No. 67572-2</b>		2. EPA Product Manager <b>Cynthia Giles-Parker</b>	
4. Company/ Product (Name) <b>CP Multi-Purpose Fungicide Ready-to-Use</b>		3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted	
5. Name and Address of Applicant (Include ZIP Code) <b>Contract Packaging, Inc. c/o RegWest Company, LLC 30856 Rocky Road Greeley, CO 80631-9375</b>		6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	
<input type="checkbox"/> Check if this is a new address			
<b>Section – II</b>			
<input type="checkbox"/> Amendment – Explain below.		<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated <b>04/04/01</b>	
<input type="checkbox"/> Resubmission in response to Agency letter dated _____		<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification – Explain below.		<input type="checkbox"/> Other – Explain below.	
<b>NOTIFICATION</b> <b>JUN 06 2003</b>			
<b>Explanation:</b> Use additional page(s) if necessary. (For Section I and Section II.)			
<b>Submit notification of label revision primarily to add selling copy and update per PR Notice 2001-6.</b>			
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR § 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR § 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.			
<b>Section – III</b>			
1. Material This Product Will Be Packaged In:			
Child Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify): _____
<b>*Certification must be submitted</b>		If "Yes" Unit Packaging Wgt. No. Per Container	If "Yes" Packaging Wgt. No. Per Container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling Accompanying Product
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued	<input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____
<b>Section - IV</b>			
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application).			
Name <b>Kim Davis, CC, APC, Manager</b>		Title <b>Consultant/Agent</b>	Telephone No. (Include Area Code) <b>(970) 353-0611</b>
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title <b>Consultant/Agent</b>	
4. Typed Name <b>Kim Davis, CC, APC, Manager</b>		5. Date <b>May 27, 2003</b>	

2/15

U.S. Environmental Protection Agency  
1200 Pennsylvania Avenue, N.W.  
Washington, DC 20460-0001



**Notification Acknowledgement**

The following notification submission was received and accepted by the U.S. Environmental Protection Agency as of

(Date)

Identification of notification submission:

<b>Registrant Name:</b>	Contract Packaging, Inc.
<b>Product Name:</b>	CP Multi-Purpose Fungicide Ready-to-Use
<b>EPA Reg. No.:</b>	67572-2
<b>Notification Date:</b>	May 27, 2003
<b>Notification Purpose:</b>	<ol style="list-style-type: none"> <li>1. Add selling copy (taken from existing text)</li> <li>2. Update <i>Disposal</i> directions per PR Notice 2001-6</li> <li>3. Other minor changes as identified on the <i>Reference Label</i></li> </ol>

Please return this date-stamped form to:  
Kim Davis, CC, APC, Manager  
RegWest Company, LLC  
30856 Rocky Road  
Greeley, CO 80631-9375

(Fold and return postage prepaid acknowledgement – see reverse side)

30856 Rocky Road  
Greeley, CO 80631-9375  
970-353-0611  
970-353-0513 Fax  
regwestco@aol.com

3/15

# CP Multi-Purpose Fungicide Ready-To-Use

**[NEW!] Ready-to-Use Fungicide Disease Control**  
**Contains Echo™ LITE**  
**CHLOROTHALONIL FUNGICIDE**

**CONTROLS DISEASES:**  
**Leaf Spots, Black Spots, Rust[s], Blights, Fruit Rots, Mildew[s], Scab, Molds, Lawn Fungus, etc.**  
**on: Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees**

**Use on Roses**  
**For Use on Vegetables, Flowers and Shrubs**

**Ready-To-Use      For Healthier Plants      For Control of Plant Disease**

<b>Active Ingredient:</b>	
Chlorothalonil (CAS No. 1897-45-6) .....	0.087%
<b>Other Ingredients</b> .....	<b>99.913%</b>
Total .....	100.000%

**Keep Out of Reach of Children**  
**CAUTION**  
See back panel for **additional** precautionary statements.

EPA Reg. No. 67572-2                      EPA Est. No. 67572-GA-1

Manufactured [Distributed] by  
**CONTRACT PACKAGING, INC.**  
P.O. Box 1508  
Covington, GA 30015-1508

**NOTIFICATION**  
**JUN 06 2003**

**Net Contents: 1 qt. (32 fl. oz.) [24 fl. oz.] [1 gallon]**

**(Booklet Back Panel)**

Peel Back Tab for Additional Information

**Ornamental Diseases**

Cercospora, Phytophthora Die-back, Ovulinia Flower Blight, Leaf Blotch, Scab, Anthracnose, Fabraea (Entomosporium) Leaf Spot, Rhizoctonia Web Blight, Powdery Mildew, Rust, Botrytis, Alternaria Leaf Spot/Branch Rot, Septoria Leaf Spot, Black Spot, Helminthosporium Leaf Spot, Botrytis Gray Mold (foliar phase), Phytophthora Blight, Cylindrocladium Leaf Spots, Marssonina Leaf Spot and many more listed inside.

**Vegetable Disease**

Rust, Botrytis Blight (Gray Mold), Alternaria Leaf Spot, Downy Mildew, Ring Spot, Cercospora (Early) Blight, Alternaria (Late) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression), Helminthosporium Leaf Blight, Anthracnose, Scab, Purple Blotch, Botrytis Vine Rot and many more listed inside.

**Fruit Diseases**

Peach Leaf Curl, Coryneum Blight (Shothole), Cherry Leaf Spot, Brown Rot Blossom Blight, Scab.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

**Storage:** Store in a cool, dry (preferably locked) place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this empty container. Place wrap and put in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### General Precautions and Restrictions

Not for use on plants being grown for: sale or other commercial use, commercial seed production or research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in interior landscapes and ornamental gardens or parks. Also for use on fruit trees and vegetables. Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets either directly or through drift. Use of this product on home lawns is prohibited.

#### Application Instructions

#### SHAKE WELL BEFORE USING Read Entire Label and Use Strictly in Accordance With Precautionary Statements and Directions

**Personal Protective Equipment (PPE):** Wear long sleeved shirt, long pants and rubber gloves while mixing and applying this product.

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when used according to the label directions. Thorough uniform coverage of plant surfaces is essential for good disease control. Keep children and pets off treated area until dry.

#### Maximum Application Rates

For each site listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased. **Note:** AI = active ingredient; A = acre.

Site	Maximum Individual Application Rate (Lbs./AI/A)	Minimum Retreatment Interval Days	Maximum Seasonal Total Application Rate (Lbs./AI/A/Season)
Bean (snap)	2.25	7	9
Carrot	1.5	7	15
Celery	2.25	7	18
Cole crops	1.5	7	12
Corn (sweet)	1.5	7	9
Cucurbits	2.25	7	15.75
Ornamentals	1.55	7	36.5 <sup>1</sup>
Roses	1.1	7	36.4 <sup>1</sup>
Onion (dry bulb) and garlic	2.25	7	15
Onion (green), leak, shallot	2.25	7	6.7
Potato	1.125	5	11.25
Stone fruits (including cherries)	3.1	10	15.5
Tomato	2.16	7	15.1

<sup>1</sup>Seasonal maximum applies only to field-grown

#### VEGETABLE DISEASES

**Note:** Do not spray plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

PHI = the number of days to wait between the last application and harvest.

{Sq. ft. not in brackets or parentheses is for the 24 fl. oz. market container; sq. ft. in brackets is for the 32 fl. oz. container; sq. ft. in parentheses is for the 1 gallon container.}

Crop(s)	Disease(s)	Area to Cover	PHI	Comments
Beans (Green)	Rust, Botrytis Blight (Gray Mold)	28 [37.5] (112) sq. ft.	7	Begin applications during early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Alternaria Leaf Spot, Downy Mildew	56 [75] (224) sq. ft.	0	Begin after transplants are set in the field or when conditions favor disease development.
Brussels Sprouts (CA only)	Ring Spot	42 [56] (168) sq. ft.	0	Begin at time of early sprout development or when conditions favor disease development.
Carrots	Cercospora (Early) Blight, Alternaria (Late) Blight	42 [56] (168) sq. ft.	0	Begin when disease threatens and repeat as required to maintain control.
Celery	Cercospora (Early) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression)	56 [75] (224) sq. ft.	7	Begin when plants are set in the field.
Corn (Sweet)	Helminthosporium Leaf Blight, Rust	42 [56] (168) sq. ft.	14	Begin when conditions favor disease development. Repeat as needed. Do not apply to sweet corn grown for processing.
Cucumbers, Melons, Pumpkins, Squash	Anthracnose, Downy Mildew, Target Spot	42 [56] (168) sq. ft.	0	Begin when plants are in the first true leaf stage or when conditions are favorable for disease.
	Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states)	28 [37.5] (112) sq. ft.		
Cucumbers	Fruit Rot (Rhizoctonia) [suppression]			
Onion (Green), Leek, Shallot	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew [suppression]	28 [37.5] (112) sq. ft.	14	Apply at first sign of disease
Onion (Dry Bulb), Garlic	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew [suppression]	28 [37.5] (112) sq. ft.	7	Apply at first sign of disease.
Potato	Early Blight, Late Blight, Botrytis Vine Rot	56 [75] (224) sq. ft.	0	Begin when plants are 6 inches high or when disease threatens and continue as needed to maintain disease control.
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot, Botrytis Gray Mold	28 [37.5] (112) sq. ft.	0	Begin when conditions favor disease development.

**FRUIT DISEASES**

Apply a full coverage spray to the point of runoff to thoroughly cover tree canopy including undersides of leaves.

Crop(s)	Disease(s)	Comments
Apricot, Nectarine, Peach	Peach Leaf Curl, Coryneum Blight (Shothole)	Make one application at leaf fall in late autumn. Make 1 or 2 applications in mid- to late-winter before buds begin to swell. Where <i>Coryneum Blight (Shothole)</i> occurs, apply an additional application 1 to 2 weeks after petal fall or at shucksplit (tiny fruit just beginning to form) to prevent fruit infections.

6/15

Apricot, Nectarine, Peach (continued)	Brown Rot Blossom Blight	Make 1 application at popcorn (pink, red or early white bud stage when first color shows) and a second application at full bloom. If cool, moist weather continues, make an additional application at petal fall.
Apricot, Peach, Nectarine Cherry	Scab	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit. Do not apply after shucksplit or before harvest.
	Cherry Leaf Spot	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit.
	Cherry Leaf Spot (post harvest)	Make 1 application to foliage within 7 days after fruit is removed. In trees with a history of high leaf spot incidence, make a second application 10 to 14 days later.

**ORNAMENTAL DISEASES**

Spray foliage to the point of runoff. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. Only for outdoor use.

Species	Diseases Controlled	First Application
<b>Broadleaf Shrubs and Trees</b>		
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring bud break
Azalea*, Rhododendron*	Phytophthora Die-back, Ovulinia Flower Blight	New leaf emergence, early bloom
Buckeye, Horsechestnut	Leaf Blotch, Anthracnose	Spring bud break
Cherry-Laurel	Cercospora Leafspot	Petal fall
Crabapple	Scab, Cedar-Apple Rust, Sphaeropsis Leaf Spot	Spring bud break
Dogwood	Anthracnose (Discula sp.), Septoria Leaf Spot	Spring bud swell, early bloom
Euonymus	Anthracnose	Spring bud break
Firethorn (Pyracantha)	Scab	Spring bud break
Flowering Almond, Quince and Cherry Hawthorn	Monilinia Blossom/Branch Blight Rust, Fabraea Leaf Spot	Early bloom, pre-bloom
Holly	Rhizoctonia Web Blight	Warm, moist conditions
Mountain Laurel	Cercospora Leaf Spot	Spring bud break
Oak (red group only)	Taphrina Blister, Actinopelte Leaf Spot, Anthracnose	Dormant bud swell
Oregon-Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea (Entomosporium) Leaf Spot	Spring bud break
Pieris (Andromeda)	Phytophthora Die-Back	New leaf emergence
Poplar	Marssonina Leaf Spot	Spring bud break
Privet	Cercospora Leaf Spot	Prolonged wet conditions
Sycamore, Planetree	Anthracnose	Spring bud break
Viburnum	Powdery Mildew	Mid-summer
<b>Bulbs and Flowering Plants</b>		
Carnation	Alternaria Leaf Spot/Branch Rot, Botrytis	Transplant of cuttings; cool, moist conditions
Chrysanthemum/Daisy	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of cuttings
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-bloom cool, moist conditions
Gladiolus	Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea* (foliage only)	Cercospora and Septoria Leaf Spots, Rust	Early propagation
Iris	Botrytis Blossom Blight, Didymellina Leaf Spot	Cool, moist conditions
Lily	Botrytis Gray Mold (foliar phase), Phytophthora Blight	Pre-bloom
Petunia*	Botrytis Gray Mold (foliar phase), Phytophthora Blight, Botrytis Blight	Pre-bloom
Rose	Black Spot, Botrytis Blight	Spring bud break
Statice	Anthracnose, Cercospora, Alternaria, Botrytis Leaf Blight	Spring bud break
Zinnia	Powdery Mildew	First sign of disease

\*Discoloration of bloom has been noted on certain varieties when applications are made during flowering.

7/15

Foliage Plants		
Dracaena	Fusarium Leaf Spot	Pre-transplant
Leatherleaf Fern	Ascochyta Blight, Cercospora Leaf Spot, Cylindrocladium Leaf Spot, Rhizoctonia Blight	Spring bud break
Parlor Palm (Chamaedorea)	Bipolaris Leaf Spot (Helminthosporium)	Cool, moist conditions
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early propagation
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early propagation
Philodendron	Phytophthora Blight, Dactylaria Leaf Spot	Moist conditions
Syngonium	Cephalosporium Leaf Spot	Warm, moist conditions

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

**First Aid**

<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably <b>by</b> mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Immediately rinse skin with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If In Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-434-9300 <b>637-8193</b> for emergency medical treatment information.</p>	

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

**Environmental Hazards**

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flood areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

**Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.**

**To obtain additional information or in case of an emergency call 770-787-1977.**

[ ] Denotes alternate/optional language

{ } Denotes language that does not appear on the market labeling

8/15

**USE ON ROSES!**

**GARDEN TECH**

**NEW! Ready-To-Use**

# FUNGICIDE DISEASE CONTROL

**For Use on Vegetables, Flowers & Shrubs**



**Black Spots**



**Mildew**



**Rust**

**FOR CONTROL OF PLANT DISEASE**

<p><b>ACTIVE INGREDIENT:</b>          Chlorothalonil (CAS No. 1897-45-6) ..... 0.087%  <b>OTHER INGREDIENTS:</b> ..... 99.913%  <b>TOTAL:</b> ..... 100.000%</p>	<p><b>KEEP OUT OF REACH OF CHILDREN</b></p> <p><small>See Back Panel for Precautionary Statements</small></p>
--	---

**NET CONTENTS 1 qt. (32 fl. oz.) (946 ml)**

EPA REG NO. 67572-2 EPA EST NO. 67572-GA-1

file: 24704-1/label                      create: 12-21-98 es  
 folder: 24704-1/garden tech            all: 10-22-01 brn  
 colors: 4-color process, blue 072, red 186

**NOTIFICATION**  
**JUN 06 2003**



9/15

GARDEN  
FUNGICIDE  
DISEASE  
CONTROL

### Ornamental Diseases

Cercospora, Phytophthora Die-back, Ovulinia Flower Blight, Leaf Blotch, Scab, Anthracnose, Fabraea (Entomosporium) Leaf Spot, Rhizoctonia Web Blight, Powdery Mildew, Rust, Botrytis, Alternaria Leaf Spot/Branch Rot, Septoria Leaf Spot, Black Spot, Helminthosporium Leaf Spot, Botrytis Gray Mold (foliar phase), Phytophthora Blight, Cylindrocladium Leaf Spots, Marssonina Leaf Spot & many more listed inside.

### Vegetable Diseases

Rust, Botrytis Blight (Gray Mold), Alternaria Leaf Spot, Downy Mildew, Ring Spot, Cercospora (Early) Blight, Alternaria (Late) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression), Helminthosporium Leaf Blight, Anthracnose, Scab, Purple Blotch, Botrytis Vine Rot & many more listed inside.

### Fruit Diseases

Peach Leaf Curl, Coryneum Blight (Shothole), Cherry Leaf Spot, Brown Rot Blossom Blight, Scab.

Fuel Bot - 10/15/14 - 10/15/14 (10/15/14)

© 2014 Bayer CropScience  
All rights reserved. Bayer CropScience is a registered trademark of Bayer CropScience AG.  
Bayer CropScience is a registered trademark of Bayer CropScience AG.  
Bayer CropScience is a registered trademark of Bayer CropScience AG.



10/15

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

**STORAGE:** Store in a cool, dry (preferably locked) place, inaccessible to children and pets.

**DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### GENERAL PRECAUTIONS AND RESTRICTIONS

Not for use on plants being grown for sale or other commercial use, commercial seed production or research purposes.

For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes and ornamental gardens or parks. Also for use on fruit trees and vegetables. Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets either directly or through drift. Use of this product on home lawns is prohibited.

#### Application Instructions

#### SHAKE WELL BEFORE USING

#### READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

**Personal Protective Equipment (PPE):** Wear long sleeved shirt, long pants, and rubber gloves while mixing and applying this product.

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when it is used according to the label directions. Thorough uniform coverage of plant surfaces is essential for good disease control. Keep children and pets off treated area until dry.

#### Maximum Application Rates

For each site listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased. Note: AI = active ingredient; A = acre.

Site	Maximum Individual Application Rate (Lbs./AI/A)	Minimum Retreatment Interval Days	Maximum Seasonal Total Application Rate (Lbs./AI/A/Season)
Bean (snap)	2.25	7	9

(continued)

NOTES

11/15

**FIRST AID**

**IF SWALLOWED:** • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

**IF INHALED:** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING:** • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

**IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-434-9300 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

**Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.**

GardenTech is a trademark for Gulfstream Home & Garden, Inc.

(040401EPA/PR2001-6)

**Maximum Application Rates (continued)**

Site	Maximum Individual Application Rate (Lbs./A/A)	Maximum Retreatment Interval Days	Maximum Seasonal Total Application Rate (Lbs./A/A/Season)
Carrot	1.5	7	15
Celery	2.25	7	18
Cole crops	1.5	7	12
Corn (sweet)	1.5	7	9
Cucurbits	2.25	7	15.75
Ornamentals	1.55	7	36.5 <sup>1</sup>
Roses	1.1	7	36.4 <sup>1</sup>
Onion (dry bulb) and garlic	2.25	7	15
Onion (green), leek, shallot	2.25	7	6.7
Potato	1.125	5	11.25
Stone fruits (including cherries)	3.1	10	15.5
Tomato	2.16	7	15.1

<sup>1</sup>Seasonal maximum applies only to field-grown

**VEGETABLE DISEASES**

Note: Do not spray plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

PHI = the number of days to wait between the last application and harvest.

Crop(s)	Disease(s)	Area to Cover	PHI	Comments
Beans (Green)	Rust, Botrytis Blight (Gray Mold)	37.5 sq. ft.	7	Begin applications during early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary.

(continued)

12/15

### FRUIT DISEASES

Apply a full coverage spray to the point of runoff to thoroughly cover tree canopy including undersides of leaves.

Crop(s)	Disease(s)	Comments
Apricot, Nectarine, Peach	Peach Leaf Curl, Coryneum Blight (Shothole)	Make one application at leaf fall in late autumn. Make 1 or 2 applications in mid- to late-winter before buds begin to swell. Where <i>Coryneum Blight (Shothole)</i> occurs, apply an additional application 1 to 2 weeks after petal fall or at shucksplit (tiny fruit just beginning to form) to prevent fruit infections.
	Brown Rot Blossom Blight	Make 1 application at popcorn (pink, red or early white bud stage when first color shows) and a second application at full bloom. If cool, moist weather continues, make an additional application at petal fall.
Apricot, Peach, Nectarine, Cherry	Scab	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit. Do not apply after shucksplit or before harvest.
	Cherry Leaf Spot	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit.
	Cherry Leaf Spot (post harvest)	Make 1 application to foliage within 7 days after fruit is removed. In trees with a history of high leaf spot incidence, make a second application 10 to 14 days later.

### ORNAMENTAL DISEASES

Spray foliage to the point of runoff. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. Only for outdoor use.

Species	Diseases Controlled	First Application
<b>BROADLEAF SHRUBS AND TREES</b>		
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring Bud Break
Azalea*	Phytophthora Die-back, Ovipollinia Flower Blight	New Leaf Emergence, Early Bloom
Buckeye, Horsechestnut	Leaf Blotch, Anthracnose	Spring Bud Break
Cherry-Laurel	Cercospora Leafspot	Petal Fall
Crabapple	Scab, Cedar-Apple Rust, Sphaeropsis Leaf Spot	Spring Bud Break
Dogwood	Anthracnose (Discula Sp.), Septoria Leaf Spot	Spring Bud Swell, Early Bloom
Euonymus	Anthracnose	Spring Bud Break
Firethorn (Pyracantha)	Scab	Spring Bud Break
Flowering Almond, Quince and Cherry Hawthorn	Monilinia Blossom/Branch Blight, Rust, Fabraea Leaf Spot	Early Bloom, Pre-Bloom
Holly	Rhizoctonia Web Blight	Warm, Moist Conditions
Mountain Laurel	Cercospora Leaf Spot	Spring Bud Break
Oak (red group only)	Taphrina Blister, Actinopelte Leaf Spot, Anthracnose	Dormant Bud Swell
Oregon-Grape (Mahonia)	Rust	Spring Bud Break
Photinia	Fabraea (Entomosporium) Leaf Spot	Spring Bud Break
Pieris (Andromeda)	Phytophthora Die-Back	New Leaf Emergence
Poplar	Marssonina Leaf Spot	Spring Bud Break

(continued)

**VEGETABLE DISEASES (continued)**

Crop(s)	Disease(s)	Area to Cover	PH	Comments
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Alternaria Leaf Spot, Downy Mildew	75 sq. ft.	0	Begin after transplants are set in the field or when conditions favor disease development.
Brussels Sprouts (CA only)	Ring Spot	56 sq. ft.	0	Begin at time of early sprout development or when conditions favor disease development.
Carrots	Cercospora (Early) Blight, Alternaria (Late) Blight	56 sq. ft.	0	Begin when disease threatens and repeat as required to maintain control.
Celery	Cercospora (Early) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression)	75 sq. ft.	7	Begin when plants are set in the field.
Corn (Sweet)	Helminthosporium Leaf Blight, Rust	56 sq. ft.	14	Begin when conditions favor disease development. Repeat as needed. Do not apply to sweet corn grown for processing.
Cucumbers, Melons, Pumpkins, Squash	Anthracnose, Downy Mildew, Target Spot	56 sq. ft.	0	Begin when plants are in the first true leaf stage or when conditions are favorable for disease.
	Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states)	37.5 sq. ft.		

(continued)

**FOLIAGE PLANTS**

Species	Diseases Controlled	First Application
Dracaena	Fusarium Leaf Spot	Pre-Transplant
Leatherleaf Fern	Ascochyta Blight, Cercospora Leaf Spot, Cylindrocladium Leaf Spot, Rhizoctonia Blight	Spring Bud Break
Parlor Palm (Chamaedorea)	Bipolaris Leaf Spot (Helminthosporium)	Cool, Moist Conditions
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early Propagation
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early Propagation
Philodendron	Phytophthora Blight, Dactylaria Leaf Spot	Moist Conditions
Syngonium	Cephalosporium Leaf Spot	Warm, Moist Conditions

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

14/15

Species	Diseases Controlled	First Application
<b>BROADLEAF SHRUBS AND TREES (continued)</b>		
Privet	Cercospora Leaf Spot	Prolonged Wet Conditions
Sycamore, Planetree	Anthracoese	Spring Bud Break
Viburnum	Powdery Mildew	Mid-Summer
<b>BULBS AND FLOWERING PLANTS</b>		
Carnation	Alternaria Leaf Spot/Branch Rot, Botrytis	Transplant of Cuttings, Cool, Moist Conditions
Chrysanthemum/Daisy	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of Cuttings
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-Bloom Cool, Moist Conditions
Gladiolus	Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot	Early Propagation
Hollyhock	Rust	Early Seedling Stage
Hydrangea* (foliage only)	Cercospora & Septoria Leaf Spots, Rust	Early Propagation
Iris	Botrytis Blossom Blight, Dicyemellina Leaf Spot	Cool, Moist Conditions
Lily	Botrytis Gray Mold (foliar phase), Phytophthora Blight	Pre-Bloom
Petunia*	Botrytis Gray Mold (foliar phase), Phytophthora Blight, Botrytis Blight	Pre-Bloom
Rose	Black Spot, Botrytis Blight	Spring Bud Break
Statice	Anthracoese, Cercospora, Alternaria, Botrytis Leaf Blight	Spring Bud Break
Zinnia	Powdery Mildew	First Sign of Disease

\*DISCOLORATION OF BLOOM HAS BEEN NOTED ON CERTAIN VARIETIES WHEN APPLICATIONS ARE MADE DURING FLOWERING.

**VEGETABLE DISEASES (continued)**

Crop(s)	Disease(s)	Area to Cover	P/W	Comments
Cucumbers	Fruit Rot (Rhizoctonia) [suppression]	37.5 sq. ft.	0	Begin when plants are in the first true leaf stage or when conditions are favorable for disease.
Onion (Green), Leak, Shallot	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	37.5 sq. ft.	14	Apply at first sign of disease.
Onion (Dry Bulb), Garlic	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	37.5 sq. ft.	7	Apply at first sign of disease.
Potato	Early Blight, Late Blight, Botrytis Vine Rot	75 sq. ft.	0	Begin when plants are 6 inches high or when disease threatens and continue as needed to maintain disease control.
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot, Botrytis Gray Mold	37.5 sq. ft.	0	Begin when conditions favor disease development.



May 27, 2003

Ms. Sherada Hobgood  
Document Processing Desk – **NOTIF**  
Office of Pesticide Programs – **7504C**  
**U.S. Environmental Protection Agency**  
1921 Jefferson Davis Highway  
Crystal Mall #2, Room 266A  
Arlington, VA 22202-4501

Dear Ms. Hobgood:

**Subject:** Contract Packaging, Inc.  
CP Multi-Purpose Fungicide Ready-to-Use  
EPA Reg. No. 67572-2

On behalf of its client **Contract Packaging, Inc.**, RegWest Company, LLC is submitting the attached material as notification of label revision for subject product. RegWest Company, LLC will act as sole agent in this endeavor.

Attached are the following:

1. Application for Pesticide: Other
2. Notification Acknowledgement
3. One **Reference Label** (illustrates all label revisions)
4. Two copies of market labeling (incorporating the revisions)

We are primarily adding selling copy (taken from existing text) and updating the label PR Notice 2001-6.

We are using market labeling with our additional brand name, *GardenTech Ready-to-Use Fungicide Disease Control*. We understand the "NEW!" can appear only for 6 months after introduction of the product into the marketplace.

Please contact me at (970) 353-0611 or at regwestco@aol.com if you have any questions concerning this submission or require additional information.

Regards,  
**RegWest Company, LLC**



Kim Davis, CC, APC, Manager  
Consultant/Agent

Attachments

pc: CPI – Hacker, S.

30856 Rocky Road  
Greeley, CO 80631-9375  
970-353-0611  
970-353-0611 Fax  
regwestco@aol.com