Section – III							
1. Material This Product Will Be Packaged In:							
Child Resistant Packaging	Unit Packaging			Water Soluble Pack	aging	2. Type of Co	ontainer
☐ Yes*	Yes			Yes		☐ Metal	
ĺ □ No	□ No			□ No		☐ Plastic	
*Certification must	If "Yes"	No. Po	<u>er</u>	If "Yes"	No. Per	—— ☐ Glass	
be submitted	Unit Packaging Wgt.	Conta	* .	Backgring Mat Container D Pape		r   🗅 Paper	
					1	Other (S	
Location of Net Contents	Information	4. Size(s)	Retail	Container		5. Location of Labe	el Directions
<b>  -</b>	<b>–</b>					On Label	
Label	☐ Container					On Labeling A	ccompanying Product
6. Manner in Which Label is	Affixed to Product	C	<b>J</b> Lith	nograph	Stenci	led	
ļ		[	<b>⊐</b> Pap	per glued	Other		
				ection - IV	. "		
1. Contact Point (Comp	lete items directly be	low for idea	ntificat	tion of Individual to	be contacte	d, if necessary, to p	rocess this application).
Name		Title				Telephone No. (Incl	ude Area Code)
Kim Davis CC Al	PC Manager		Cor	nsultant/Agent		(970) 353-0611	/ / kim@regwest.com
Kim Davis, CC, APC, Manager Certifica						(370) 333-001	6. Date Application
				Received			
accurate and complete. I acknowledge that any knowingly false or misleading statement may be (Stamped)							
punishable by fine or imprisonment or both under applicable law.							
			3. Title				
			Consultant/Agent			; !	
4. Typed Name			5. Dat	te			
Kim Davis CC APC Manager				Decemb	ner 8 200	14	

# **CP Multi-Purpose Fungicide Ready-To-Use**

[NEW!] Ready-to-Use Fungicide Disease Control Contains Echo™ LITE CHLOROTHALONIL FUNGICIDE NOTIFICATION
JAN 07 2005

Controls/Stops and Prevents over 70 Diseases on Flowers, Vegetables, Shrubs, Fruit and Shade Trees

# **CONTROLS DISEASES:**

Leaf Spots, Black Spots, Rust[s], Blights, Fruit Rots, Mildew[s], Scab, Molds, Lawn Fungus, etc. on: Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees

Use on Roses
For Use on Vegetables, Flowers and Shrubs

Ready-To-Use

For Healthier Plants

For Control of Plant Disease

**Active Ingredient:** 

 Chlorothalonil (CAS No. 1897-45-6)
 0.087%

 Other Ingredients
 99.913%

 Total
 100.000%

Keep Out of Reach of Children CAUTION

See back panel for additional precautionary statements.

EPA Reg. No. 67572-2

EPA Est. No. 67572-GA

Manufactured [Distributed] by

P.O. Box 1508
Covington, GA 30015-1508

Contents: 1 at (32 fl oz ) (956 m

Net Contents: 1 qt. (32 fl. oz.) (956 ml) [24 fl. oz.] [1 gallon]

{Booklet}

Peel Back Tab for Additional Information

# **Ornamental Diseases**

Cercospora, Phytophthora Die-back, Ovulinia Flower Blight, Leaf Blotch, Scab, Anthracnose, Fabraea (Entomosporium) Leaf Spot, Rhizoctonia Web Blight, Powdery Mildew, Rust, Botrytis, Alternaria Leaf Spot/Branch Rot, Septoria Leaf Spot, Black Spot, Helminthosporium Leaf Spot, Botrytis Gray Mold (foliar phase), Phytophthora Blight, Cylindrocladium Leaf Spots, Marssonina Leaf Spot and many more listed inside.

### Vegetable Disease

Rust, Botrytis Blight (Gray Mold), Alternaria Leaf Spot, Downy Mildew, Ring Spot, Cercospora (Early) Blight, Alternaria ate) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression), Helminthosporium Leaf Blight, Anthracnose, ab, Purple Blotch, Botrytis Vine Rot and many more listed inside.

## Fruit Diseases

Peach Leaf Curl, Coryneum Blight (Shothole), Cherry Leaf Spot, Brown Rot Blossom Blight, Scab.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

# Storage and Disposal

**Storage:** Store in a cool, dry (preferably locked) place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### **General Precautions and Restrictions**

Not for use on plants being grown for sale or other commercial use, commercial seed production or research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in interior plantscapes and ornamental gardens or parks. Also for use on fruit trees and vegetables. Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets either directly or through drift. Use of this product on home lawns is prohibited.

# **Application Instructions**

# SHAKE WELL BEFORE USING Read Entire Label and Use Strictly in Accordance With Precautionary Statements and Directions

**Personal Protective Equipment (PPE):** Wear long sleeved shirt, long pants and rubber gloves while mixing and applying this product.

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when used according to the label directions. Thorough uniform coverage of plant surfaces is essential for good disease control. Keep children and pets off treated area until dry.

# **Maximum Application Rates**

or each site listed below, the listed maximum individual and seasonal application rates must not be exceeded and the asted minimum retreatment intervals must not be decreased. **Note:** Al = active ingredient; A = acre.

Site	Maximum Individual Application Rate (Lbs./Al/A)	Minimum Retreatment Interval Days	Maximum Seasonal Total Application Rate (Lbs./Al/A/Season)
Bean (snap)	2.25	7	9
Carrot	1.5	7	15
Celery	2.25	7	18
Cole crops	1.5	7	12
Corn (sweet)	1.5	7	9
Cucurbits	2.25	7	15.75
Ornamentals	1.55	7	36.5
Roses	1.1	7	36.41
Onion (dry bulb) and garlic	2.25	7	15
Onion (green), leak, shallot	2.25	7	6.7
Potato	1.125	5	11.25
Stone fruits (including cherries)	3.1	10	15.5
Tomato	2.16	7	15.1

Seasonal maximum applies only to field-grown

### **VEGETABLE DISEASES**

**Note:** Do not spray plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

PHI = the number of days to wait between the last application and harvest.

Sq. ft. not in brackets or parentheses is for the 24 fl. oz. market container; sq. ft. in brackets is for the 32 fl. oz. container; 4. ft. in parentheses is for the 1 gallon container.

67572-2 **New Label**: Page 3 of 6 04/04/01 SAL + 05/27/03 (05/27/03) + 12/08/04 NOTIFs

Crop(s)	Disease(s)	Area to Cover	PHI	Comments
Beans (Green)	Rust, Botrytis Blight (Gray Mold)	28 [37.5] (112) sq. ft.	7	Begin applications during early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Alternaria Leaf Spot, Downy Mildew	56 [75] (224) sq. ft.	0	Begin after transplants are set in the field or when conditions favor disease development.
Brussels Sprouts (CA only)	Ring Spot	42 [56] (168) sq. ft.	0	Begin at time of early sprout development or when conditions favor disease development.
Carrots	Cercospora (Early) Blight, Alternaria (Late) Blight	42 [56] (168) sq. ft.	0	Begin when disease threatens and repeat as required to maintain control.
Celery	Cercospora (Early) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression)	56 [75] (224) sq. ft.	7	Begin when plants are set in the field.
Corn (Sweet)	Helminthosporium Leaf Blight, Rust	42 [56] (168) sq. ft.	14	Begin when conditions favor disease development. Repeat as needed. Do not apply to sweet corn grown for processing.
Cucumbers, Melons, Pumpkins, Squash	Anthracnose, Downy Mildew, Target Spot Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states)	42 [56] (168) sq. ft. 28 [37.5] (112) sq. ft.	0	Begin when plants are in the first true leaf stage or when conditions are favorable for disease.
Cucumbers	Fruit Rot (Rhizoctonia) [suppression]			
Onion (Green), Leek, Shallot	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	28 [37.5] (112) sq. ft.	14	Apply at first sign of disease
Onion (Dry Bulb), Garlic	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	28 [37.5] (112) sq. ft.	7	Apply at first sign of disease.
Potato	Early Blight, Late Blight, Botrytis Vine Rot	56 [75] (224) sq. ft.	0	Begin when plants are 6 inches high or when disease threatens and continue as needed to maintain disease control.
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot, Botrytis Gray Mold	28 [37.5] (112) sq. ft.	0	Begin when conditions favor disease development.

# **FRUIT DISEASES**

Apply a full coverage spray to the point of runoff to thoroughly cover tree canopy including undersides of leaves.

Crop(s)	Disease(s)	Comments
Apricot, Nectarine, Peach	Peach Leaf Curl, Coryneum Blight (Shothole)	Make one application at leaf fall in late autumn. Make 1 or 2 applications in mid- to late-winter before buds begin to swell. Where Coryneum Blight (Shothole) occurs, apply an additional application 1 to 2 weeks after petal fall or at shucksplit (tiny fruit just beginning to form) to prevent fruit infections.

67572-2 **New Label**: Page 4 of 6 04/04/01 SAL + 05/27/03 (05/27/03) + 12/08/04 NOTIFs

Crop(s)	Disease(s)	Comments
Apricot, Nectarine, Peach (continued)	Brown Rot Blossom Blight	Make 1 application at popcorn (pink, red or early white bud stage when first color shows) and a second application at full bloom. If cool, moist weather continues, make an additional application at petal fall.
Apricot, Peach, Nectarine, Cherry	Scab	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit. Do not apply after shucksplit or before harvest.
	Cherry Leaf Spot	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit.
	Cherry Leaf Spot (post harvest)	Make 1 application to foliage within 7 days after fruit is removed. In trees with a history of high leaf spot incidence, make a second application 10 to 14 days later.

# **ORNAMENTAL DISEASES**

Spray foliage to the point of runoff. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. Only for outdoor use.

Species	Diseases Controlled	First Application		
Broadleaf Shrubs and Trees				
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring bud break		
Azalea*, Rhododendron*	Phytophthora Die-back, Ovulinia Flower Blight	New leaf emergence, early		
	<u> </u>	bloom		
uckeye, Horsechestnut	Leaf Blotch, Anthracnose	Spring bud break		
_herry-Laurel	Cercospora Leafspot	Petal fall		
Crabapple	Scab, Cedar-Apple Rust, Sphaeropsis Leaf Spot	Spring bud break		
Dogwood	Anthracnose (Discula sp.), Septoria Leaf Spot	Spring bud swell, early bloom		
Euonymus	Anthracnose	Spring bud break		
Firethorn (Pyracantha)	Scab	Spring bud break		
Flowering Almond, Quince	Monilinia Blossom/Branch Blight Rust, Fabraea Leaf Spot	Early bloom, pre-bloom		
and Cherry Hawthorn				
Holly	Rhizoctonia Web Blight	Warm, moist conditions		
Mountain Laurel	Cercospora Leaf Spot	Spring bud break		
Oak (red group only)	Taphrina Blister, Actinopelte Leaf Spot, Anthracnose	Dormant bud swell		
Oregon-Grape (Mahonia)	Rust	Spring bud break		
Photinia	Fabraea (Entomosporium) Leaf Spot	Spring bud break		
Pieris (Andromeda)	Phytophthora Die-Back	New leaf emergence		
Poplar	Marssonina Leaf Spot	Spring bud break		
Privet	Cercospora Leaf Spot	Prolonged wet conditions		
Sycamore, Planetree	Anthracnose	Spring bud break		
Viburnum	Powdery Mildew	Mid-summer		
	Bulbs and Flowering Plants			
Carnation	Alternaria Leaf Spot/Branch Rot, Botrytis	Transplant of cuttings; cool,		
	' ' '	moist conditions		
Chrysanthemum/Daisy	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of cuttings		
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-bloom cool, moist conditions		
Gladiolus	Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot	Early propagation		
Hollyhock	Rust	Early seedling stage		
Hydrangea* (foliage only)	Cercospora and Septoria Leaf Spots, Rust	Early propagation		
Iris	Botrytis Blossom Blight, Didymellina Leaf Spot	Cool, moist conditions		
' ily	Botrytis Gray Mold (foliar phase), Phytophthora Blight	Pre-bloom		
etunia*	Botrytis Gray Mold (foliar phase), Phytophthora Blight, Botrytis Blight Pre-bloom			
Rose	Black Spot, Botrytis Blight	Spring bud break		
Statice	Anthracnose, Cercospora, Alternaria, Botrytis Leaf Blight	Spring bud break		
Zinnia Powdery Mildew First sign of disease				

Foliage Plants			
racaena	Fusarium Leaf Spot	Pre-transplant	
Leatherleaf Fern	Ascochyta Blight, Cercospora Leaf Spot, Cylindrocladium Leaf Spot, Rhizoctonia Blight	Spring bud break	
Parlor Palm (Chamaedorea)	Bipolaris Leaf Spot (Helminthosporium)	Cool, moist conditions	
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early propagation	
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early propagation	
Philodendron	Phytophthora Blight, Dactylaria Leaf Spot	Moist conditions	
Syngonium	Cephalosporium Leaf Spot	Warm, moist conditions	

### PRECAUTIONARY STATEMENTS

# **Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

### First Aid

	Tilat Ald
If Swallowed	Immediately call a poison control center or doctor for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by the poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
If Inhaled	Move person to fresh air.
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
•	Call a poison control center or doctor for further treatment advice.
lf on Skin	Take off contaminated clothing.
or Clothing	Immediately rinse skin with plenty of water for 15-20 minutes.
_	Call a poison control center or doctor for treatment advice.
If In Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
•	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
	ct container or label with you when calling a poison control center or doctor or going for treatment. You ot 1-800-420-9341 for emergency medical treatment information.

(Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.)

### **Environmental Hazards**

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

{Code}

If you have a medical emergency, need consultation or diagnosis, therapy and analysis please call (800) 420-9347 If you have questions relating to product use or application, please call (800) 969-7200

67572-2 **New Label**: Page 6 of 6 04/04/01 SAL + 05/27/03 (05/27/03) + 12/08/04 NOTIFs

If you have a product spill or transportation emergency call (800) 424-9300 For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200

[ ] Denotes alternate/optional language { } Denotes language that does not appear on the market labeling



December 8, 2004

Ms. Sherada Hobgood
Document Processing Desk – NOTIF
Office of Pesticide Programs – 7504C
U.S. Environmental Protection Agency
1801 South Bell Street
Crystal Mall #2, Room 266A
Arlington, VA 22202-4501

Dear Ms. Hobgood:

Subject: Contract Packaging, Inc.

CP Multi-Purpose Fungicide Ready-to-Use

EPA Reg. No. 67572-2

n behalf of its client *Contract Packaging, Inc.*, RegWest Company, LLC is submitting the attached materials as notification of label revision for subject product. RegWest Company will act as sole agent in this endeavor.

Attached are the following:

- 1. Application for Pesticide: Other
- 2. One Reference Label (illustrates all changes)
- 3. Three **New Labels** (incorporating all changes)

Please refer to the **Reference Label** which illustrates all the changes. We are primarily adding selling copy consumer contact numbers. The additional selling copy on the front panel is substantially similar to a statement which was accepted by the EPA for the chlorothalonil concentrate product, EPA Reg. No. 67572-82.

Please contact me at (970) 353-0611 or kim@regwest.com if you have any questions concerning this submission or require additional information.

Regards,

RegWest Company, LLC

Kim Davis, CC, APC, Manager

Onsultant/Agent

**Attachments** 

pc: CPI - Hacker, S.

30856 Rocky Road Creeley, CO 80631-9375 970-353-0311

070 050 0113 0

970-353-0313 Fax

Kim@regwest.com