

67572-2

06-16-2010

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mrs. Kim Davis  
Consultant/Agent  
Contract Packaging, Inc.  
c/o RegWest Company, LLC  
8203 West 20<sup>th</sup> St., Suite A  
Greeley, CO 80634-4696

JUN 16 2010

RE: Notification to Add the California "Prop 65" Verbiage  
EPA Registration Number: 67572-2  
Date of Submission: May 13, 2010

Dear Mrs. Davis:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, May 13, 2010, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs

|  |   |  |
|--|---|--|
| United States<br><b>Environmental Protection Agency</b><br>1200 Pennsylvania Avenue, N.W.<br>Washington, DC 20460-0001 | <input type="checkbox"/> <b>Registration</b><br><input type="checkbox"/> <b>Amendment</b><br><input checked="" type="checkbox"/> <b>Other</b> | OPP Identifier Number<br><br><h2 style="text-align: center;">Notification</h2> |
|--|---|--|

**Application for Pesticide – Section I**

|  |   |  |
|--|---|--|
| 1. Company/Product Number<br><p style="text-align: center;"><b>EPA Reg. No. 67572-2</b></p>  | 2. EPA Product Manager<br><p style="text-align: center;"><b>Tony Kish</b></p>   | 3. Proposed Classification<br><input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/ Product (Name)<br><p style="text-align: center;"><b>CP Multi-Purpose Fungicide Ready-to-Use</b></p>  | PM #<br><p style="text-align: center;"><b>10</b></p>  |  |
| 5. Name and Address of Applicant (Include ZIP Code)<br><b>Contract Packaging, Inc.</b><br><b>c/o RegWest Company, LLC</b><br><b>8203 West 20<sup>th</sup> St., Suite A</b><br><b>Greeley, CO 80634-4696</b><br><input type="checkbox"/> Check if this is a new address | 6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to:<br><br>EPA Reg. No. _____<br><br>Product Name _____ |  |

**Section – II**

|  |   |
|--|---|
| <input type="checkbox"/> Amendment – Explain below.<br><input type="checkbox"/> Resubmission in response to Agency letter dated _____<br><input checked="" type="checkbox"/> Notification – Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____<br><input type="checkbox"/> "Me Too" Application.<br><input type="checkbox"/> Other – Explain below. |
|--|---|

NOTIFICATION  
JUN 16 2010

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)

Submit label revision notification per PR Notice 98-10 to add the California "Prop 65" verbiage

*This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR § 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR § 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.*

**Section – III**

|   |   |  |                         |   |   |
|---|---|--|-------------------------|---|---|
| 1. Material This Product Will Be Packaged In:   |   |  |                         | 2. Type of Container  |   |
| Child Resistant Packaging<br><input type="checkbox"/> Yes*<br><input type="checkbox"/> No   | Unit Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No | Water Soluble Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No |                         | <input type="checkbox"/> Metal  | <input type="checkbox"/> Plastic                |
|   |   |  |                         | <input type="checkbox"/> Glass  | <input type="checkbox"/> Paper                  |
| <b>*Certification must be submitted</b>   | If "Yes" Unit Packaging Wgt.  | No. Per Container  | If "Yes" Packaging Wgt. | No. Per Container   | <input type="checkbox"/> Other (Specify): _____ |
| 3. Location of Net Contents Information<br><input type="checkbox"/> Label <input type="checkbox"/> Container  |   | 4. Size(s) Retail Container  |                         | 5. Location of Label Directions<br><input type="checkbox"/> On Label<br><input type="checkbox"/> On Labeling Accompanying Product |   |
| 6. Manner in Which Label is Affixed to Product<br><input type="checkbox"/> Lithograph <input type="checkbox"/> Stenciled<br><input type="checkbox"/> Paper glued <input type="checkbox"/> Other _____ |   |  |                         |   |   |

**Section - IV**

1. **Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application).**

|   |   |  |
|---|---|--|
| Name<br><p style="text-align: center;"><b>Kim Davis</b></p> | Title<br><p style="text-align: center;"><b>Consultant/Agent</b></p> | Telephone No. (Include Area Code)<br><p style="text-align: center;"><b>970.353.0611</b><br/><b>kim@regwest.com</b></p> |
|---|---|--|

|  |  |  |  |
|--|--|--|--|
| <b>Certification</b>   |  |  | 6. Date Application Received<br><b>(Stamped)</b> |
| I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. |  |  |  |
| 2. Signature<br>   | 3. Title<br><p style="text-align: center;"><b>Consultant/Agent</b></p> |  |  |
| 4. Typed Name<br><p style="text-align: center;"><b>Kim Davis</b></p>   | 5. Date<br><p style="text-align: center;"><b>May 13, 2010</b></p>      |  |  |



May 13, 2010

Mr. Tony Kish, Product Manager 22  
Document Processing Desk – **NOTIF**  
Office of Pesticide Programs – **7504P**  
**U.S. Environmental Protection Agency**  
One Potomac Yard (South Building), Room S-4900  
2777 South Crystal Drive  
Arlington, VA 22202

Dear Mr. Kish:

**Subject:** Contract Packaging, Inc.  
CP Multi-Purpose Fungicide Read-to-Use  
EPA Reg. No. 67572-2

On behalf of **Contract Packaging, Inc.**, RegWest Company, LLC is submitting the attached materials as a notification of label revision for subject product. RegWest will act as sole agent in this endeavor.

Attached are the following:

1. Application for Pesticide: Other (Notification)
2. One *Reference Label* (illustrating the revision)
3. Two *New Labels* (incorporating the revision)

We are adding the California "Prop 65" verbiage to the label. No other changes are being made at this time.

Market labeling is not included in this submission as this product is marketed by supplemental distributors, and not by Contract Packaging.


Please contact me at (970) 353-0611 or kim@regwest.com if you have any questions concerning this submission or require additional information.

Regards,  
RegWest Company, LLC

Kim Davis,  
Consultant/Agent

Attachments

cc: CPI – Hacker, S.



4  
11

# CP Multi-Purpose Fungicide Ready-To-Use

**NOTIFICATION**

Ready-to-Use Fungicide Disease Control  
Contains Echo™ LITE  
CHLOROTHALONIL FUNGICIDE

**JUN 16 2010**

Controls/Stops and Prevents over 70 Diseases on Flowers, Vegetables, Shrubs, Fruit and Shade Trees

**CONTROLS DISEASES:**

Leaf Spots, Black Spots, Rust[s], Blights, Fruit Rots, Mildew[s], Scab, Molds, Lawn Fungus, etc.  
on: Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees

Use on Roses

For Use on Vegetables, Flowers and Shrubs

Ready-To-Use      For Healthier Plants      For Control of Plant Disease

**Active Ingredient:**

|  |                 |
|--|-----------------|
| Chlorothalonil (CAS No. 1897-45-6) ..... | 0.087%          |
| <b>Other Ingredients</b> .....           | <b>99.913%</b>  |
| <b>Total</b> .....                       | <b>100.000%</b> |

**Keep Out of Reach of Children**  
**CAUTION**  
See back panel [booklet] for additional precautionary statements.

EPA Reg. No. 67572-2

EPA Est. No. 67572-GA-1

Manufactured [Distributed] by  
**Contract Packaging, Inc.**  
P.O. Box 1508  
Covington, GA 30015-1508


**Net Contents: 1 qt. (32 fl. oz.) (946 ml)**  
**[24 fl. oz.] [1 gallon]**

{Booklet}

Peel Back Tab for Additional Information

**Ornamental Diseases**

Cercospora, Phytophthora Die-back, Ovulinia Flower Blight, Leaf Blotch, Scab, Anthracnose, Fabraea (Entomosporium) Leaf Spot, Rhizoctonia Web Blight, Powdery Mildew, Rust, Botrytis, Alternaria Leaf Spot, Branch Rot, Septoria Leaf Spot, Black Spot, Helminthosporium Leaf Spot, Botrytis Gray Mold (foliar phase), Phytophthora Blight, Cylindrocladium Leaf Spots, Marssonina Leaf Spot and many more listed inside.



5  
11

### Vegetable Diseases

Rust, Botrytis Blight (Gray Mold), Alternaria Leaf Spot, Downy Mildew, Ring Spot, Cercospora (Early) Blight, Alternaria (Late) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression), Helminthosporium Leaf Blight, Anthracnose, Scab, Purple Blotch, Botrytis Vine Rot and many more listed inside.

### Fruit Diseases

Peach Leaf Curl, Coryneum Blight (Shothole), Cherry Leaf Spot, Brown Rot Blossom Blight, Scab.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### General Precautions and Restrictions

Not for use on plants being grown for sale or other commercial use, commercial seed production or research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in interior landscapes and ornamental gardens or parks. Also for use on fruit trees and vegetables. Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets either directly or through drift. Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle or high schools), campgrounds, churches and theme parks.

### Application Instructions

#### SHAKE WELL BEFORE USING

Read Entire Label and Use Strictly in Accordance  
 With Precautionary Statements and Directions

**Personal Protective Equipment (PPE):** Wear long sleeved shirt, long pants and chemical-resistant gloves made of any waterproof material while mixing and applying this product.

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when used according to the label directions. Thorough uniform coverage of plant surfaces is essential for good disease control. Keep children and pets off treated area until dry.

### Maximum Application Rates

For each site listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased. **Note:** AI = active ingredient; A = acre.

| Site                        | Maximum Individual Application Rate (Lbs./AI/A) | Minimum Retreatment Interval Days | Maximum Seasonal Total Application Rate (Lbs./AI/A/Season) |
|-----------------------------|---|-----------------------------------|--|
| Bean (snap)                 | 2.25  | 7                                 | 9  |
| Carrot                      | 1.5   | 7                                 | 15   |
| Celery                      | 2.25  | 7                                 | 18   |
| Cole crops                  | 1.5   | 7                                 | 12   |
| Corn (sweet)                | 1.5   | 7                                 | 9  |
| Cucurbits                   | 2.25  | 7                                 | 15.75  |
| Ornamentals                 | 1.55  | 7                                 | 36.5 <sup>1</sup>  |
| Roses                       | 1.1   | 7                                 | 36.4 <sup>1</sup>  |
| Onion (dry bulb) and garlic | 2.25  | 7                                 | 15   |

| Site                              | Maximum Individual Application Rate (Lbs./AI/A) | Minimum Retreatment Interval Days | Maximum Seasonal Total Application Rate (Lbs./AI/A/Season) |
|-----------------------------------|---|-----------------------------------|--|
| Onion (green), leak, shallot      | 2.25  | 7                                 | 6.7  |
| Potato                            | 1.125   | 5                                 | 11.25  |
| Stone fruits (including cherries) | 3.1   | 10                                | 15.5   |
| Tomato                            | 2.16  | 7                                 | 15.1   |

<sup>1</sup>Seasonal maximum applies only to field-grown

### VEGETABLE DISEASES

**Note:** Do not spray plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

PHI = the number of days to wait between the last application and harvest.

{Sq. ft. not in brackets or parentheses is for the 24 fl. oz. market container; sq. ft. in brackets is for the 32 fl. oz. container; sq. ft. in parentheses is for the 1 gallon container.}

| Crop(s)  | Disease(s)   | Area to Cover           | PHI | Comments   |
|--|--|-------------------------|-----|--|
| Beans (Green)                                    | Rust, Botrytis Blight (Gray Mold)  | 28 [37.5] (112) sq. ft. | 7   | Begin applications during early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary. |
| Broccoli, Brussels Sprouts, Cabbage, Cauliflower | Alternaria Leaf Spot, Downy Mildew   | 56 [75] (224) sq. ft.   | 0   | Begin after transplants are set in the field or when conditions favor disease development.                               |
| Brussels Sprouts (CA only)                       | Ring Spot  | 42 [56] (168) sq. ft.   | 0   | Begin at time of early sprout development or when conditions favor disease development.                                  |
| Carrots  | Cercospora (Early) Blight, Alternaria (Late) Blight  | 42 [56] (168) sq. ft.   | 0   | Begin when disease threatens and repeat as required to maintain control.   |
| Celery   | Cercospora (Early) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression) | 56 [75] (224) sq. ft.   | 7   | Begin when plants are set in the field.  |
| Corn (Sweet)                                     | Helminthosporium Leaf Blight, Rust   | 42 [56] (168) sq. ft.   | 14  | Begin when conditions favor disease development. Repeat as needed. Do not apply to sweet corn grown for processing.      |

| Crop(s)                                      | Disease(s)   | Area to Cover              | PHI | Comments  |
|--|--|----------------------------|-----|---|
| Cucumbers,<br>Melons,<br>Pumpkins,<br>Squash | Anthrachnose, Downy Mildew,<br>Target Spot   | 42 [56] (168)<br>sq. ft.   | 0   | Begin when plants are in the<br>first true leaf stage or when<br>conditions are favorable for<br>disease.                     |
|  | Scab, Gummy Stem Blight<br>(Black Rot), Leaf Blight,<br>Cercospora Leaf Spot,<br>Powdery Mildew (except<br>Southwestern states)  | 28 [37.5] (112)<br>sq. ft. |     |   |
| Cucumbers                                    | Fruit Rot (Rhizoctonia)<br>[suppression]   |                            |     |   |
| Onion (Green),<br>Leek, Shallot              | Botrytis Leaf Blight (Blast),<br>Purple Blotch, Downy Mildew<br>(suppression)  | 28 [37.5] (112)<br>sq. ft. | 14  | Apply at first sign of disease.   |
| Onion (Dry Bulb),<br>Garlic                  | Botrytis Leaf Blight (Blast),<br>Purple Blotch, Downy Mildew<br>(suppression)  | 28 [37.5] (112)<br>sq. ft. | 7   | Apply at first sign of disease.   |
| Potato                                       | Early Blight, Late Blight,<br>Botrytis Vine Rot  | 56 [75] (224)<br>sq. ft.   | 0   | Begin when plants are 6<br>inches high or when disease<br>threatens and continue as<br>needed to maintain disease<br>control. |
| Tomato                                       | Early Blight, Late Blight, Gray<br>Leaf Spot, Gray Leaf Mold,<br>Septoria Leaf Spot,<br>Anthrachnose, Alternaria Fruit<br>Rot (Black Mold), Rhizoctonia<br>Fruit Rot, Botrytis Gray Mold | 28 [37.5] (112)<br>sq. ft. | 0   | Begin when conditions favor<br>disease development.   |

### FRUIT DISEASES

Apply a full coverage spray to the point of runoff to thoroughly cover tree canopy including undersides of leaves.

| Crop(s)                   | Disease(s)                                     | Comments  |
|---------------------------|--|---|
| Apricot, Nectarine, Peach | Peach Leaf Curl, Coryneum Blight<br>(Shothole) | Make one application at leaf fall in<br>late autumn. Make 1 or 2<br>applications in mid- to late-winter<br>before buds begin to swell. Where<br><i>Coryneum Blight (Shothole)</i> occurs,<br>apply an additional application 1 to 2<br>weeks after petal fall or at shucksplit<br>(tiny fruit just beginning to form) to<br>prevent fruit infections. |
|                           | Brown Rot Blossom Blight                       | Make 1 application at popcorn (pink,<br>red or early white bud stage when<br>first color shows) and a second<br>application at full bloom. If cool,<br>moist weather continues, make an<br>additional application at petal fall.  |

| Crop(s)                           | Disease(s)                      | Comments  |
|-----------------------------------|---------------------------------|---|
| Apricot, Peach, Nectarine, Cherry | Scab                            | Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit. Do not apply after shucksplit or before harvest. |
|                                   | Cherry Leaf Spot                | Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit.  |
|                                   | Cherry Leaf Spot (post harvest) | Make 1 application to foliage within 7 days after fruit is removed. In trees with a history of high leaf spot incidence, make a second application 10 to 14 days later. |

**ORNAMENTAL DISEASES**

Spray foliage to the point of runoff. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. Only for outdoor use.

| Species                                      | Diseases Controlled                                     | First Application               |
|--|---|---------------------------------|
| <b>Broadleaf Shrubs and Trees</b>            |   |                                 |
| Ash (Fraxinus)                               | Cercospora, Cercosporidium, Cylindrosporium Leaf Spots  | Spring bud break                |
| Azalea*, Rhododendron*                       | Phytophthora Die-back, Ovulinia Flower Blight           | New leaf emergence, early bloom |
| Buckeye, Horsechestnut                       | Leaf Blotch, Anthracnose                                | Spring bud break                |
| Cherry-Laurel                                | Cercospora Leafspot                                     | Petal fall                      |
| Crabapple                                    | Scab, Cedar-Apple Rust, Sphaeropsis Leaf Spot           | Spring bud break                |
| Dogwood                                      | Anthracnose (Discula sp.), Septoria Leaf Spot           | Spring bud swell, early bloom   |
| Euonymus                                     | Anthracnose   | Spring bud break                |
| Firethorn (Pyracantha)                       | Scab  | Spring bud break                |
| Flowering Almond, Quince and Cherry Hawthorn | Monilinia Blossom/Branch Blight Rust, Fabraea Leaf Spot | Early bloom, pre-bloom          |
| Holly  | Rhizoctonia Web Blight                                  | Warm, moist conditions          |
| Mountain Laurel                              | Cercospora Leaf Spot                                    | Spring bud break                |
| Oak (red group only)                         | Taphrina Blister, Actinopelte Leaf Spot, Anthracnose    | Dormant bud swell               |
| Oregon-Grape (Mahonia)                       | Rust  | Spring bud break                |
| Photinia                                     | Fabraea (Entomosporium) Leaf Spot                       | Spring bud break                |
| Pieris (Andromeda)                           | Phytophthora Die-Back                                   | New leaf emergence              |



|  |   |  |
|--|---|--|
| Poplar   | Marssonina Leaf Spot  | Spring bud break                               |
| Privet   | Cercospora Leaf Spot  | Prolonged wet conditions                       |
| Sycamore, Planetree  | Anthrachnose  | Spring bud break                               |
| Viburnum   | Powdery Mildew  | Mid-summer                                     |
| <b>Bulbs and Flowering Plants</b>  |   |  |
| Carnation  | Alternaria Leaf Spot/Branch Rot, Botrytis   | Transplant of cuttings; cool, moist conditions |
| Chrysanthemum/Daisy  | Mycosphaerella Ray Blight, Septoria Leaf Spot   | Transplant of cuttings                         |
| Geranium   | Botrytis Flower Blight (Gray Mold), Rust  | Pre-bloom cool, moist conditions               |
| Gladiolus  | Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot                                | Early propagation                              |
| Hollyhock  | Rust  | Early seedling stage                           |
| Hydrangea* (foliage only)  | Cercospora and Septoria Leaf Spots, Rust  | Early propagation                              |
| Iris   | Botrytis Blossom Blight, Didymellina Leaf Spot  | Cool, moist conditions                         |
| Lily   | Botrytis Gray Mold (foliar phase), Phytophthora Blight                                | Pre-bloom                                      |
| Petunia*   | Botrytis Gray Mold (foliar phase), Phytophthora Blight, Botrytis Blight               | Pre-bloom                                      |
| Rose   | Black Spot, Botrytis Blight   | Spring bud break                               |
| Statice  | Anthrachnose, Cercospora, Alternaria, Botrytis Leaf Blight                            | Spring bud break                               |
| Zinnia   | Powdery Mildew  | First sign of disease                          |
| *Discoloration of bloom has been noted on certain varieties when applications are made during flowering. |   |  |
| <b>Foliage Plants</b>  |   |  |
| Dracaena   | Fusarium Leaf Spot  | Pre-transplant                                 |
| Leatherleaf Fern   | Ascochyta Blight, Cercospora Leaf Spot, Cylindrocladium Leaf Spot, Rhizoctonia Blight | Spring bud break                               |
| Parlor Palm (Chamaedorea)  | Bipolaris Leaf Spot (Helminthosporium)  | Cool, moist conditions                         |
| Prayer Plant (Maranta)   | Helminthosporium Leaf Spot  | Early propagation                              |
| Oyster Plant (Rhoeo)   | Tan Leaf Spot   | Early propagation                              |
| Philodendron   | Phytophthora Blight, Dactylaria Leaf Spot   | Moist conditions                               |
| Syngonium  | Cephalosporium Leaf Spot  | Warm, moist conditions                         |

**Storage and Disposal**

**Pesticide Storage:** Store in a cool, dry (preferably locked) place inaccessible to children and pets. **Pesticide Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

19  
11

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

### First Aid

|  |   |
|--|---|
| <b>If Swallowed</b>  | <ul style="list-style-type: none"><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li><li>• Immediately call a poison control center or doctor for treatment advice.</li></ul> |
| <b>If Inhaled</b>  | <ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>  |
| <b>If on Skin or Clothing</b>  | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Immediately rinse skin with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If in Eyes</b>  | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-420-9347 for emergency medical treatment information. |   |

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

### Environmental Hazards

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

**Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.**

{Code}

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347  
If you have questions relating to product use or application, please call (800) 969-7200

11  
11

If you have a product spill or transportation emergency call (800) 424-9300  
For Material Safety Data Sheets please go to [www.gardentech.com](http://www.gardentech.com), or call (800) 969-7200

NOTICE {ATTENTION}: This product contains chlorothalonil, a chemical known to the State of California to cause cancer in laboratory animals.

[ ] Denotes alternate/optional language  
{ } Denotes language that does not appear on the market labeling

