1912010



1512-2

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Ms. Kim Davis RegWest Company, LLC Agent for Contract Packaging, Inc. 8203 West 20th. St., Suite A Greely, CO 80634-4696 SEP - 9 2010

Subject: Notification for Label Revisions under PRN 98-10, 2008-1 and 2007-4 Storage & Disposal, Environmental Hazards Statement and Other General Changes

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 98-10, 2008-1 and 2007-4 Storage & Disposal, Environmental Hazards Statement and Other Changes dated August 19, 2010 for:

EPA Registration 67572-2 "CP Multi-Purpose Fungicide Ready-To-Use"

The Registration Division (RD) has conducted a review of the request(s) for applicability under PRN 98-10, 2008-1 and 2007-4 and finds that the label changes requested fall within the scope of PRN 98-10, 2008-1 and 2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on <u>non-refillable</u> containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.


If you have any questions, please call me directly at 703-305-5335 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely. Hotrodone

Paul J. Mastradone, Ph.D., Acting Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

	• • • •	<i>٦</i>				~~~~			201
D))	_		
Please read instructions on	reverse before comple United State								MB No. 2070-0060. Intifier Number
Enviro	nmental Prote		v		_	Registrat			
	0 Pennsylvania Av		У	C]	Amendm	ent	No	tification
	Vashington, DC 20					Other		110	
		pplication fo	r Pestici	ide – Se	ecti	onl			
1. Company/Product Numb				EPA Produ			3. Pr	oposed C	lassification
	PA Reg. No. 67572-	2		То	ny K	lish			
	pose Fungicide Re			Λ#	22			None	Restricted
5. Name and Address of A		ode)							th FIFRA Section
Contract Packaging, In				c)(3)(b)(i) i beling to:	my p	oroduct is simi	lar or ic	dentical i	n composition and
c/o RegWest Company				Jenny to.					
8203 West 20th St., Suit			EF	PA Reg. N	o				
Greeley, CO 80634-469	90			Ū.					
Ch	eck if this is a new add				ne				
		S	<u>ection –</u>						
Amendment – Explain	below.			Final printed	d labe	els in response	to Agen	cy_letter_d	ated
Resubmission in respo	onse to Agency letter da	ated		"Me Too" A	pplica	ation.	IVVI	TUA	11 1 11
Notification – Explain b				Other – Exp			sei	P09;	วกเก
Explanation: Use ad	······	san (For Section					Que have h		
the labeling or the confidential I further understand that if this FIFRA and I may be subject to o	notification is not consist	tent with the terms of nalties under sections	PR Notices 98-	10, 2007-4 aı IFRA.					
1. Material This Product \	Vill Be Packaged In:			111					
Child Resistant Packaging	Unit Packaging		Water Solub	e Packagin	g			ontainer	
☐ Yes*	Yes		🗖 Yes				Aetal		
No	No No		No No			■	Plastic		
*Certification must be	If "Yes"	No. Per	If "Yes"		No. P		Slass		
submitted	Unit Packaging Wt.	Container	Packaging W	/t.	Conta		Paper		
3. Location of Net Contents		4. Size(s) Retail	Container	1		5. Location	Other (S		
	sinonnauon		Juntainer			On Lat		er Directio	
Label	Container	1 qt., 24	fl. oz. and	1 gallon				ccompan	ying Product
6. Manner in Which Label i	s Affixed to Product	🗖 Lith	ograph		Ster	nciled		,	
			er glued		Oth		: Self-ad	dhesive	,
			ection - I	V					
1. Contact Point (Comp	olete items directly be				ontac	cted, if necess	ary, to L	process t	
Name		Title				Telephone	No. (Incl	lude Areá	Ccdo)
								0.353.0e	
Kim Dav	/is		nsultant/Ag	ent			kim@	pregwes	
certify that the state	ements I have ma	Certification ade on this for		attachme	ents	thereto are	true,	Rec	e Application eived
accurate and complete	e. I acknowledge	that any knowin	igly false o						(Stamped)
<u>pun</u> ishable by fine or i		oth under applic	able law.						
2. Signature		3. Title							
			~	onsultant	1.	ont			
4. Typed Name		5. Date		Unsuitail	<u>arge</u>	51fL		1	
			-						
Kim Davis			August 19, 2010						

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.



Label Revision Notification



August 19, 2010

Mr. Tony Kish, Product Manager 22 Document Processing Desk - *NOTIF* Office of Pesticide Programs – *7504P* **U.S. Environmental Protection Agency** One Potomac Yard (South Building), Room S-4900 2777 South Crystal Drive Arlington, VA 22202

Dear Mr. Kish:

Subject: Contract Packaging, Inc. CP Multi-Purpose Fungicide Ready-To-Use EPA Reg. No. 67572-2

On behalf of *Contract Packaging, Inc.*, RegWest Company, LLC is submitting the attached documents per PR Notices 98-10, 2007-4 and 2008-1. RegWest will act as sole agent in this endeavor.

Attached are the following documents:

- 1. Application for Pesticide: Notification
- 2. One Reference Label (illustrating the label revisions)
- 3. Two New Labels (incorporating the revisions)

In addition to the PR Notices changes, we are deleting the "General Precautions and Restrictions" and replacing it with "Product Precautions and Restrictions". Please refer to the *Reference Label* for an illustration of all revisions.

Please contact me at 970.353.0611 or kim@regwest.com if you have any questions or comments concerning this submission.

Regards, RegWest Company, LLC

Kim Davis, Consultant/Agent

Attachments

cc: CPI - Hacker, S.

8203 West 20th St., Suite A Greeley, C0 80634-4696 p 970-353-0611 f 970-353-0613

ງເງລວັ ວ ງງງງວ່າ

.....



67572-2 New Label: Page 1 of 7 11/02/2007 SAL + 05/13/2010 & 08/19/2010 Notification

CP Multi-Purpose Fungicide Ready-To-Use

Ready-to-Use Fungicide Disease Control Contains Echo™ LITE CHLOROTHALONIL FUNGICIDE

Controls/Stops and Prevents over 70 Diseases on Flowers, Vegetables, Shrubs, Fruit and Shade Trees

Controls Diseases:

Leaf Spots, Black Spots, Rust[s], Blights, Fruit Rots, Mildew[s], Scab, Molds, Lawn Fungus, etc. on: Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees NOTIFICATION

> Use on Roses For Use on Vegetables, Flowers and Shrubs

SEP 0 9 2010

Ready-To-Use For Healthier Plants For Control of Plant Disease

Active Ingredient:

Chlorothalonil (CAS No. 1897-45-6)	. 0.087%
Other Ingredients	<u>99.913%</u>
Total	100.000%

Keep Out of Reach of Children CAUTION

See back panel [booklet] for First Aid and additional Precautionary Statements.

EPA Reg. No. 67572-2


{Booklet}

EPA Est. No. 67572-GA-1

Net Contents: 1 qt. (32 fl. oz.) (946 ml) [24 fl. oz.] [1 gal.]

First Aid If Swallowed: Have person sip a glass of water if able to swallow. . Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. • Immediately call a poison control center or doctor for treatment advice. . If Inhaled: Move person to fresh air. ٠ If person is not breathing, call 911 or an ambulance, then give artificial respiration, . preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. If on Skin Take off contaminated clothing. • or Clothing: Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-420-9347 for emergency medical treatment information.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

Environmental Hazards

This pesticide is toxic to aquatic invertebrates and wildlife. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Peel Back Tab for Additional Information

Ornamental Diseases

Cercospora, Phytophthora Die-back, Ovulinia Flower Blight, Leaf Blotch, Scab, Anthracnose, Fabraea (*Entomosporium*) Leaf Spot, Rhizoctonia Web Blight, Powdery Mildew, Rust, Botrytis, Alternaria Leaf Spot/Branch Rot, Septoria Leaf Spot, Black Spot, Helminthosporium Leaf Spot, Botrytis Gray Mold (foliar phase), Phytophthora Blight, Cylindrocladium Leaf Spots, Marssonina Leaf Spot and many more listed inside.

Vegetable Diseases

Rust, Botrytis Blight (Gray Mold), Alternaria Leaf Spot, Downy Mildew, Ring Spot, Cercospora (Early) Blight, Alternaria (Late) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression), Helminthosporium Leaf Blight, Anthracnose, Scab, Purple Blotch, Botrytis Vine Rot and many more listed inside.

Fruit Diseases

Peach Leaf Curl, Coryneum Blight (Shothole), Cherry Leaf Spot, Brown Rot Blossom Blight, Scab.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Product Precautions and Restrictions

Not for use on plants being grown for sale or other commercial use, commercial seed production or research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in interior plantscapes and ornamental gardens or parks. Also for use on fruit trees and vegetables. Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets either directly or through drift. Do not use on home lawns and turf sites associated with apartment buildings,



67572-2 New Label: Page 3 of 7 11/02/2007 SAL + 05/13/2010 & 08/19/2010 Notification

daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle or high schools), campgrounds, churches and theme parks.

To use the squeeze and pour bottle, remove cap and inner seal from the calibrated dosing chamber. Then replace cap and tighten, then loosen ¼ turn. Gently squeeze bottle to fill chamber to the appropriate dose. Tilt the bottle to the direction indicated by the arrow on the bottle. Pour concentrate directly into sprayer and slowly add the appropriate amount of water. Replace cap.

Application Instructions

Shake Well Before Using Read Entire Label and Use Strictly in Accordance With Precautionary Statements and Directions

Personal Protective Equipment (PPE): Wear long sleeved shirt, long pants and chemical-resistant gloves made of any waterproof material while mixing and applying this product.

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when used according to the label directions. Thorough uniform coverage of plant surfaces is essential for good disease control. Keep children and pets off treated area until dry.

Maximum Application Rates

For each site listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased. **Note:** AI = active ingredient; A = acre.

Crop(s)	Maximum Individual Application Rate (Lbs./Al/A)	Minimum Retreatment Interval Days	Maximum Seasonal Total Application Rate (Lbs./Al/A/Season)
Bean (snap)	2.25	7	9
Carrot	1.5	7	15
Celery	2.25	7	18
Cole crops	1.5	7	12
Corn (sweet)	1.5	7	9
Cucurbits	2.25	7	15.75
Ornamentals	1.55	7	36.5 ¹
Roses	1.1	7	36.4 ¹
Onion (dry bulb) and garlic	2.25	7	15
Onion (green), leak, shallot	2.25	7	6.7
Potato	1.125	5	11.25
Stone fruits (including cherries)	3.1	10	15.5
Tomato	2.16	7	15.1

¹Seasonal maximum applies only to field-grown



Vegetable Diseases

Note: Do not spray plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

PHI = the number of days to wait between the last application and harvest.

{Sq. ft. not in brackets or parentheses is for the 24 fl. oz. market container; sq. ft. in brackets is for the 32 fl. oz. container; sq. ft. in parentheses is for the 1 gallon container.}

Crop(s)	Disease(s)	Area to Cover	PHI	Comments
Beans (Green)	Rust, Botrytis Blight (Gray Mold)	28 [37.5] (112) sq. ft.	7	Begin applications during early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Alternaria Leaf Spot, Downy Mildew	56 [75] (224) sq. ft.	0	Begin after transplants are set in the field or when conditions favor disease development.
Brussels Sprouts (CA only)	Ring Spot	42 [56] (168) sq. ft.	0	Begin at time of early sprout development or when conditions favor disease development.
Carrots	Cercospora (Early) Blight, Alternaria (Late) Blight	42 [56] (168) sq. ft.	0	Begin when disease threatens and repeat as required to maintain control.
Celery	Cercospora (Early) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (Suppression)	56 [75] (224) sq. ft.	7	Begin when plants are set in the field.
Corn (Sweet)	Helminthosporium Leaf Blight, Rust	42 [56] (168) sq. ft.	14	Begin when conditions favor disease development. Repeat as needed. Do not apply to sweet corn grown for processing.
Cucumbers, Melons, Pumpkins, Squash	Anthracnose, Downy Mildew, Target Spot Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states)	42 [56] (168) sq. ft. 28 [37.5] (112) sq. ft.	0	Begin when plants are in the first true leaf stage or when conditions are favorable for disease.
Cucumbers	Fruit Rot (Rhizoctonia) [suppression]			
Onion (Green), Leek, Shallot	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	28 [37.5] (112) sq. ft.	14	Apply at first sign of disease.
Onion (Dry Bulb), Garlic	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	28 [37.5] (112) sq. ft.	7	Apply at first sign of disease.



67572-2 New Label: Page 5 of 7 11/02/2007 SAL + 05/13/2010 & 08/19/2010 Notification

Crop(s)	Disease(s)	Area to Cover	PHI	Comments
Potato	Early Blight, Late Blight, Botrytis Vine Rot	56 [75] (224) sq. ft.	0	Begin when plants are 6 inches high or when disease threatens and continue as needed to maintain disease control.
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot, Botrytis Gray Mold	28 [37.5] (112) sq. ft.	0	Begin when conditions favor disease development.

Fruit Diseases

Apply a full coverage spray to the point of runoff to thoroughly cover tree canopy including undersides of leaves

Crop(s)	Disease(s)	Comments
Apricot, Nectarine, Peach	Peach Leaf Curl, Coryneum Blight (Shothole)	Make one application at leaf fall in late autumn. Make 1 or 2 applications in mid- to late-winter before buds begin to swell. Where <i>Coryneum</i> <i>Blight (Shothole)</i> occurs, apply an additional application 1 to 2 weeks after petal fall or at shucksplit (tiny fruit just beginning to form) to prevent fruit infections.
	Brown Rot Blossom Blight	Make 1 application at popcorn (pink, red or early white bud stage when first color shows) and a second application at full bloom. If cool, moist weather continues, make an additional application at petal fall.
Apricot, Peach, Nectarine, Cherry	Scab	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit. Do not apply after shucksplit or before harvest.
	Cherry Leaf Spot	Make 1 application at popcorn and a second application at full bloom. In addition, make one application at shucksplit.
	Cherry Leaf Spot (post harvest)	Make 1 application to foliage within 7 days after fruit is removed. In trees with a history of high leaf spot incidence, make a second application 10 to 14 days later.



Ornamental Diseases

Spray foliage to the point of runoff. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. For outdoor use only.

Species	Diseases Controlled	First Application
	Broadleaf Shrubs and Trees	
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring bud break
Azalea*, Rhododendron*	Phytophthora Die-back, Ovulinia Flower Blight	New leaf emergence, early bloom
Buckeye, Horsechestnut	Leaf Blotch, Anthracnose	Spring bud break
Cherry-Laurel	Cercospora Leafspot	Petal fall
Crabapple	Scab, Cedar-Apple Rust, Sphaeropsis Leaf Spot	Spring bud break
Dogwood	Anthracnose (Discula sp.), Septoria Leaf Spot	Spring bud swell, early bloom
Euonymus	Anthracnose	Spring bud break
Firethorn (Pyracantha)	Scab	Spring bud break
Flowering Almond, Quince and Cherry Hawthorn	Monilinia Blossom/Branch Blight Rust, Fabraea Leaf Spot	Early bloom, pre-bloom
Holly	Rhizoctonia Web Blight	Warm, moist conditions
Mountain Laurel	Cercospora Leaf Spot	Spring bud break
Oak (red group only)	Taphrina Blister, Actinopelte Leaf Spot, Anthracnose	Dormant bud swell
Oregon-Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea (Entomosporium) Leaf Spot	Spring bud break
Pieris (Andromeda)	Phytophthora Die-Back	New leaf emergence
Poplar	Marssonina Leaf Spot	Spring bud break
Privet	Cercospora Leaf Spot	Prolonged wet conditions
Sycamore, Planetree	Anthracnose	Spring bud break
Viburnum	Powdery Mildew	Mid-summer
	Bulbs and Flowering Plants	
Carnation	Alternaria Leaf Spot/Branch Rot, Botrytis	Transplant of cuttings; cool, moist conditions
Chrysanthemum/Daisy	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of cuttings
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-bloom cool, moist conditions
Gladiolus	Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea* (foliage only)	Cercospora and Septoria Leaf Spots, Rust	Early propagation
Iris	Botrytis Blossom Blight, Didymellina Leaf Spot	Cool, moist conditions

11/02/2007 SAL + 05/13/2010 & 08/19/2010 Notification

Species	Diseases Controlled	First Application			
Lily	Botrytis Gray Mold (foliar phase), Phytophthora Blight	Pre-bloom			
Petunia*	Botrytis Gray Mold (foliar phase), Phytophthora Blight, Botrytis Blight	Pre-bloom			
Rose	Black Spot, Botrytis Blight	Spring bud break			
Statice	Anthracnose, Cercospora, Alternaria, Botrytis Leaf Blight	Spring bud break			
Zinnia	Powdery Mildew	First sign of disease			
*Discoloration of bloom has been	n noted on certain varieties when application	ons are made during flowering.			
Foliage Plants					
Dracaena	Fusarium Leaf Spot	Pre-transplant			
Leatherleaf Fern	Ascochyta Blight, Cercospora Leaf Spot, Cylindrocladium Leaf Spot, Rhizoctonia Blight	Spring bud break			
Parlor Palm (Chamaedorea)	Bipolaris Leaf Spot (Helminthosporium)	Cool, moist conditions			
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early propagation			
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early propagation			
Philodendron	Phytophthora Blight, Dactylaria Leaf Spot	Moist conditions			
Syngonium	Cephalosporium Leaf Spot	Warm, moist conditions			

Storage and Disposal

Pesticide Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets. **Pesticide/Container Disposal:** Non-refillable container; do not reuse or refill this container. **If empty:** Offer for recycling, if available; otherwise, discard in trash. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

{Per PR Notice 2007-4 the batch code/lot number will appear on the label or container.}

To the extent not prohibited by applicable law, buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

{Code}

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347 If you have questions relating to product use or application, please call (800) 969-7200 If you have a product spill or transportation emergency call (800) 424-9300 For Material Safety Data Sheets please call (800) 969-7200

Notice (Attention): This product contains chlorothalonil, a chemical known to the State of California to cause cancer in laboratory animals.

Manufactured [Distributed] by Contract Packaging, Inc. P.O. Box 1508 Covington, GA 30015-1508

[] Denotes alternate/optional language

{ } Denotes language that does not appear on the market labeling