

R & M ALOE EAR MITE TREATMENT

With Aloe and Lanolin
Dogs • Puppies • Cats • Kittens



For Dogs,



Cats



and Rabbits

ACCEPTED
NOV 13 2000
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67572-40

Kills Ear Mites on Contact

ACTIVE INGREDIENTS:

Pyrethrins (CAS No. 8003-34-7)	0.05%
Piperonyl Butoxide, Technical (CAS No. 51-03-6)	0.50%
OTHER INGREDIENTS	99.45%
Total	100.00%

Keep Out of Reach of Children
CAUTION
See back panel for additional precautionary statements.

EPA Reg. No. 67572-40

EPA Est. 67572-GA-1

Net Contents: ____ fl. oz.

Manufactured by:
CONTRACT PACKAGING, INC.
P.O. Box 1508
Covington, GA 30015-1508

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS, CATS AND RABBITS**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets and away from heat or open flame.
Disposal: Do not reuse empty container; wrap and put in trash.

Ear mites usually induce the formation of a dry, dark brown, waxy exudate with crusts in the ears of pets. Ear mites may be detected by placing some of the ear exudate on a dark surface and carefully watching for the movement of tiny white specks away from the exudate. Inflamed, watery or blocked ear canals indicate a more serious condition that requires the services of a veterinarian.

Do not use on puppies or kittens under 12 weeks. Consult a veterinarian before using this product on debilitated, medicated, aged, pregnant or nursing animals.

To Kill Ear Mites: Clean ears to remove built-up wax and dirt. While firmly holding pet, fill ear canal with recommended number of drops in each ear. Massage base of ear to ensure insecticidal action penetrates ear wax. Gently dry ear with a cotton ball, soft cloth or cotton swab. Reapply every 2 weeks to reduce the possibility of ear mite infestation and the accumulation of ear wax. If no improvement is seen after 2 applications, consult a veterinarian.

BODY WEIGHT	DOSAGE
5 to 15 lbs.	4 to 5 drops
15 to 30 lbs.	5 to 10 drops
30 lbs. and over	10 to 15 drops

If condition for which this preparation is used persists or if an irritation develops, discontinue use and consult a veterinarian.

For use with a swabstick: Open packet, remove swabstick. While firmly holding pet, fill ear with liquid from packet. Massage base of ear to ensure that insecticidal action penetrates ear wax. Use swabstick to remove as much waxy exudate as possible from the outer ear. **IMPORTANT:** Do not use the swabstick in the ear canal (i.e., swab part should always be visible!) to avoid damaging the delicate tissue of the ear. Reapply every 2 weeks.

**PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals**

CAUTION: Humans: Harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes. Thoroughly wash with soap and water after use. Do not apply directly to food or contaminate food or feedstuffs with this product. **Animals:** Avoid contact with eyes. If in eyes, flush with water. Do not use on meat or milk producing animals. For external use only. Do not use for a prolonged period of time. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, immediately consult a veterinarian.

First Aid

If Swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible • Call a poison control center or doctor for further treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.

Buyer assumes all risk of use, storage or handling of this product not in strict accordance with this label.

To obtain additional information or in case of an emergency contact 770-787-1977.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}