
67572 -56



67572-56 New Label: Page 1 of 3 12/20/99

> registered under HPA Reg. No. 67

CP SNAIL & BUG BAIT

For the Control of Sowbugs, Snails, Ants, Grasshoppers, Slugs, Cutworms and Crickets

This pesticide may be fatal to children and dogs or other pets if eaten. Keep children and pets out of treated areas.

ACTIVE INGREDIENTS:

Keep Out of Reach of Children		Under t Fungleic as amer	he Fr is, az Ided,	deret 1 d Rode for the	nsectici nticida pestici	id o , Act, Ida
SEVIN is a registered trademark of Rhône-Poulenc Ag Compan	M	IAR .	3 0	2000		
Contains SEVIN brand carbaryl insecticide and metaldehyde.	100.0%	AU	U.	出上	TE	L D
INERT INGREDIENTS	95.0%	TA M	2	13 14	-	
Metaldehyde	1.0%					
Carbaryl (1-naphthyl N-methylcarbamate)	4.0%					

Keep Out of Reach of Children WARNING See back panel for additional precautionary statements.

Net Weight: ____ pounds

EPA Reg. No. 67572-56 EPA Est. 67572-GA-1

CONTRACT PACKAGING, INC. P.O. Box 1508 Covington, GA 30015

Rack Panelj

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: May be fatal if inhaled or swallowed. Harmful dust. Avoid breathing dust. Wash thoroughly after handling. Wash hands and face before eating. Avoid contact with skin, eyes and clothing. Do not take internally. Wear regular long-sleeved work clothes. Change to clean clothes daily. Keep out of reach of children and animals.

First Aid

If Swallowed:	Call a poison control center or doctor immediately for treatment advice.						
	• Have person sip a glass of water if able to swallow.						
	• Do not induce vomiting unless told to by a poison control center or doctor.						
If Inhaled:	Move person to fresh air.						
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mout to-mouth if possible.						
	Call a poison control center or doctor for further treatment advice.						
If on Skin:	Take off contaminated clothing.						
	• Rinse skin immediately with plenty of water for 15-20 minutes.						
	Call a poison control center or doctor for treatment advice.						
If in Eyes:	 Hold eye open and rinse slowly with water for 15-20 minutes. Remove contact lenses, if prosent, after the first 5 minutes, then continue rinsing. 						
	Call a poison control center or doctor for treatment advice.						
Have the produ	uct container or label with you when calling a poison control center or doctor or going for treatment.						
	cian: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.						

67572-56 New Label: Page 2 of 3 12/20/99

Environmental Hazards

,

This product is highly toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Bee Caution: This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treated area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This bait may be used to protect flowers, ornamentals, shrubs, deciduous fruits and the vegetables named below. Apply on the ground and adjacent to plants. Do not apply to edible parts of the plants. Broadcast applications may be used when there is a heavy infestation. Repeat applications as necessary.

Storage and Disposal

Storage: Store in a cool, dry (preferably locked) place inaccessible to children and domestic animals. Disposal: Do not reuse empty container. Wrap container and put in trash.

Сгор	Maximum Single Application Rate (lbs. / 1000 sq. ft.)	Maximum Seasonal Application Rate (lbs. / 1000 sq. ft.)	Retreatment Interval (days)	PHI (days)	- Comments
Asparagus (spears)	1.1	3.4	3	1	Total maximum seasonal application rate is
Asparagus (ferns*)	1.1	2.3	7	1	23 lbs. per 1000 sq. ft. per crop. The total number of applications is 3 prior to harvest with 2 additional applications only to ferns. *after harvest of spears.
Beans (fresh & dried):	.85	3.4	3	3 (S)	S = succulents
[snap, navy & kidney],				21 (D)	D = dried
Peas (fresh & dried).				14 (F)	$\mathbf{F} = \mathbf{forage}$
Cowpeas, Lentils				21 (H)	H = hay
Garden beets (roots),	1.1	3.4	7	7	Total number of applications per crop can be
Carrots, Horseradish,	3		1		6 (some of which must be at lower than
					maximum single application rate)
Brussels sprouts,	1.1	3.4	7	3 (A)	A = Brussels sprouts, broccoli, cabbage,
Broccoli, Cabbage,				14 (B)	kohlrabi and cauliflower
Cauliflower,					B = Chinese cabbage, kale and collards
Chinese cabbage,					Total number of applications per crop can be
Collards, Kale, Kohlrabi					4 (some of which must be at lower than maximum single application rate)
Cucumber, melon,	.57	3.4	7	3	
squash))) 2 2 4 2 3)
Lettuce (head & leaf),	1.1	3.4	7	14-15	Total number of applications 'per 'crop can be
Endive, Celery,					5 (some of which must be at lower than
Dandelion, Parsley,					maximum single application rate)
Spinach, Garden Beets					
(tops), Turnips (tops)					
Potatoes	1.1	3.4	7	7	Total number of applications.per.crop can be
					6 (some of which must be at lover than
					maximum single application rate)
Tomato, Pepper,	1.1	4.6	7	3	Total number of applications' pet 'crop can be
Eggplant	•)			7 (some of which must be at lower than
Omomentale handers	1.2	2.4	None		maximum single application rate)
Ornamentals, borders	1.2	<u> </u>	None	None	<u> </u>

67572-56 New Label: Page 3 of 3 12/20/99

Special Instructions for Improved Performance are:

- 1. Slugs and snails favor dark, damp places. Bait where these pests stay during daylight hours.
- 2. Be sure area to be baited is damp. In dry weather, water area well before baiting.
- 3. Apply bait in evening as these pests feed mainly at night.
- 4. In ornamental and border areas, until infestation is under control, bait should be renewed at least every 7 to 10 days or more frequently if necessary.
- 5. Since snails and slugs migrate extensively, when invasions from adjacent areas are anticipated a barrier of the bait will aid in control.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.