



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 19 2001

Kim Davis, Agent for Contract Packaging, Inc.
RegWest Company
30856 Rocky Road
Greeley, CO 80631-9375

Subject: CP Snail & Bug Bait
EPA Registration No. 67572-56
Label revisions submission dated October 5, 2000

Dear Ms. Davis:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

1. On page 3 of the copies of the label submitted for stamping the crop "Beans" appears before the heading that contains the crop, application rate, retreatment interval, etc. The information contained in the heading must appear before any crops in the final printed labeling if this portion of the label is split as in the submitted copies.

Submit two copies of your final printed labeling before you release the product for shipment. A copy of the labeling stamped "Accepted with Comments" is enclosed for your records.

Sincerely,

George Tompkins
George Tompkins, Ph.D., Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505C)

CP SNAIL & BUG BAIT

For the Control of Sowbugs, Snails, Ants, Grasshoppers, Slugs, Cutworms and Crickets

This pesticide may be fatal to children and dogs or other pets if eaten.
Keep children and pets out of treated areas.

ACTIVE INGREDIENTS:

Carbaryl (CAS No. 63-25-2)	4%
Metaldehyde (CAS No. 108-62-3).....	1%
OTHER INGREDIENTS	95%
Total	100%

Contains SEVIN® brand carbaryl insecticide and metaldehyde.
SEVIN is a registered trademark of Aventis Environmental Science

Keep Out of Reach of Children
WARNING
See back panel for additional precautionary statements.

ACCEPTED
with COMMENTS
In EPA Letter Dated:

JAN 19 2001


Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
67572-56

EPA Reg. No. 67572-56

EPA Est. 67572-GA-1

Net Weight: ___ pounds

Manufactured by:
CONTRACT PACKAGING, INC.
P.O. Box 1508
Covington, GA 30015-1508



XXXXXX = additions; XXXXXX = deletions

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: May be fatal if inhaled or swallowed. Harmful dust. Avoid breathing dust. Wash thoroughly after handling. Wash hands and face before eating. Avoid contact with skin, eyes and clothing. Do not take internally. Wear regular long-sleeved work clothes. Change to clean clothes daily. Keep out of reach of children and animals.

First Aid

If Swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency treatment information.	
Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.	

Environmental Hazards

This product is highly toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. **Bee Caution:** This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treated area. Time applications to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets. **Disposal:** Do not reuse empty container; wrap and put in trash.

Use this bait to protect flowers, ornamentals, shrubs, deciduous fruits and the vegetables named below. Apply on the ground and adjacent to plants. Do not apply to edible parts of the plants. When there is a heavy infestation use broadcast applications. Repeat applications as necessary.

Crop	Maximum Single Application Rate (lbs. / 1000 sq. ft.)	Maximum Seasonal Application Rate (lbs. / 1000 sq. ft.)	Retreatment Interval (days)	PHI (days)	Comments
Asparagus (spears)	1.1	3.4	3	1	Total maximum seasonal application rate is 23 lbs. per 1000 sq. ft. per crop. The total number of applications is 3 prior to harvest with 2 additional applications only to ferns. *after harvest of spears.
Asparagus (ferns*)	1.1	2.3	7	1	

Beans (fresh & dried): [snap, navy & kidney], Peas (fresh & dried), Cowpeas, Lentils	.85	3.4	3	3 (S) 21 (D) 14 (F) 21 (H)	S = succulents D = dried F = forage H = hay
Crop	Maximum Single Application Rate (lbs. / 1000 sq. ft.)	Maximum Seasonal Application Rate (lbs. / 1000 sq. ft.)	Retreatment Interval (days)	PHI (days)	Comments
Garden beets (roots), Carrots, Horseradish,	1.1	3.4	7	7	Total number of applications per crop can be 6 (some of which must be at lower than maximum single application rate)
Brussels sprouts, Broccoli, Cabbage, Cauliflower, Chinese cabbage, Collards, Kale, Kohlrabi	1.1	3.4	7	3 (A) 14 (B)	A = Brussels sprouts, broccoli, cabbage, kohlrabi and cauliflower B = Chinese cabbage, kale and collards Total number of applications per crop can be 4 (some of which must be at lower than maximum single application rate)
Cucumber, melon, squash	.57	3.4	7	3	
Lettuce (head & leaf), Endive, Celery, Dandelion, Parsley, Spinach, Garden Beets (tops), Turnips (tops)	1.1	3.4	7	14-15	Total number of applications per crop can be 5 (some of which must be at lower than maximum single application rate)
Potatoes	1.1	3.4	7	7	Total number of applications per crop can be 6 (some of which must be at lower than maximum single application rate)
Tomato, Pepper, Eggplant	1.1	4.6	7	3	Total number of applications per crop can be 7 (some of which must be at lower than maximum single application rate)
Ornamentals, borders	1.2	2.4	None	None	

Special Instructions for Improved Performance are:

1. Slugs and snails favor dark, damp places. Bait where these pests stay during daylight hours.
2. Be sure area to be baited is damp. In dry weather, water area well before baiting.
3. Apply bait in evening as these pests feed mainly at night.
4. In ornamental and border areas, until infestation is under control, renew bait at least every 7 to 10 days or more frequently if necessary.
5. Since snails and slugs migrate extensively, when invasions from adjacent areas are anticipated a barrier of the bait will aid in control.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

To obtain additional information or in case of an emergency call 770-787-1977.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}