



067572-00081-062399

Printed: 11:17:18 Thursday, 22 Jul, 1999 # 43 / 3745

Systems Integration Group, Inc.

PM04

67572-81

6/23/99

1 of 4

**SEVIN® GRANULES (1% Sevin®)**  
**ANT, FLEA, TICK & GRUB KILLER**

For Home Lawns and Ornamental/Recreational Turf

To Kill and Control Fire Ants, Ticks (including those that Vector Lyme Disease)  
and Nuisance Pests Around Homes

\* Kills Japanese Beetles \* Slow Release Granules \* Bug Killer

\* Kills Numerous [Over 20] Pests on Lawns and Ornamentals \* Covers Up To \_\_\_\_\_ Square Feet

**ACTIVE INGREDIENT:**

Carbaryl* .....	1.0%
<b>OTHER INGREDIENTS</b> .....	<b>99.0%</b>
*CAS Number 63-25-2	Total ..... 100.0%

**Keep Out of Reach of Children**  
**CAUTION**  
 See back panel for additional precautionary statements.

**ACCEPTED**  
 JUN 23 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67572-81

Net Weight: 5 pounds [Covers up to 1,111 square feet]  
 Net Weight: 10 pounds [Covers up to 2,222 square feet]  
 Net Weight: 20 pounds [Covers up to 4,444 square feet]

EPA Reg. No. 67572-81      EPA Est. 67572-GA-1

**CONTRACT PACKAGING, INC.**  
 P.O. Box 1508  
 Covington, GA 30015

[Back Panel]

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**Personal Protective Equipment (PPE)**

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks and household or latex rubber gloves. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables are provided, use detergent and hot water. Keep and wash PPE separately from other laundry.

**First Aid**

**If Swallowed:** Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting. **If Inhaled:** Remove person to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. **If in Eyes:** Flush with plenty of water. Call a physician if irritation persists. **If on Skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**Note to Physician:** Carbaryl is a moderate cholinesterase inhibitor. Atropine is antidotal.

**User Safety Recommendations**

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until granules have been absorbed into the soil.

**Environmental Hazards**

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for commercial seed production, research purposes, sale or other commercial use such as sod.

**Read Entire Label Before Using This Product**

Strictly observe label directions and cautions. Do not apply this product in a way that will contact applicators or other persons, either directly or through drift. Unprotected persons must vacate the area being treated.

{For product packaged in containers of 5 pounds or less:}

**Storage and Disposal**

**Storage:** Store in original container in a cool, dry place away from children and animals. Avoid excess heat. **Disposal:** Do not reuse empty container [bag]; wrap and put in trash.

{For product packaged in containers greater than five pounds:}

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container in a cool, dry place away from children and animals. Avoid excess heat. In case of spill, confine spill and avoid contact; isolate and keep away from animals and unprotected persons. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** **Plastic:** Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**GENERAL DIRECTIONS**

**FOR LAWNS, ORNAMENTAL AND RECREATIONAL TURF**

Use when insects first appear and/or plant damage is imminent. For best results, mow lawn first and then apply material uniformly to dry turfgrass. Irrigate 1/4" to wash insecticide into the grass. Do not allow use of treated areas during application. Keep children and pets off treated areas until grass is dry. Apply when air is calm to avoid drift and contact with skin and eyes. Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.

Treatment Areas	Pests	Pounds of This Product Per 1000 Sq. Ft.	Comments
Lawns, Ornamental and Recreational Turf Areas	Ants	4.5	For effective control, irrigate after application to remove chemical from granules.  For heavy infestations of Chinch bugs, increase rate to 10 lbs. per 1000 sq. ft. and treat the entire area.
	Lucerne moth		
	Armyworms		
	Millipedes		
	Centipedes		
	Mosquitoes (adults)		
	Chiggers		
	Scarab beetle adults:		
	Chinch bugs such as May beetle,		
	Cutworms June beetle,		
	Earwigs Japanese beetle,		
	Essex skipper Green June beetle		
European chafer Sod webworm			
Fall armyworm (lawn moths)			
Fiery skipper Sowbugs			
Grasshoppers Spittlebugs			

Green June beetle Larvae Leafhoppers	Springtails Ticks Yellowstriped Armyworms		
Billbug larvae European crane fly larvae Fleas White grubs: Such as Japanese beetle larvae, Chafer beetle larvae, <i>Phyllophaga</i> spp. larvae, Annual white grubs	Black turfgrass ataenius beetle larvae Hyperodes weevil (bluegrass weevil larvae)	18	For effective control of subsurface pests, apply when insects are actively feeding near the soil surface, usually during late March through May, or July through early September, or as recommended by local agricultural extension service agents. Irrigate thoroughly after application to remove chemical from granules and move it down into the thatch.

#### Control of Ticks That Vector Lyme Disease

*For control of juvenile and adult Ixodes spp. ticks (Deer Tick and Black Legged Tick) and Amblyomma spp. ticks (Lone Star Tick) which vector Lyme Disease:* Apply granules at the rate of 7 to 9 pounds of product per 500 sq. ft. with appropriate equipment to ensure uniform coverage. Target applications to control juvenile ticks (larvae and nymphs) in late spring or early summer (May/June). Control of adult ticks can be achieved with late summer and fall applications (August/September). Make applications to lawns, shrub beds, ornamental plantings, wooded areas and around outside perimeters of home and other buildings. Do not make spot applications. Treat entire lawn and perimeter wooded areas and property boundaries where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove chemical from the granules. Ticks may be reintroduced from surrounding areas or host animals. Reapply as necessary to maintain adequate control, but not more often than once every 7 days.

#### Imported Fire Ants

For use as a mound treatment to control imported fire ants occurring in lawns, turf and premise areas, apply 1 cup of granules over and around each mound (approximately 2' x 2'). Thoroughly irrigate mound immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be sprinkled over each mound, but do not use more than 2 gallons of water as flooding of the mound may result. Apply water gently to avoid disturbing ants and cause migration before chemical can be effective. For best results apply in cool weather (65°F to 80°F), early morning or late afternoon. Treat new mounds as they appear.

#### For Flowering Plants and Shrubs (Ornamental Plantings) in Landscape and Foundation Plantings

Pests	Pounds This Product Per 1000 Sq. Ft.	Specific Directions
Ants Armyworms Centipedes Crickets Cutworms Earwigs	4.5 to 10	Application to wet foliage during periods of high humidity may cause injury to tender foliage.  Apply granules in a 6-foot band around the building, to shrub and flowerbeds, foundations, ornamental plantings and lawn or soil areas immediately adjacent to the building. For most effective control, treated area should be lightly watered after application to remove the chemical from the granules.  Repeat as needed for control up to 3 times per year but do not apply more often than once every 7 days.  For plants above chest height, consider using a liquid carbaryl product.
Firebrats Fleas Millipedes Scorpions Silverfish Spiders Ticks		

### Nuisance Pest Control Around the Outside of Homes, Porches, Patios and Buildings

For control of Ants, Armyworms, Centipedes, Crickets, Cutworms, Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders and Ticks: Apply granules uniformly in an 8-foot band around the building. Apply granules at the rate of 4 2/3 lbs. per 1000 sq. ft. to shrub and flowerbeds, foundations, ornamental plantings and lawn and soil areas immediately adjacent to the building. For most effective control, treated area should be lightly watered after application to remove the chemical from the granules. Repeat application as necessary to control new infestations.

### Nuisance Pest Control in Vegetables and In Small Fruits and Berries (Not for Use in California)

For control of ants, centipedes, crickets, cutworms, earwigs, grasshoppers, millipedes, sowbugs (pillbugs), springtails and ticks, apply 5 pounds per 1000 sq.ft. evenly over the soil surface. Water lightly after application to release the chemical from the granules. Applications may be made at any time before, during or after the growing season. Do not make more than 4 applications per year or apply more often than once every 7 days during the growing season. Do not use on crops other than those listed below. Observe the following preharvest intervals for applications during the growing season:


Crops	Preharvest Interval (Days)
Asparagus	1
Sweet Corn	2
Broccoli, Brussel sprouts, Cabbage, Cauliflower, Kohlrabi	3
Fruiting Vegetables (including tomato, eggplant, pepper)	3
Root and Tuber Vegetables (including beets, carrots, parsnips, rutabagas, salsify, turnips, potatoes)	7
Sweet potatoes	7
Small Fruit and Berry crops (including grapes, strawberries, blackberries, dewberries, raspberries, blueberries)	7
Leafy Vegetables (including celery, dandelion, endive, lettuce, parsley, rhubarb, spinach, swiss chard)	14
Chinese cabbage, Collards, Kale, Mustard	14
Peanuts	14

Do not plant rotational crops not listed on this or other carbaryl labels in carbaryl-treated soil.

Spreader	Spreader Settings			
	Pounds / 1000 Sq. Ft.		Pounds / 500 Sq. Ft.	
	4.5 lbs.	10 lbs.	7 lbs.	9 lbs.
Scotts EvenGreen 18"	16	17*	16	17
Scotts Speedy Broadcast	15	16*	15	16
Red Devil Broadcast	13	14*	13	14
Red Devil Drop 20"	13	14*	13	14
Republic EZ Drop 18"	9	11	9	11
Republic EZ Broadcast	10	14*	10	14
Republic EZ Tractor Broadcast	10	14*	10	14
[Other Spreaders As Appropriate]	TBD	TBD	TBD	TBD

\*These numbers indicate a criss-cross application will be required to apply the number of pounds shown for 1,000 sq.ft.

Example:



Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

Sevin® is a registered trademark of Rhone-Poulenc Nederland BV for Carbaryl Insecticide