

67572-82

1-07-2005

1/8

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060.

United States Environmental Protection Agency 1200 Pennsylvania Avenue, N.W. Washington, DC 20460-0001	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	--	-----------------------

Application for Pesticide – Section I

1. Company/Product Number EPA Reg. No. 67572-82	2. EPA Product Manager Cynthia Giles-Parker	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/ Product (Name) CPI Chlorothalonil Multipurpose Fungicide	PM # 22	
5. Name and Address of Applicant (Include ZIP Code) Contract Packaging, Inc. c/o RegWest Company, LLC 30856 Rocky Road Greeley, CO 80631-9375 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section – II

<input type="checkbox"/> Amendment – Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification – Explain below.	<input type="checkbox"/> Other – Explain below.

NOTIFICATION
JAN 07 2005

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)


Submit notification of label revision per PR Notice 98-10.
(Refer to the Reference Label for an illustration of all revisions)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR § 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR § 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section – III

1. Material This Product Will Be Packaged In:			
Child Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify): _____
*Certification must be submitted		If "Yes" Unit Packaging Wgt. No. Per Container	If "Yes" Packaging Wgt. No. Per Container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling Accompanying Product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Stenciled <input type="checkbox"/> Paper glued <input type="checkbox"/> Other			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application).		
Name Kim Davis, CC, APC, Manager	Title Consultant/Agent	Telephone No. (Include Area Code) (970) 353-0611 / kim@regwest.com
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Star.iped)
2. Signature 	3. Title Consultant/Agent	
4. Typed Name Kim Davis, CC, APC, Manager	5. Date December 8, 2004	

CPI Chlorothalonil Multipurpose Fungicide

Concentrate
For Residential Use

NOTIFICATION
JAN 07 2005

Only For Outdoor Use
Water Based

Makes up to 64 Gallons Diluted Spray {pint}
Makes up to 128 Gallons Diluted Spray {quart}

Controls Diseases on: Shrubs, Vegetables, Fruit Trees, Flowers and Shade Trees
Controls Diseases – Leaf Spots, Rust, Blights, Fruit Rots, Mildews, Scab, Molds, etc.
on Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees

Controls/Stops & Prevents over 130 [130+] Diseases on Vegetables, Fruits, Roses, Flowers, Shrubs
and Trees/Ornamentals
Controls/Prevents/Stops Garden Diseases

Active Ingredient:

Chlorothalonil (CAS No. 1897-45-6)	29.6%
Other Ingredients	70.4%
Total	100.0%

Contains 2.88 pounds Chlorothalonil per gallon

Peel open for use directions and further information.
Open resealable label for directions and precautions.
Pull here to open label. Press to reseal.

Keep Out of Reach of Children
WARNING
See back panel [booklet] for additional
precautionary statements.

EPA Reg. No. 67572-82

EPA Est. 67572-GA-1

Net [Contents]: _____

Manufactured by:
Contract Packaging, Inc.
10115 Highway 142 North
Covington, GA 30014
{website}

{Booklet}

Product Usage	
Controls/Prevents/ Stops Diseases	Leaf Spots, Rust, Blights, Fruit Rots, Mildews, Scab, Molds {or select others from use directions}
Use On	Vegetables, fruits, roses, flowers, shrubs, trees and ornamentals {or select others from use directions}
Dilution Rates	Vegetables (depending on crop/disease): 1½ tsp to 1 Tbs per gallon of water Fruits (depending on fruit/disease): ¾ tsp to 2¼ tsp per gallon of water

	Ornamentals: 2¼ tsp to 3 Tbs per gallon of water Conifers: 1½ tsp to 3 Tbs per gallon of water
--	---

Ornamental Diseases

Cercospora, Phytophthora Die-back, Ovulinia Flower Blight, Leaf Blotch, Scab, Anthracnose, Fabraea Leaf Spot, Rhizoctonia Web Blight, Powdery Mildew, Rust, Botrytis, Alternaria Leaf Spots, Branch Rot, Septoria Leaf Spot, Black Spot, Helminthosporium Leaf Spot, Botrytis Gray Mold (foliar phase), Phytophthora Blight, Ascochyta Blight, Cylindrosporium Leaf Spots, Cercosporidium, Marssonina Leaf Spot, Fabraea (*Entomosporium*) Leaf Spot, Taphrina Blister, Actinopelte Leaf Spot, Monilinia Blossom/Branch Blight & many more listed inside.

Vegetable Diseases

Rust, Botrytis Blight (Gray Mold), Alternaria Leaf Spot, Downy mildew, Ring Spot, Cercospora (Early) Blight, Alternaria (Late) Blight, Septoria (Late) Blight, Basal Stalk rot, Pink Rot (Suppression), Helminthosporium Leaf Blight, Anthracnose, Target Spot, Scab, Purple Blotch, Botrytis Vine Rot, Gray Leaf Spot, Gray Leaf Mold, Rhizoctonia Fruit Rot, Gummy Stem Blight (Black Rot), Botrytis Leaf Blight (Blast), Cercospora Leaf Spot & many more listed inside.

Fruit Diseases

Peach Leaf Curl, Coryneum Blight (Shothole), Cherry Leaf Spot, Brown Rot, Blossom Blight, Scab & more listed inside.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[Shake Well Before Using]

Read Entire Label and Use Strictly in Accordance with Label Precautionary Statements and Directions.

tsp = teaspoon Tbs = Tablespoon

Readily mixes with water and can be used in a hand-held, hose-end or tank-type sprayer. Shake well before using. Fine pray required for proper coverage; use a hose-end sprayer for best results.

Use of this product on home lawns is prohibited. Do not apply this product in a way that will contact other persons or pets, either directly or through drift. Keep people and pets off treated areas until dry.

Vegetable Diseases

Apply the amount of concentrate in the table below in one gallon of water to cover 200 sq ft (10' x 20'). Thoroughly spray to cover all plant surfaces. **Note:** Avoid spraying plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to animals.

Crop	Diseases Controlled	Amount of Concentrate	Application Notes	PHI*
Beans (Green)	Rust, Botrytis Blight (Gray Mold)	1 Tbs	Start spraying at first sign of disease. Apply every 7 days to maintain control.	7
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Alternaria Leaf Spot, Downy Mildew	1½ tsp	Start spraying when conditions favor disease, after transplanting or when sprouts appear. Apply every 7 to 10 days.	0
Brussels Sprouts (only in CA)	Ring Spot	2 tsp		
Carrots	Cercospora (Early) Blight, Alternaria (Late) Blight	2 tsp	Start spraying at first sign of disease. Apply every 7 to 10 days to maintain control	0
Celery	Cercospora (Early) Blight, Septoria (Late) Blight	1½ tsp	Start spraying after transplanting. Apply every 7 days.	7
	Basal Stalk Rot, Pink Rot (suppression)	1 Tbs		
Corn (Sweet)	Helminthosporium Leaf Blight, Rust	2 tsp	Start spraying when conditions favor disease. Apply every 7 to 10 days to maintain control. Do not apply to corn if you plan to process it.	14

Crop	Diseases Controlled	Amount of Concentrate	Application Notes	PHI*
Cucumbers, Melons, Pumpkins, Squash	Anthracnose, Downy Mildew, Target Spot	2 tsp	Start spraying when plants are in the first true leaf stage or when conditions favor disease. Apply every 7 days to maintain control.	0
	Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states)	1 Tbs		
Cucumbers	Fruit Rot (suppression)			
Onion (Green), Leek, Shallot	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	1 Tbs	Start spraying at first sign of disease. Apply every 7 to 10 days. Do not use more than 3 times per season.	14
Onion (Dry Bulb), Garlic			Start spraying at first sign of disease. Apply every 7 to 10 days.	7
Potato	Early Blight, Late Blight, Botrytis Vine Rot	1½ tsp	Start spraying when plants are 6 inches high or at first sign of disease. Apply every 5 days to maintain control.	0
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot, Botrytis Gray Mold	1 Tbs	Start spraying when conditions favor disease. Apply every 7 to 10 days to maintain control	0

*PHI = the days required between last application and harvest.

Fruit Diseases

Apply the amount of concentrate in the table below in one gallon of water. Thoroughly spray to cover all plant surfaces.

Fruit Tree	Diseases Controlled	Amount of Concentrate	Application Notes
Peach, Nectarine, Apricot	Peach Leaf Curl, Coryneum Blight (Shothole)	2¼ tsp	Do not apply between fruit formation and harvest. Spray at leaf fall in late autumn. Reapply 1 to 2 more times in mid- to late winter before buds begin to swell. For Shothole, to prevent fruit infections, also apply 1 to 2 weeks after petal fall when fruit just begins to form.
	Brown Rot, Blossom Blight, Scab		Do not apply between fruit formation and harvest. Spray when buds begin to show color. Reapply at full bloom. If cool, moist weather continues, make a third application at petal fall. Apply once more when fruit just begins to form.
Cherry, Plum, Prune	Brown Rot Blossom Blight, Cherry Leaf Spot (cherries)	2¼ tsp	Do not apply between fruit formation and harvest. Spray when buds begin to show color. Reapply at full bloom. If cool, moist weather continues, make a third application at petal fall. For Cherry Leaf Spot, also apply when fruit just begins to form and again 7 days after fruit is removed; apply once more 10 to 14 days later, if tree has a history of leaf spot.
Papaya	Alternaria Fruit Spot, Anthracnose, Stem End Rot	1½ tsp	Start spraying when fruit spots appear (April to July). Spray every 14 days until dry weather returns.
Passion Fruit (only in HI)	Alternaria Fruit and Leaf Spot (Passion Fruit Brown Spot)	¾ tsp	Start spraying before spots appear on fruit (April to July). Spray every 14 days until dry weather returns.

Ornamental Diseases

Unless otherwise directed in the table below, mix 2¼ tsp per gallon of water or 3 Tbs (1½ fl oz) per 4 gallons of water and spray foliage to point of runoff. Begin applications as directed for each species and disease condition cited and apply every 7 to 14 days (unless otherwise specified) until conditions no longer favor disease. During periods when conditions favor severe disease, generally cloudy or wet weather, use the shorter interval between applications.

Species	Diseases Controlled	Suggested First Application
[Broadleaf Shrubs and Trees]		
Ash	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring bud break
Azalea*, Rhododendron*	Phytophthora Die-Back, Ovinia Flower Blight	New leaf emergence; early bloom
Buckeye (Horsechestnut)	Leaf Blotch, Anthracnose	Spring bud break
Crabapple	Scab, Cedar-apple Rust, Sphaeropsis Leaf Spot	Spring bud break
Dogwood	Anthracnose (<i>Discula</i> sp.), Septoria Leaf Spot	Spring bud swell; early bloom
Euonymus	Anthracnose	Spring bud break
Firethorn (Pyracantha)	Scab	Spring bud break
Flowering Almond, Quince and Cherry Hawthorn	Monilinia Blossom/Branch Blight, Rust, Fabraea Leaf Spot	Early bloom; pre-bloom
Holly	Rhizoctonia Web Blight	Warm, moist conditions
Mountain Laurel	Cercospora Leaf Spot	Spring bud break
Oak (only red group)	Taphrina Blister, Actinopelte Leaf Spot, Anthracnose	Dormant bud swell
Oregon-Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea (<i>Entomosporium</i>) Leaf Spot	Spring bud break
Pieris (Andromeda)	Phytophthora Die-Back	New leaf emergence
Poplar	Marssonina Leaf Spot	Spring bud break
Privet	Cercospora Leaf Spot	Prolonged wet conditions
Sycamore (Planetree)	Anthracnose	Spring bud break
Viburnum	Powdery Mildew	Mid-summer
[Bulbs and Flowering Plants]		
Carnation	Alternaria Leaf Spot, Branch Rot, Botrytis	Transplant of cuttings; cool, moist conditions
Chrysanthemum (Daisy)	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of cuttings
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-bloom; cool, moist conditions
Gladiolus	Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydranga* (only foliage)	Cercospora and Septoria Leaf Spots, Rust	Early propagation
Iris	Botrytis Blossom Blight, Didymellina Leaf Spot	Cool, moist conditions
Lily	Botrytis Gray Mold (foliar phase), Phytophthora Blight	Pre-bloom
Petunia*	Botrytis Gray Mold (foliar phase), Phytophthora Blight, Botrytis Blight	Pre-bloom
Rose – Use 1¼ fl oz (2½ Tbs) per 4 gallons of water	Black Spot, Botrytis Blight	Spring bud break
Statice	Anthracnose, Cercospora, Alternaria Leaf Spots, Botrytis Leaf Blight	Spring bud break
Zinnia	Powdery Mildew	First sign of disease
[Foliage Plants]		
Dracaena	Fusarium Leaf Spot	Pre-transplant
Pachysandra – Use 5 tsp per gallon of water	Volutella Leaf Blight	Spring bud break
Leatherleaf fern	Ascochyta Blight, Cercospora Leaf Spot, Cylindrocladium Leaf Spot, Rhizoctonia Blight	Spring bud break
Parlor Palm (Chamaedorea)	Bipolaris Leaf Spot (<i>Helminthosporium</i>)	Cool, moist conditions
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early propagation
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early propagation
Philodendron	Phytophthora Blight, Dactylaria Leaf Spot	Moist conditions
Synonium	Cephalosporium Leaf Spot	Warm, moist conditions

Discoloration of blooms has been noted on certain varieties when applications are made during flowering.

Conifer Diseases

Tree	Diseases Controlled	Amount of Concentrate	Application Notes
Pine	Scleroderris Canker, Lophodermium Needlecast	1½ tsp per gallon of water	Start spraying in spring when new shoots are ½ to 2 inches long. Spray every 3 to 4 weeks until dry weather returns. For Lophodermium Needlecast in Northcentral and Northeastern states: Start spraying in mid-July to early August before disease occurs; spray every 3 to 4 weeks until dry weather returns.
	Schirrhia Brown Spot	3 Tbs per gallon of water	
Douglas Fir	Needlecast, Rhabdocline, Swiss Needlecast, Lophodermium Needlecast	1½ tsp per gallon of water	
Spruce	Lophodermium Needlecast	1½ tsp per gallon of water	
	Rhizosphaera Needlecast	3 Tbs per gallon of water	

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool dry (preferably locked) place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: Causes eye irritation. May be a potential skin sensitizer. This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, long sleeve shirt, long pants and rubber gloves when mixing and applying this product. Do not breathe spray mist. Harmful if swallowed or absorbed through the skin. Thoroughly wash skin and hands after using. Remove and wash contaminated clothing before reuse.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [You may also contact {appropriate 800 number} for emergency medical treatment information.]	
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If Swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. This chemical can contaminate surface water through spray drift. Under some conditions it may also have high potential for runoff into surface water for several days to weeks after application; these include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely

shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips and areas overlaying tile draining systems that drain to surface water.

Buyer assumes all risk of use, storage or handling of this product not in strict accordance with this label.

{Code}

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347

If you have questions relating to product use or application, please call (800) 969-7200

If you have a product spill or transportation emergency call (800) 424-9300

For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200

[] Denotes optional/alternate language

{ } Denotes language that does not appear on the market label



December 8, 2004

Ms. Sherada Hobgood
Document Processing Desk – **NOTIF**
Office of Pesticide Programs – **7504C**
U.S. Environmental Protection Agency
1801 South Bell Street
Crystal Mall #2, Room 266A
Arlington, VA 22202-4501

Dear Ms. Hobgood:

Subject: Contract Packaging, Inc.
CPI Chlorothalonil Multipurpose Fungicide
EPA Reg. No. 67572-82

On behalf of its client **Contract Packaging, Inc.**, RegWest Company, LLC is submitting the attached materials as notification of label revision for subject product. RegWest Company will act as sole agent in this endeavor.

Attached are the following:

1. Application for Pesticide: Other
2. One **Reference Label** (illustrates all changes)
3. Three **New Labels** (incorporating all changes)

Please refer to the **Reference Label** which illustrates all the changes. We are primarily adding selling copy (taken directly from the use directions) and consumer contact numbers.

This product is not sold only by Contract Packaging's subregistrants; therefore no market labeling is attached.

Please contact me at (970) 353-0611 or kim@regwest.com if you have any questions concerning this submission or require additional information.

Regards,
RegWest Company, LLC

Kim Davis, CC, APC, Manager
Consultant/Agent

Attachments

pc: CPI – Hacker, S.

30856 Rocky Road
Greeley, CO 80631-9375
970-353-0611
970-353-0613 Fax
Kim@regwest.com