

1818  
1-20-85  
B  
181

# DRYACIDE

DIATOMACEOUS EARTH  
INSECTICIDE

FOR THE PROTECTION OF STORED GRAIN,  
THE DISINFESTATION OF GRAIN STORAGE FACILITIES AND  
GRAIN HANDLING MACHINERY.

CONTROLS ALL STORED-GRAIN INSECT PESTS INCLUDING  
FLOUR BEETLES, GRAIN BORERS, GRANARY WEEVILS, RICE WEEVILS,  
GRAIN BEETLES AND WAREHOUSE MOTHS

ACTIVE INGREDIENT	
Silicon Dioxide, from Diatomaceous Earth	97 %
INERT INGREDIENTS	
Silica Gel	3 %
Total	<u>100 %</u>

**\*\* CAUTION \*\***  
--- KEEP OUT OF REACH OF CHILDREN ---

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes. Use adequate ventilation and avoid breathing dust. Wear a suitable dust mask when using this product during prolonged periods.

#### STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with water. If irritation develops consult a physician.

NET WEIGHT OF THIS CONTAINER 15 kg or 33 pounds


## DRYACIDE

Imported and Distributed by:

EPA Registration No. 57595-1

EPA Establishment No. 67595

DRYACIDE USA  
3536 Emerson Street  
San Diego, CA 92106  
Telephone 619-222-1680



# DRYACIDE

## STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

STORAGE: Store in the closed original container in a dry, well-ventilated area and out of direct sunlight.

PESTICIDE Wastes resulting from the use of this product may be disposed of on site or at DISPOSAL: an approved waste disposal facility.

CONTAINER Completely empty bag into application equipment. Then dispose of empty DISPOSAL: bag in a sanitary landfill or by incineration or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

### DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DRYACIDE is intended for application with hand or power duster, pickle applicator or other suitable means to places where insects are found. Suggestions regarding the application of the material are listed below:

SITES	RATE	COMMENTS
Stored Grain	2 lb/ton	Apply to grain as it is transported to the storage location in a manner to give a uniform distribution throughout the grain. Treat grain immediately after harvest.
-----		
Grain Silos and Storage areas		
1 Dust	0.4 lb per 1000 sq Ft	Distribute dust to give an even coating on interior surfaces of storage areas paying particular attention to cracks and crevices where debris might remain. A hand operated duster may be used but for silos and large storage areas a power operated duster is recommended. For vertical silos, apply from TOP manholes or access points. (Dust downward into silo, not upwards from the base.)
2 Slurry	1.25 lb per 1000 sq ft (Dry basis)	Spray as a 10 to 20 percent slurry using a pressure of at least 40-60 psi using a centrifugal pump or venturi type sand blaster and flat, fan nozzle of at least 1 3 GPM capacity. Use a tank mixer or recirculate slurry through slurry tank throughout operation to maintain good slurry dispersion. Rinse nozzle, lines, and equipment with clean water after use.
-----		
Grain Handling Machinery, eg Field Bins, Augers, Seeders, Conveyors and Elevators	0.4 lb per 1000 sq Ft	Before applying, thoroughly clean all debris from machine. Use a vacuum cleaner or compressed air if available. Dust throughout machine, ensuring distribution to hidden or relatively inaccessible parts. Clean out excess before using machinery.

Restriction: Do not treat grain with a moisture content greater than 12% of in any operation where water is added to grain.

181