

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 67603-1	Date of Issuance: APR 16 1997
	Term of Issuance: Conditional	
	Name of Pesticide Product: TSD Multi-Purpose Insect Spray	
Name and Address of Registrant (include ZIP Code): Sherwin Williams Company 31500 Solon Road Solon, MA 44139		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. Make the following label changes: <ol style="list-style-type: none"> Revise the EPA Registration Number to read, "EPA Reg. No. ". 67603-1 Revise the Environmental Hazard statement as follows: "This product is toxic to fish and other organisms. Do not apply directly to water." 		
Signature of Approving Official:		Date:

c. Since you propose to use this product in industrial areas which could include commercial food handling establishments revise the paragraph beginning "Nonfood areas of food/feed handling xxx etc" to read "Do not use in the food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas. For use only in the non food/feed areas of these establishments such as (garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning and bottling).".

d. On the front panel add "Not for use in USDA Meat and Poultry Plants".

e. Qualify the heading Fleas and Ticks to read "Fleas and Ticks (Pet Premises)".

f. Revise the front panel statements to read "Control Named Residential, Industrial, Livestock and Garden insects".

g. Revise your label in accordance with PR Notice 96-6 concerning pet pesticide product label statements.

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

PERMANONE TSD Multi-Purpose Insect Spray
MULTI-PURPOSE 0.25% SPRAY

- Contains the Synthetic Pyrethroid PERMANONE
- Controls Named Residential, Industrial and Garden Insects
- For Use On Dogs And On Premises To Control Fleas And Ticks
- Controls Fleas And Ticks On Dogs For 14 Days
- A Water-Based Aerosol
- Controls Pests For Up To Four Weeks

-Not For Use On Humans

ACTIVE INGREDIENT:

*Permethrin.....0.25%

INERT INGREDIENTS:.....99.75%
100.00%

(3-Phenoxyphenyl)methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate (Cis/Trans ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.)

PERMANONE - Registered Trademark of Roussel Uclaf Corporation

EPA REG. NO. ~~4816-714~~ ⁶⁷⁶⁰³⁻¹ EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF IN EYES: Flush with plenty of water. Consult a physician if irritation persists.

IF ON SKIN: Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

See Side Panel For Additional
Precautionary Statements

NET CONTENTS:

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 16 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

67603-1

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION

Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin Wash hands thoroughly after use. Vacate room after treatment ant ventilate before reoccupying. Avoid contamination of feed and foodstuffs Cover or remove fishbowls. This product is not for use on humans.

Animals: Do not spray directly in/on eyes, mouth or genitalia of pets. Do not treat or cause exposure to puppies less than four weeks old. **DO NOT USE ON CATS.**

Environmental Hazards

This product is toxic to fish, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

Physical Or Chemical Hazards

Contents under pressure Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130.F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in a cool dry area. Do not transport or store below 32°F.

Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Nonfood areas of food/feed handling establishments are areas such as garage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Remove buds and cover fish aquariums before spraying.

Houseflies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion Repeat treatment as necessary.

Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around chins and plumbing, hitting insects with spray when ever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space Keep area closed for 15 minutes following treatment. Repeat application as necessary.

Exposed stages of *Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites:* Spray into cracks and crevices, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards moldings and floors. Repeat treatment as necessary. Allow all sprayed articles to dry thoroughly before use.

To control *Clothes Moths* and *Carpet Beetles* on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for seven hours in sunlight, if possible. Apply spray liberally to empty cheers, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spaying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of *House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small Flying Moths*, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results,

keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

TO TREAT LIVESTOCK To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of *Stable Flies*, *Horse Flies*, *Deer Flies*, *Face Flies*, *House Flies*, *Horn Flies*, *Mosquitoes and Gnats*: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from *Face Flies*, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

FOR SWINE Do not ship animals for slaughter within 5 days of last treatment.

Fleas and Ticks: (Adults and Larvae: Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place. *Wore use.*

To control *Fleas, Ticks* on dogs. Starting at the tail, stroke against the lay of the hair, spraying the parted hair in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application weekly if necessary to control infestation. **NOTE:** Not Recommended for use on puppies 1 ~ than four weeks old.

FOR THE CONTROL OF GARDEN INSECTS; For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* kill by spray, as well as *Whiteflies*, *Aphids*, *exposed Thrips*, *Spiders*, *Red Mites* and *Leafminers*.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums; dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against *Whiteflies*, *Aphids*, *Spider and Red Mites*, *exposed Thrips*, *Lace Bugs*, *Japanese Beetles*, *Pavement Ants*, *Clover Mites* and *Army Worms*. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 P. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

FOR RESIDENTIAL, COMMERCIAL, INDUSTRIAL USE