

7/7/97

PM 13

67603-1

pg 125

# TSD MULTI-PURPOSE INSECT SPRAY

- Contains the Synthetic Pyrethroid PERMANONE
- Controls Named Residential, Industrial, Livestock and Garden Insects
- For Use On Dogs And On Premises To Control Fleas And Ticks
- Controls Fleas And Ticks On Dogs For 14 Days
- A Water-Based Aerosol
- Controls Pests For Up To Four Weeks
- Kills Lice And Their Eggs
- For Use On Bedding, Furniture And Other Inanimate Objects Infested With Lice
- Not For Use On Humans
- Not For Use In Federally Inspected Meat and Poultry Plants

## ACTIVE INGREDIENT:

\*Permethrin.....0.25%

INERT INGREDIENTS:.....99.75%

100.00%

(3-Phenoxyphenyl)methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate (Cis/Trans ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.)

PERMANONE - Registered Trademark of AgrEvo Environmental Health, Inc.

EPA REG. NO. 67603-1    EPA EST. NO. 10900-OH-1, 10900-CA-1  
33915-MI-1, 257-MD-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**FIRST AID**

**IF INHALED:** Remove affected person to fresh air. Apply artificial respiration if indicated.

**IF IN EYES:** Flush with plenty of water. Consult a physician if irritation persists.

**IF ON SKIN:** Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

See Side Panel For Additional  
First Aid Statements

NET CONTENTS:

<b>ACCEPTED</b>
JUL - 7 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67603-1

**FIRST AID**

**Hazards To Humans & Domestic Animals**

**CAUTION**

Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin Wash hands thoroughly after use. Vacate room after treatment ant ventilate before reoccupying. Avoid contamination of feed and foodstuffs Cover or remove fishbowls. **This product is not for use on humans.**

**Animals:** Do not spray directly in/on eyes, mouth or genitalia of pets. Do not treat or cause exposure to puppies less than twelve weeks old. **DO NOT USE ON CATS.**

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other organisms. Do not apply directly to water.

**PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

**READ ENTIRE LABEL BEFORE EACH USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Store in a cool dry area. Do not transport or store below 32°F. Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

**NOTE**

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Do not use in the food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in

serving areas. For use only in the non food/feed areas of these establishments such as (garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning and bottling).

### **SHAKE WELL BEFORE USING**

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying.

***Houseflies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths:*** Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

***Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs:*** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Repeat treatment as necessary.

***Ants:*** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

***Indian Meal Moths, Angoumois Grain Moths:*** Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

***Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites:*** Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

***Carpet Beetles:*** Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

***Bedbugs*** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all sprayed articles to dry thoroughly before reuse. Repeat treatment as necessary.

To control ***Clothes Moths and Carpet Beetles*** on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles

may be treated by spaying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

**In Animal Quarters And Milkrooms:** For rapid control of *House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small Flying Moths*, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

**To Treat Livestock:** To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of *Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes and Gnats*: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from *Face Flies*, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

**For Swine:** Do not ship animals for slaughter within 5 days of last treatment.

**To Control Blood Sucking Lice:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**To Control Poultry Lice:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

**For Control Of Bedbugs And Mites In Poultry Houses:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**Fleas and Ticks: (Pet Premises) Adults and Larvae:** Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

**USE ONLY ON DOGS**

**To control Fleas and Ticks on Dogs:** Starting at the tail, stroke against the lay of the hair, spraying the parted hair in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat every 14 days if necessary to control infestation. NOTE: Do Not Use On Puppies Under 12 Weeks Old. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or on animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**To Kill Lice And Louse Eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only

those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

**For The Control Of Garden Insects:** For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray, as well as *Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers*.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums; dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against *Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms*. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

**FOR RESIDENTIAL, COMMERCIAL, INDUSTRIAL USE**

Sherwin-Williams Diversified Brands, Inc.  
Solon, OH 44139

For Emergency Health Information Call:  
1-216-566-2917