

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO. 67619-1

DATE OF ISSUANCE

APR 28 1998

TERM OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION
 REREGRISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

CPPC Bleach

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

The Clorox Professional Products Company
C/o PS & RC
P. O. Box 493
Pleasanton, Ca 94566

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.


This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants to similar products to submit such data.
2. Add the phrase "EPA Registration No. 67619-1" to your label before you release the product for shipment.

Submit one copy of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

A stamped copy of the label is enclosed for your records.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.


Robert S. Brennis
Product Manager (32)
Antimicrobial Program Branch
Regulatory Division (7504C)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

2 pte 26

(57619-03) CPPC BLEACH (1 of 24)

CPPPC BLEACH

ACTIVE INGREDIENT:
 SODIUM HYPOCHLORITE 5.25%
 INERT INGREDIENTS: 94.75%
 [TOTAL 100.00%]

Contains No Phosphates -or- Phosphorus
 (Yields -or- contains 5% available chlorine)


ACCEPTED
 APR 28 1998
 Under the Federal Insecticide, Fungicide, and
 Rodenticide Act, as amended, for the
 pesticide registered under
 EPA Reg. No. 67619-1

KEEP OUT OF REACH OF CHILDREN
WARNING: (See back panel for other cautions)

ADVERTENCIA: (Véase el panel posterior
 para otras precauciones)

NET CONTENTS:

for front or back
• unbeatable whitening



[* For Use Only In The PowerPack™ cleaning system]

**GERMICIDAL APPLICATIONS
ESPECIALLY RECOMMENDED FOR FOOD SERVICE APPLICATIONS**

- REMOVES STAINS
- CLEANS
- DISINFECTS
- SANITIZES
- DEODORIZES

MANTENGA ESTE PRODUCTO FUERA DEL ALCANCE DE LOS NIÑOS

COMMERCIAL SOLUTIONS

- GERMICIDAL
- BACTERIOCIDE
- BACTERIOCIDAL
- VIRUCIDE
- VIRUCIDAL
- FUNGICIDE
- FUNGICIDAL
- STREPTOCIDAL
- STAPHYLOCIDAL
- PSEUDOMONACIDAL
- MILDEWCIDE
- MILDEWCIDAL
- TUBERCULOCIDE
- TUBERCULOCIDAL

ACCEPTED
 APR 28 1998
 Under the Federal Insecticide, Fungicide, and
 Rodenticide Act, as amended. For the
 pesticide registered under
 EPA Reg. No. 67619-1

CPPC BLEACH

For a cleaner, fresher laundry and household

This product, a germicide -and/or- broad spectrum disinfectant, is a 5.25% sodium hypochlorite solution containing 5% available chlorine by weight. This product can be used on hard non-porous surfaces in homes, hospitals, commercial, institutional, and eating establishments, pet kennels, and veterinary premises.

- Whitens, and removes stains -or- • Brightens whites and removes stains • Safe for most color-fast washables
- Boosts cold water cleaning power • Disinfects and deodorizes by killing most odor-causing germs and their odors.
- Whitens bleachable fabrics • Removes [Tough] Stains • Cleans

One Tablespoon of this product in a gallon of water is equivalent to 200 parts per million (ppm) available chlorine.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR SANITIZING

	Amount of this product	Amount water	Instructions
Work Surfaces	1 Tbsp 1/2 oz.	1 Gallon	Wash, rinse, wipe surface area with bleach solution for at least 1 minute. Let air dry.
Dishes, Glassware, Utensils	1 Tbsp 1/2 oz.	1 Gallon	Wash and rinse. After washing, soak for at least 1 minute in bleach solution, drain and air dry.
Refrigerators, Freezers	1 Tbsp 1/2 oz.	1 Gallon	Wash, rinse, wipe surface area with bleach solution for at least 1 minute. Let air dry.
Mops, Brushes, Brooms & Rags	3/4 Cup 6 oz.	1 Gallon	Pre-wash items, then soak them in bleach solution for at least 1 minute. Rinse well.

FOR DISINFECTING

Floors & Walls	3/4 Cup 6 oz.	1 Gallon	Pre-wash surface, mop or wipe with bleach solution. Allow solution to contact surface for at least 2 minutes. Rinse well.
Bathtubs & Showers	3/4 Cup 6 oz.	1 Gallon	Pre-wash surface [and] wipe with bleach solution. Allow solution to contact surface for at least 2 minutes. Rinse well.
Toilet Bowl	3/4 Cup 6 oz.	1 Gallon	Flush toilet. Pour product into bowl. Brush entire bowl including under rim with brush or mop. Let stand 2 minutes. Flush.
Laundry	1 Cup 8 oz.	Standard Washer	Use 1 1/2 cups bleach for extra large washers. Use a detergent.

FOR DEODORIZING

Garbage Cans	1/4 Cup 2 oz.	1 Gallon	After washing and rinsing, brush inside with bleach solution. Let drain.
Drains	1 Cup 8 oz.	—	Flush drains. Pour into drain. Flush with hot water.

FOR BLEACHING/WHITENING

Wooden Surfaces	3/4 Cup 6 oz.	1 Gallon	Apply for [at least] 2 minutes, rinse.

FOR MOLD, MILDEW & STAIN REMOVAL

All Surfaces	3/4 Cup 6 oz.	1 Gallon	Add bleach to [powdered] detergent solution. Apply, let stand 2 or 5-15 minutes. Wipe and rinse.

Una Cucharada del producto en un galón de agua es el equivalente de 200 partes por millón (ppm) de cloro disponible.

INSTRUCCIONES PARA EL USO

Es una violación de la ley federal usar este producto de una manera inconsistente con la designación de la etiqueta sellada.

PARA LA HIGIENE DE

	Cantidad del producto	Cantidad de Agua	Instrucciones
Áreas de trabajo	1 Cucharada 1/2 onza	1 Galón	Lave, enjuague, frote la superficie con una solución de blanqueador por lo menos 1 minuto, deje secar al aire.
Platos, vasos, utensilios	1 Cucharada 1/2 onza	1 Galón	Lave y enjuague. Después de lavarlos, sumérgalos en una solución de blanqueador por los menos 1 minuto, enjuagar y deje secar al aire.
Refrigeradores, congeladores	1 Cucharada 1/2 onza	1 Galón	Lave, enjuague, frote la superficie con una solución de blanqueador por lo menos 1 minuto, deje secar al aire.
Trapeadores, cepillos, escobas y trapos	3/4 de Taza 6 onza	1 Galón	Lave los artículos antes del tratamiento, sumérgalos en la solución de blanqueador por lo menos 1 minuto. Enjuague bien.

PARA DESINFECTAR

Pisos y paredes	3/4 de Taza 6 onza	1 Galón	Lave la superficie antes del tratamiento. Friegue o frote con la solución de blanqueador. Deje que la solución contacte la superficie por lo menos 2 minutos. Enjuague bien.
Bañeras, duchos	3/4 de Taza 6 onza	1 Galón	Lave la superficie antes del tratamiento [y] frote con la solución de blanqueador. Deje que la solución contacte la superficie por lo menos 2 minutos. Enjuague bien.
Tasa del baño	3/4 de Taza 6 onza	1 Galón	Jalar para inundar la cubeta del inodoro. Verter el producto en la cubeta. Cepillar toda la cubeta, incluso por debajo del reborde, con un cepillo o una aljofifa. Dejar actuar durante dos minutos y volver a jalar-para enjuagar.
Lavanderia	1 Taza 8 onza	Lavadora Standard	Use una y media taza de blanqueador para lavadas grandes. Use tambien un detergente.

PARA DESODORIZAR

Basureros	1/4 de Taza 2 onza	1 Galón	Después de lavarlos y enjuagarlos, cepillelos por dentro con la solución de blanqueador. Deje secar.
Desagues, vertederos, sumideros	1 Taza 8 onza	—	Enjuague. Vierta la solución en el desague. Luego deje fluir agua caliente.

PARA BLANQUEAR/DECOLORAR

Superficies de madera	3/4 de Taza 6 onza	1 Galón	Aplique la solución [por lo menos] -o- [durante] 2 minutos. Luego enjuague la superficie.

PARA QUITAR MOHO Y MANCHAS

En cualquier superficie	3/4 de Taza 6 onza	1 Galón	Agregue blanqueador a la solución detergente. Aplíquelo y deje en remojo 2, -o- 5-15 minutos. Enjuague y frote. Luego enjuague la superficie.

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

5026

(67619-xx) CPCC BLEACH (4 of 24)



HOUSEHOLD HINT
Disinfect Flower Pots and Planters
 Cleaning flower containers helps prevent the transfer of molds and diseases from old plants to new ones. Wash and [thoroughly] rinse pots and planters. Soak 2 minutes in a solution of 3/4 cup of this product to one gallon of water, then rinse.



HOUSEHOLD HINT
Sanitize Pet's Food and Water Bowls
 To sanitize pet food containers, wash bowls with detergent and rinse. Fill bowls with a solution of 1 Tablespoon of this product per gallon of water. Let stand 1 minute, drain and air dry.



HOUSEHOLD HINT
Deodorizing Cat's Litter Box
 Unpleasant cat box odors can be eliminated when this product is used to kill odor-causing germs. Wash litter box with sudsy water and rinse. Then wipe with a solution of 3/4 cup of this product per gallon of water. Let solution stand 2 minutes before rinsing thoroughly.



HOUSEHOLD HINT
Keep Wading Pools Sanitary
 This product is excellent for chlorinating wading pools. As a general rule, use 1/2 cup per 100 gallons of water. For example, an 8-foot diameter pool holding 1 foot of water would require 1/2 cup of this product. To chlorinate, mix required amount of this product with 2 gallons of water and scatter over surface of empty pool. Fill remainder of pool with water. Empty small pools daily. Stabilized pools should maintain a residual of 1.0 to 1.5ppm available chlorine. Test the pH, available chlorine residual and alkalinity of the water frequently with appropriate test kits. Frequency of water treatment will depend upon temperature and number of swimmers.



HOUSEHOLD HINT
Cleaning Mold and Mildew
 Mold and mildew in the bathroom can be removed easily and effectively using this product. Simply wipe down surfaces using a solution of 3/4 cup of this product to each gallon of warm water. Keep surface wet [at least] 2 minutes; then rinse thoroughly and wipe dry. Repeat, if necessary, on heavily soiled surfaces.



HOUSEHOLD HINT
Removing Patio Moss and Mildew Stains
 Patio moss and mildew stains can be unsightly, slippery and dangerous. Hose patio to remove loose debris. Then use this product to remove moss and mildew stains by washing the area with a solution of 3/4 cup of this product to 1 gallon of water. Reapply the solution as needed to keep the area wet for at least 2 minutes. Brush as needed to remove moss and then rinse thoroughly. [Do not use on painted wood.]



HOUSEHOLD HINT
Eliminating Garbage Can Odors
 This product can deodorize and sanitize your garbage cans by eliminating the bacteria that cause odors. Wash garbage cans with soapy water and rinse. Then to deodorize and sanitize, swish a solution of 3/4 cup of this product per gallon of water over the inside of the can. Let the solution stand 2 minutes before rinsing.



HOUSEHOLD HINT
Keep Cut Flowers Fresh Longer
 Fresh cut flowers will stay beautiful longer if you add 1/4 teaspoon of this product to each quart of cold water. This product can also be used to remove flower vase stains and odors. Wash the vase thoroughly and then fill with a solution of 3/4 cup of this product to one gallon water. Let stand 5 minutes before rinsing.



HOUSEHOLD HINT
Sanitizing Kitchen Cloths
 This product can help you deodorize and sanitize dishcloths and synthetic sponges while cleaning your sink at the same time. Fill sink with a gallon of water. Add 3/4 cup of this product. Soak kitchen cloths in solution for [at least] 2 minutes, then rinse sink and cloths. Allow to air dry.



HOUSEHOLD HINT
Eliminating Refrigerator Odors
 This product kills odor causing bacteria and leaves your refrigerator smelling fresh and clean. Use it inside and out. Wash surfaces with a solution of 3/4 cup this product per gallon of water. Let stand 2 minutes. Rinse and then air dry interior surfaces a few minutes before replacing food.




HOUSEHOLD HINT
Sanitizing Baby Items
 Baby bottles, nipples and dishes can be easily sanitized using this product. Soak washed items for 1 minute in a solution of 1 Tablespoon this product per gallon of water. Pour solution through nipples; rinse and air dry.



HOUSEHOLD HINT
Christmas Trees Fresher Longer
 To prolong the life of a fresh cut tree, instead of using plain water in the tree stand bowl, use a solution of 2 teaspoons of this product, 1/2 gallon hot water, 1 cup corn syrup and 1/2 cup powdered chelated iron (available from local nurseries).



HOUSEHOLD HINT
Removing Exterior Mold
 Mold [growing] on washable and colorfast exterior surfaces of your home, like siding, tile roofs, brick, stucco and patio stone, can be easily removed using this product. First, hose surfaces to remove loose soil. Then apply a solution of 3/4 cup of this product per 1 gallon of water to wet surfaces. Reapply the solution as needed to keep the area wet for at least 2 minutes. Rinse thoroughly to remove residue. [Avoid applying solution in direct sunlight or to unfinished wood.] Rinse quickly and thoroughly if solution comes in contact with aluminum window frames or gutters since metal corrosion may occur.



HOUSEHOLD HINT
Sanitize and Remove Stains from Kitchenware
 Tough stains can be removed from dinnerware, dishes, plastic and glassware with this product. Plus, this product cleans and sanitizes. Wash items; then rinse, and soak for 2 minutes in a solution of 1 Tablespoon of this product to each gallon of water. Then drain and air dry.



HOUSEHOLD HINT
Disinfecting Baby Furniture and Toys
 Painted and enameled cribs, changing tables and high chairs, plastic mattress covers and bumpers, and washable colorfast toys are disinfected quickly and easily with this product. Plus, this product kills 99.9% of all common household germs, including those that cause odors. This product leaves baby's room clean and fresh smelling. Disinfect with a solution of 3/4 cup of this product in one gallon of water. Let stand 2 minutes. Rinse and [air] dry.



HOUSEHOLD HINT
Sanitize Wooden Cutting Boards
 Wash counter or cutting board with hot, sudsy water. Rinse. Then to sanitize surface thoroughly with solution of 3 Tablespoons of this product per 1 gallon of water. Keep surface wet for at least 2 minutes. Rinse then drain and air dry.

ACCEPTED
 APR 28 1998
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

R-10233-1

10026

(57619-xx) CPFC BLEACH (p of 24)

HOUSEHOLD USE

WHERE TO USE: -or- WHERE DO I USE THIS PRODUCT?

HOUSEHOLD USE DIRECTIONS FOR DISINFECTION: [Pre-clean to remove heavy soils.] Use 1/4 cup of this product in -or- per one gallon of water. Wash, or wipe items with the [cleaning] solution [then rinse]. For difficult stains, toilet bowls or to disinfect, pre-clean to remove heavy soils. Apply disinfecting -or- bleach solution. Let stand [for] 2 minutes [before rinsing]. Rinse [thoroughly] [and air] dry.

KITCHEN: Refrigerators, sinks, appliances, plastic laminate, countertops, tile, vinyl, linoleum.

Kitchen Sinks: Cover stains with 2/3 gallon of water before adding 1/2 cup of this product. -or- Cover stains with 2/3 gallon of water; then add 1/2 cup of this product. Let stand 2 minutes before rinsing.

BATHROOM: Bathtubs, toilets, showers, sink, countertop, glazed porcelain, cat litter boxes, and mold/mildew removal. Bathtubs and showers, floors, vinyl, tile, woodwork and appliances: Flush drain, if applicable. Clean with a solution of 3/4 cup of this product per gallon of [warm] water. Let stand 2 minutes before rinsing.


TOILET BOWLS: Flush toilet to remove heavy soil. Pour 3/4 -or- 1 cup of this product into bowl. Brush entire bowl including under the rim with a scrub brush or mop. Let stand 2 minutes before flushing again.

BABY'S NURSERY: Toys, changing tables, painted cribs, high chairs, plastic mattress covers and bumpers.

OUTDOORS: Use to remove -or- Removes mold, moss, and mildew [stains] from [outdoor] siding, tile, brick, stucco, and patios. Also use on flower pots and planters.

HOUSEHOLD USE DIRECTIONS FOR SANITIZATION:

GARBAGE CANS/DIAPER PAILS: Pre-clean garbage can/diaper pail with a cleaning product prior to sanitization. Rinse with water and drain. Pour in 3/4 cup of this product to each gallon of water. Mix. Let stand [at least] 2 minutes. Rinse and air dry.



LAUNDRY USE

Sort laundry by color and fabric. For the cleanest, whitest whites, use this product in every bleachable load. Most white fabrics and some colored fabrics can be safely washed with this product. In fact, many fabrics that have "Do not bleach" labels are bleach safe.

-or-

Sort laundry and fill washer with water. Add 1 cup of this product. Use 1 1/2 cups for extra large or heavily soiled loads.* Add detergent then laundry.

*Use bleach dispenser if available.

If unsure about colorfastness of garment or if garment says, "No bleach" or "Only non-chlorine bleach": Mix solution of 1 Tablespoon of this product with 1/4 cup water. Apply a drop of solution to a hidden part of garment. (Be sure to check all colors.) Wait one minute and blot dry.

-or-

If uncertain about colorfast dye, test fabric by applying one drop of a solution made of 1 Tablespoon of this product plus 1/4 cup water to hidden part of seam. Be sure to check all colors. After 1 minute, blot dry.

No color change means [the] article can be safely bleached. [Always] avoid bleaching spandex, wool, [mohair], silk, and non-fast colors.

For Best Laundry Results:

- Dilute 1 cup of this product in 1 quart of cool water.
- Add to wash 5 minutes after wash cycle has begun.

To Pretreat Stains:

- Mix 1/4 cup of this product with one gallon cold water.
- Soak garment for 5 minutes.
- Rinse, then wash as usual.

To Handwash:

- Mix 1/4 cup of this product with 2 gallons water.
- Soak garment for 5 minutes.
- Rinse, then wash as usual.

Do not use full strength to pretreat.

-or-

To handwash, [disinfect] or pretreat stains [and heavy soils], rinse to remove loose soil and fully soak garment for 5 minutes in a solution of 1/4 cup of this product to 1 gallon of cool -or- cold water.

Add this product to dispenser, if available. If not, add bleach and detergent with the wash water before the laundry is put in. Or for best results, dilute this product with a quart of water and add to wash 5 minutes after the wash cycle has begun.

To Disinfect:

- add 1/4 cup of this product to a gallon of water.
- add 1 cup to a standard size washing machine.



for use with PowerPack™ cleaning system:

System Set-Up:

1. Fill water bottle with cool, clean water. Remove cap from bottle.
2. With the water bottle on the left hand side, stand both bottles together so that the flat sides touch each other.
3. Rotate the white tabs on the bottom of the sprayer clockwise to the open position.
4. With the logo on the sprayer facing you, mount the sprayer onto both bottles insuring that the long plastic tube attached to the sprayer is inserted into the water bottle.
5. Lock the bottles onto sprayer by rotating the white tabs counterclockwise to the locked position.
6. Adjust the sprayer nozzle [tip] to either spray or stream and squeeze the trigger 15 times, only on initial installation to insure the proper dilution (2400 ppm available chlorine) will be used. (This product is now ready for use.)

APR 28 1998
Under a Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

7/25/96

(57619-xx) CPPC BLEACH (6 of 24)

How to Sanitize and Disinfect with this product Como Hacer Saneamiento y Desinfectar con este producto

This product is an effective multi-purpose sanitizer/disinfectant that kills bacteria that may cause food poisoning. This product is authorized by USDA for use in federally inspected meat and poultry plants. One tablespoon of

This product in a gallon of water is equivalent to 200 ppm available chlorine. One tablespoon of this product in a gallon of water meets U.S. Public Health Service recommended levels of hard surface sanitation.

To Sanitize
Pots & Pans • Glasses, Dishes, Utensils

DIP 1 MINUTE
SUMERJA 1 MINUTO

WASH LIME RINSE ENJUAGUE IMMERSE IN this product SUMERJA EN EL este producto DRAIN DRY DRENE HASTA SECAR

To Disinfect
Floors & Walls

PRE-WASH SURFACE TO BE TREATED LAVE LA SUPERFICIE DE TRATAMIENTO

2 1/2 CUPS 2 1/2 TAZAS of this product

3 GAL 3 GALONES WARM WATER CALDA AGUA

MOP FOR 2 MINUTES ALMOFRE 2 MINUTOS

RINSE & LET DRY ENJUAGUE Y DEJE SECAR

Food Contact Surfaces

CLEAN AND RINSE LAVE Y ENJUAGUE

1 1/2 CUPS this product

3 GAL 3 GALONES WATER AGUA

1 MINUTE 1 MINUTO

RINSE ENJUAGUE ALLOW TO DRY PERMITA SECAR

Toilets

SOAK 2 MINUTES EMPAPE 2 MINUTOS

1/2 - or - 1 CUP 1/2 - or - 1 TAZA of this product

FLUSH DEJE IR EL AGUA

BRUSH CEPILLE

FLUSH DEJE IR EL AGUA

Food Equipment

CLEAN AND RINSE LAVE Y ENJUAGUE

1 1/2 CUPS this product

3 GAL 3 GALONES WATER AGUA

1 MINUTE 1 MINUTO

WIPE FROTE EL REBANADOR

AIR DRY SEQUE AL AIRE

Refrigerators & Freezers

WASH LIME

WIPE WITH this product FROTE CON este producto

1 GAL 1 GALON WATER AGUA

1 MINUTE 1 MINUTO

DRY SOQUE

AIR DRY FOR 1 MINUTE SECAR AL AIRE POR 1 MINUTO

Mops, Brushes, Rags

5 GAL SINK FREGADERO DE 5 GALONES

2 GAL 2 GALONES 75° WATER AGUA

2 MINUTES 2 MINUTOS

WASH/RINSE LAVE/ENJUAGUE

SOAK EMPAPE

RINSE ENJUAGUE

AIR DRY SEQUE AL AIRE

For additional information regarding CPPC BLEACH training tools and dispensing systems, please call 1-800-685-9128 -or-

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 61619-1




80706

(67619-00) CPPG BLEACH (7 of 24)

How to Sanitize and Disinfect with this product


Como Hacer Saneamiento y Desinfectar con este producto

1 pump = 16 mL (0.54 oz)
1 bombeo = 16 mL (0.54 oz)

APPLICATIONS	AMOUNT/CANTIDAD	INSTRUCTIONS	INSTRUCCIONES
Sanitizing in the sink Saneamiento Del Fregadero	Bleach/ Blanqueador 8 Pumps 8 Bombeos 	Water/ Agua 10 Gallons 10 Galones • Add 8 pumps of bleach to 10 gallons of 65-75° F (room temperature) water in the sanitizing sink compartment • Check chlorine concentration with standard test strip Concentration should not exceed 200 ppm of available chlorine • Wash and rinse • Soak pots/pans/serving trays/utensils and food equipment for at least 1 minute in bleach solution • Drain & air dry	• Agregue 8 bombeos de blanqueador a 10 galones de agua a 65-75° F (temperatura del ambiente) en el compartimiento para saneamiento del fregadero. • Examine la concentración de cloro con una tira (banda) estándar de pruebas para cloro. La concentración no debe exceder 200 ppm de cloro disponible. • Lave y enjuague. • Remoje las ollas/cacerolas/charolas (bandejas) de servir/utensilios y equipo de comida por lo menos por 1 minuto en la solución de blanqueador. • Déjelos secar al aire.
Wipe Cloths Limpiones	Bleach/ Blanqueador 4 Pumps 4 Bombeos 	Water/ Agua 5 Gallons 5 Galones • Wash wipe cloths prior to sanitizing in a bleach solution for at least 5 minutes Store wipe cloths in bleach solution when not in use • Fill 5 gallon dispenser with 65-75° F (room temperature) water • Add 4 pumps of bleach • Draw bleach solution from dispenser as needed into well bowls, pans or spray bottles • Check concentration frequently with standard chlorine test strip Concentration should not exceed 200 ppm of available chlorine	• Antes de limpiar y hacer saneamiento con los limpiones, lave los limpiones y remójelos en una solución de blanqueador por lo menos por 5 minutos. Siempre mantenga los limpiones en una solución de blanqueador cuando no estén en uso. • Llene el dispensador de 5 galones con agua a 65-75° F (temperatura del ambiente). • Agregue 4 bombeos de blanqueador. • Use la solución de blanqueador del dispensador según se necesite poniéndola en los tazones, las cacerolas o las botellas de rociar. • Examine la concentración de cloro con una tira (banda) estándar de prueba para cloro. La concentración no debe exceder 200 ppm de cloro disponible.
Food Contact Surfaces (Beverage centers/prep-boards, holding wells/ service counters & tables/ refrigerators/ freezers) Superficies en Contacto con la Comida (Centros de bebidas/ mesas de preparación, fuentes de soporte/ mostradores y mesas de servicio/ refrigeradores/ congeladores) Stationary Equipment or other Food Contact Surfaces Equipo Estacionario u otras	Bleach/ Blanqueador 2 Pumps 2 Bombeos 	Water/ Agua 2 1/2 Gallons 2 1/2 Galones • Add 2 pumps of bleach to 2 1/2 gallon bucket of 65-75° F (room temperature) water • Check concentration frequently with chlorine test strip Concentration should not exceed 200 ppm of available chlorine • Wash, rinse and wipe surface with bleach solution • Wet all drink surfaces for [at least] 1 minute • Drain & air dry NOTE: During food preparation periods, wipe food contact surfaces frequently with sanitized wipe cloths. Store cloths in bleach solution when not in use.	• Agregue 2 bombeos de blanqueador en un balde de 2 1/2 galones de agua con una temperatura de 65-75° F (temperatura del ambiente). • Examine la concentración frecuentemente con la banda (tira) de prueba del cloro. La concentración no debe exceder 200 ppm de cloro disponible. • Lave, enjuague y limpie la superficie con la solución del blanqueador. • Moje las superficies de beber por lo menos por 1 minuto. • Déjelas secar al aire. NOTA: Durante los períodos de preparación de comida, limpie las superficies en contacto con la comida frecuentemente con limpiones saneados. Mantenga los limpiones en una solución de blanqueador cuando no estén en uso.

Fresh sanitizing solution should be prepared daily or more often if the solution becomes diluted or soiled.

La solución usada para sanitizar debe ser preparada diariamente ó más frecuente, si la solución se ve sucia ó si esta muy diluida.




9/2/26

(67619-xx) CPPC BLEACH (8 of 24)

How to Sanitize and Disinfect with this product (continued) Como Hacer Saneamiento y Desinfectar con este producto

**TOILETS AND URINALS
DISINFECTION**

**PARA DESINFECTAR
INDOROS Y ORINALES**



Lift up pump. Push down to dispense.
Levante la bomba. Apriete para dispensar.

Toilet

- Flush toilet to remove heavy soil
- Add 12 pumps of bleach to toilet bowl
- Brush to thoroughly wet all surfaces, including under the rim
- Let stand for 2 minutes, then flush

Inodoros

- Tire la cadena del inodoro para remover la suciedad
- Agregue 12 bombeos de blanqueador a la taza del inodoro
- Cepille para mojar completamente todas las superficies incluyendo debajo del borde
- Déjelo tranquilo durante 2 minutos, luego tire de la cadena.

Urinal

- Flush urinal to remove heavy soil
- Add 24 pumps of bleach
- Brush thoroughly for 1 minute
- Let stand for 2 minutes, then flush

Orinales

- Tire la cadena del orinale para remover la suciedad
- Agregue 24 bombeos de blanqueador
- Cepille completamente durante un minuto
- Déjelo tranquilo durante 2 minutos, luego tire de la cadena



**CPPC BLEACH (EPA Reg. No. 67619-xx) FOR FOOD EGG SANITATION
(USDA* Designation Q4)**

To sanitize good eggs: thoroughly clean all eggs. Completely mix 5.5 oz of this product with 10 gallons of warm water to produce a 200 ppm (parts per million) available chlorine solution. The sanitizer temperature should not exceed 130° F. Spray the warm sanitizer so that the eggs are completely wet. Allow the eggs to fully dry before carrying or breaking. Do not [apply a potable water] rinse. The solution should not be re-used to sanitize eggs.

**Measurement Table
for 200 ppm available chlorine**

1 Tbsp of this product	-to-	1 gallon water
5 1/2 oz of this product	-to-	10 gallons water
1 qt 20 oz of this product	-to-	100 gallons water
4 gallons of this product	-to-	1000 gallons water


*Authorized by USDA for use in federally inspected meat and poultry plants.

**CPPC BLEACH (EPA Reg. No. 67619-xx) FOR [MEAT & POULTRY PLANT] LAUNDRY USE
[(USDA* Designation B1)]**

This product may be used on fabric which contacts meat or poultry products directly or indirectly, provided that the fabric is thoroughly rinsed with potable water at the end of the laundering operation.

To disinfect laundry, add enough of this product to reach 200 ppm (parts per million) available chlorine. (Use 1 cup of this product per standard washer, 1 1/2 cup for extra large washers or heavily soiled loads). Use a detergent. For best results, dilute bleach with 2 quart of water and add to wash 5 minutes after the wash has begun.

*Authorized by USDA for use in federally inspected meat and poultry plants.



R-10233-1

10/2/98

67619-03 CPPC BLEACH (9 of 24)

CPPC BLEACH FOR DISINFECTION OF FLOORS, WALLS, SHOWERS AND TOILETS (USDA* Designations D2 & C2)

TO DISINFECT FLOORS, WALLS AND SHOWERS (USDA* Designation D2): For non-porous surfaces such as vinyl or ceramic tile, clean surfaces to remove gross filth. Wet surfaces thoroughly with a bleach solution containing 3/4 cup of this product per gallon of water. Allow solution to remain on the surface for 2 minutes. Rinse, let dry.

TO DISINFECT TOILETS (USDA* Designation C2): Flush toilet. Pour 3/4 -or- 1 cup of this product into bowl. Brush bowl thoroughly, making sure to get under the rim. Let solution stand for 2 minutes and flush again.

*Authorized by USDA for use in federally inspected meat and poultry plants.

CPPC BLEACH FOR DAIRY AND CREAMERY EQUIPMENT SANITATION

This product is effective as a chemical sanitizer of milk utensils, containers and equipment. This product dissolves milk solids and other protein material and is a quick and effective deodorizer.

An exposure period of at least 2 minutes to a solution of 2 oz of this product to 4 gallons water (200 p.p.m. available chlorine) should be maintained when the solution temperature is 75° F. Lower solution temperatures result in slower action; for each 18° F drop in temperature, approximately twice the exposure time is needed to achieve equivalent bactericidal action with same strength of solution. [You can also compensate for lower temperatures by increasing the concentration of this product.]

It is important to clean out large deposits of milk or other organic matter before applying bleach/water solution. A sharp decline in the available chlorine content of the solution following circulation through milk processing equipment is usually regarded as evidence of inadequate cleaning of the equipment and should be promptly investigated.

RUBBER TEAT CUPS AND TUBES - Before each milking, prepare sanitizing solution by thoroughly mixing 2 oz of this product with each 4 gallons water. This gives approximately 200 ppm available chlorine in an average 70° F solution. Sanitize teat cups with this solution for 2 minutes before transferring them from one cow to another.

To Sanitize - SOAKING METHOD: After each milking, wash cups and tubes by brushing thoroughly with detergent solution. Rinse cups and tubes with cold water. Prepare sanitizing solution by thoroughly mixing 2 oz of this product with each 4 gallons water in earthenware, glass, porcelain or stoneware containers. Submerge cups in this solution for 1 minute, holding ends of tubes; coil tubes slowly into solution between milkings; drain thoroughly and air dry before using.

To maintain sanitizing solution at proper strength, add 1/2 oz of this product daily (in hot weather, 1 oz) for each 3 gallons water; mix well. Protect solution from light. Renew solution daily. Old solution may be utilized for deodorizing and making floors and drains sanitary; for this purpose, add 1 oz of this product for each 5 gallons old solution; mix well.

To Sanitize - RACK METHOD: After each milking, rinse cups and tubes in cold water. Wash in detergent solution; then rinse. Prepare sanitizing solution by thoroughly mixing 1 oz of this product with each 2 gallons water; place solution in bottle above rack for 1 minute. Place tubes and cups in rack; fill with solution and let stand between milkings; drain thoroughly and air dry before using. Old solution may be utilized in deodorizing and making floors and drains sanitary.

METAL TEAT CUPS AND TUBES - Before each milking, prepare sanitizing solution by thoroughly mixing 2 oz of this product with each 4 gallons water. Dip teat cups into this solution before transferring them from one cow to another.

To Sanitize: After each milking, rinse cups and tubes with cold water. Wash in detergent solution; rinse in solution prepared by thoroughly mixing 2 oz of this product with each 4 gallons water for 1 minute; drain thoroughly and air dry before using. (Metal cups should not be left in bleach solution.)

TO CLEAN AND SANITIZE MILKING MACHINES AND UTENSILS: Immediately after milking, flush equipment with clean, lukewarm water. Dismantle equipment after each milking and wash it (including all rubber parts and stanchion hoses) and all utensils with a solution prepared by thoroughly mixing 2 oz of this product to each 4 gallons of your [regular] [powdered] detergent solution. Water temperature should be 100° F to 130° F. **DO NOT MIX THIS PRODUCT WITH ACID CLEANERS OR MILK STONE REMOVERS.** Rinse equipment and utensils thoroughly with clean, clear water; drain. Air dry. Immediately before use, sanitize according to directions at bottom of page.*

CLEANING IN PLACE - BULK STORAGE TANKS, DAIRY PIPELINES, TRANSFER STATIONS: Immediately after emptying milk, flush surfaces with a large volume of clear, lukewarm water until water runs completely clear. Thoroughly mix solution of 1 oz of this product with each 2 gallons of your [regular]

ACCEPTED
APR 28 1998
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 67619-1

R-10233-1

11 of 26

(57619-02) CPPC BLEACH (10 of 24)

[powdered] detergent solution. Hot water should be used if available, and the temperature of the solution should be maintained at 120-160° F throughout the entire circulation. (DO NOT USE THIS PRODUCT WITH ACID CLEANERS OR MILK STONE REMOVERS.) Circulate the sanitizing solution through the system for 10 to 15 minutes. (Brush-wash with solution all parts not coming in contact with solution as it circulates.) Rinse thoroughly with clean, clear water; allow to drain. Air dry. Seal this equipment to help protect against contamination. Immediately before use, sanitize according to directions at bottom of page.*

SEPARATORS, STRAINERS, MILK CANS, PAILS, CHURNS, PASTEURIZERS - to clean and sanitize: After using, rinse immediately with clear, cold water; then scrub or pressure-spray with solution of 1 oz of this product mixed with each 2 gallons of your [regular] [powdered] detergent solution. Rinse with clean, clear water; drain thoroughly. Air dry. Immediately before use, sanitize according to directions at bottom of page.*

MILK BOTTLES - To sanitize: Clean and rinse, then immerse for 5 minutes in solution prepared by thoroughly mixing 2 oz of this product with each 4 gallons cold or lukewarm water; drain; fill. If bottles are not filled promptly, rinse again with same strength bleach solution immediately before filling; drain thoroughly. Air dry. Ordinarily, 12 gallons of this strength solution will sanitize 5000 clean quart bottles. Keep this bleach solution clean and free from milk particles.

ICE CREAM FREEZERS - to clean and sanitize: After using, flush with warm water until water runs clear. Scrub or pressure-spray with solution prepared by thoroughly mixing 1 oz of this product with each 2 gallons of [regular] [powdered] detergent solution. Let stand 2 minutes. Rinse thoroughly with clean, clear water; drain. Air dry. Immediately before use, sanitize according to directions shown below.*

*BEFORE USE -rinse with sanitizing solution prepared by thoroughly mixing 2 oz of this product with each 4 gallons of water; drain thoroughly and air dry before use.

MEASUREMENT TABLE: PPM (Parts Per Million Available Chlorine)

1 oz of this product	- to One Gallon Water	= 400 ppm
1/2 oz (1 Tablespoon) of this product	- to One Gallon Water	= 200 ppm
1 drop of this product	- to One Gallon Water	= 2/3 ppm

CPPC BLEACH DISINFECTING GUIDE

This product is a 5.25% sodium hypochlorite solution containing approximately 5% available chlorine by weight. In addition to being a highly effective liquid chlorine bleach for laundering and household disinfecting, this product is widely used in sanitation of poultry and livestock houses and equipment, dairies, creameries, restaurants and taverns, as well as for purification of drinking water and disinfection of water for swimming and wading pools.

IMPORTANT: Always thoroughly mix with water as directed before using.

Do not allow undiluted bleach to come in contact with any fabric. (If it does, rinse out immediately with clear, cold water.)

Do not apply this product with natural sponge.

Do not use this product on steel, aluminum, silver, or chipped enamel.


If used on metal, bleach solution should be allowed to stand for no more than 5 minutes, and then rinsed off thoroughly with clear water; otherwise, it may slightly discolor and eventually corrode the metal.

If a metal sprayer is used to apply the bleach solution, rinse sprayer thoroughly after use with clear water, and then oil the plunger.

SEPTIC TANK OPERATION is not affected by regular home and farm use of of this product.

TABLE OF LIQUID MEASURES

3 teaspoons	=	1 Tablespoon	=	1/2 ounce = 1/16 cup
1 cup	=	16 Tablespoons	=	8 ounces = 1/2 pint



Boyle

67875-1 CPCC BLEACH (11 of 24)

For directions on sanitizing and disinfecting specific surfaces, write:
THE CLOROX COMPANY
 Consumer Services Department
 Oakland, California 94612-1888

CPCC BLEACH AS A FUNGICIDE FOR SEED POTATOES

This product is fungicidal to the Verticillium wilt organism *V. albo-atrum* (microsclerotial type) on seed potatoes. A bleach solution is applied to whole seed and freshly cut seed potato pieces during the cutting operation for planting. (Research at the Washington State University Irrigated Agriculture Research and Extension Center* has shown that treatment with a sodium hypochlorite solution helps to prevent the spread of organisms to uninfected soil or fields via seed potato surfaces.)

Use Instructions

Thoroughly mix a solution for spraying, in the proportion of one part of this product (5.25% sodium hypochlorite) to 9 parts of water. Use this solution to spray freshly cut seed potato pieces from the top and bottom of the cutting chain or elevator with a series of non-mist nozzles at 3 to 5 psi. Thoroughly cover all cut and uncut surfaces with the solution. The treatment will be most effective on clean seed tubers, as the organic matter in soil will reduce the effectiveness of the sodium hypochlorite.

Plant within four hours of the cutting and bleach treatment operation. If planting should be delayed, store the treated seed in clean open, well-ventilated bins or truck beds. Storing cut, wet seed in large unventilated containers will contribute to secondary breakdown from soft rot organisms.

Safety Precautions

Do not mix full-strength bleach or treatment solution with any agricultural chemical, ammonia, or acid. Avoid prolonged contact of this product with skin. If full strength or diluted bleach is splashed in the eyes, flush with water.

Conduct the spraying operations either outside, in a well-ventilated building, or under a hooded exhaust system. Use non-misting nozzles to avoid breathing of mist. Wear a face mask and plastic or rubber gloves and clothing. Because sodium hypochlorite is corrosive to many metals, chains and other machine parts should be either plastic or plastic-coated and rinsed with clear water after use.

CAUTION: DO NOT USE THE TREATED SEED FOR FOOD OR FEED. Use the bleach treatment only on crops and for the purposes recommended. Apply only as specified above. Do not apply in a dipping operation or the bleach solution may become contaminated with soil and organic matter from the potato surfaces and lose its effectiveness.

*Easton, G.D., M.E. Nagle, and D.L. Bailey, 1972. "Verticillium albo-atrum Carried by Certified Seed Potatoes into Washington and Control by Chemicals", American Potato Journal 49: 397-402.

CPCC BLEACH FOR MEAT & POULTRY PLANT PROCESSING WATER
 (USDA* Designation G4)


This product may be used in processing water of meat and poultry plants at concentrations up to 5 ppm (parts per million) calculated as available chlorine. Chlorine may be present in poultry chiller water, in water for reprocessing poultry carcasses internally contaminated with feces, and in red meat carcass final wash water at concentrations between 25 and 50 ppm calculated as available chlorine. Chlorine must be dispensed at a constant and uniform level and the method or system must be such that a controlled rate is maintained.

Measurement Table

Amount of this product to make ___ ppm available chlorine

Water volume	5 ppm	25 ppm	50 ppm
1 gallon	8 drops	40 drops	1 tsp
10 gallons	1/2 tsp	1 oz	3 tsp
100 gallons	2 oz	6 1/2 oz	13 oz
1000 gallons	13 oz	2 qt	1 gal

*Authorized by USDA for use in federally inspected meat and poultry plants.



R-10233-1

CPPC BLEACH FOR POTABLE WATER TREATMENT (USDA* Designation G4)

This product may be used in processing water of meat and poultry plants at concentrations up to 5 ppm calculated as available chlorine. Chlorine may be present in poultry chiller water, in water for reprocessing poultry carcasses internally contaminated with feces, and in red meat carcass final wash water at concentrations between 25 and 50 ppm (parts per million) calculated as available chlorine. Chlorine must be dispensed at a constant and uniform level and the method or system must be such that a controlled rate is maintained.

Measurement Table

Amount of this product to make ___ ppm available chlorine

Water volume	5 ppm	25 ppm	50 ppm
1 gallon	8 drops	40 drops	1 tsp
10 gallons	1/2 tsp	1 oz	3 tsp
100 gallons	2 oz	6 1/2 oz	13 oz
1000 gallons	13 oz	2 qt	1 gal

Use chlorine test strips to adjust to desired available chlorine level

*Authorized by USDA for use in federally inspected meat and poultry plants.

CPPC BLEACH FOR POULTRY CARE

Keeping poultry healthy, productive and profitable is largely a problem of disease prevention. Remedial measures are much more difficult and often less successful than preventing the spread of disease before it infects the flock. Regular use of this product in the sanitation and disinfection of chicken houses, brooders, and other poultry equipment is an effective aid in preventing many diseases of bacterial and viral origin.

DRINKING WATER - To disinfect, thoroughly mix 8 drops of this product with each 1 gallon clean water for 5 ppm available chlorine. Use in glass, porcelain, stoneware or concrete containers. Let stand 1 minute. Clean containers daily; rinse.

For young chicks, a solution made by thoroughly mixing 2 drops of this product with one gallon of water is satisfactory since baby chicks do not soil the water as rapidly as grown chickens, and the solution retains its effectiveness longer.

When cleaning drinking water containers, etc., a solution of 1 oz of this product to each quart water is effective in removing the slime. Do not allow birds to drink this solution.


TO CLEAN AND DISINFECT POULTRY HOUSES, BROODERS, HATCHERIES: Poultry houses should be cleaned and disinfected between cycles; hatcheries should be cleaned weekly or as necessary to keep sanitary. Metal surfaces can be satisfactorily disinfected. [Wooden surfaces are difficult to sanitize by any method.]

- (1) Remove all litter, loose dirt and debris.
- (2) Thoroughly mix solution of 1 oz [powdered] detergent and 6 oz of this product with each gallon of hot* or cold water.
- (3) Using this solution, scrub or pressure-spray all exposed areas, including floor, walls, ceiling posts and support beams. Let stand for 2 minutes.
- (4) Rinse with clean, clear, cold water.
- (5) Let dry thoroughly before introducing poultry.

METAL INCUBATORS, FEEDERS, WATER CONTAINERS, OTHER POULTRY EQUIPMENT AND UTENSILS - To clean and disinfect, remove loose dirt and debris. Scrub or pressure-spray with solution of 1 oz [powdered] detergent and 3 oz of this product thoroughly mixed with each gallon of hot* or cold water. Let stand for 2 minutes. Rinse with clear, cold water. Let dry.

*Where bleach/detergent solution is recommended for sanitizing poultry houses and equipment, use hot water (140°F or above) if available.

EGG WASHING: Wash eggs promptly after gathering. Dissolve 1 oz of detergent in each 2 gallons of water, add 1 oz this product to each 2 gallons of this solution; mix well. Washing solution temperature should be at least 20°F higher than that of eggs, but not more than 130°F. Rinse with clean or treated water at temperature at least 20°F higher than that of eggs, but not exceeding 120°F.



14 of 2

(57619-00) CPPC BLEACH (13 of 24)

(Treated water can be prepared by mixing 2 1/2 oz of this product per 5 gallons of water.)

FOR CONTINUOUS WASHERS: Prepare washing solution as above. Add an additional 1/2 oz each of detergent and this product per 4 gallons every 30 minutes. Dump wash tank and recharge every 2 hours. For manual method, soak eggs for only 1 to 2 minutes. Agitate basket. Make sure eggs are completely covered.

Air-dry eggs as rapidly as possible. Store in cool (55°F) room. Maintain relative humidity of 60-80%.

NOTE: Keep egg-washing equipment sanitary. Frequent cleaning will aid in operation and produce more sanitary eggs. While equipment is idle, bacteria can multiply. This contamination can be reduced by thoroughly flushing all equipment immediately before use with a solution of 1 oz of this product mixed with 2 gallons of water.

MEASUREMENT TABLE: PPM (Parts Per Million) Available Chlorine

1 oz of this product	-to One Gallon Water	= 400 ppm
1/2 oz (1 Tbsp) of this product	-to One Gallon Water	= 200 ppm
1 drop of this product	-to One Gallon Water	= 2/3 ppm

CPPC BLEACH SANITATION IN CARE OF LIVESTOCK, HORSES, PETS

TO CLEAN AND DISINFECT BARNs, STABLES, HUTCHES, KENNELS: Remove all litter, loose dirt and debris. Mix 1 oz [powdered] detergent and 6 oz of this product with each gallon hot* or cold water until detergent is dissolved. Using the solution, thoroughly scrub or pressure-spray all exposed areas including floor, walls, ceiling posts and support beams. Let stand for [at least] 2 minutes. Rinse with clean, clear, cold water. Let area dry thoroughly before housing animals.

LOADING AND HAULING EQUIPMENT: Loading chutes, trucks, trailers and other equipment for transportation of animals should be cleaned and disinfected prior to use. Pressure-spray or scrub with solution prepared by thoroughly mixing 1 oz [powdered] detergent and 3 oz of this product to each gallon hot* or cold water. Let stand for [at least] 2 minutes. Rinse with clean, clear, cold water. Allow to dry before use.

FEEDERS AND DRINKING WATER CONTAINERS - to clean and disinfect: Thoroughly scrub or pressure-spray with solution of 1 oz [powdered] detergent and 6 oz of this product to each gallon of hot* or cold water. Let stand for [at least] 2 minutes. Rinse thoroughly with clear, cold water; allow to drain dry. (A solution of 1 oz of this product to each quart of water is effective in removing slime which sometimes forms on drinking water containers.)

TO DISINFECT ANIMALS' DRINKING WATER: Thoroughly mix 8 drops of this product with 1 gallon clean water for 5 ppm available chlorine. Use in glass, plastic, porcelain or concrete containers daily. Let stand for [at least] 1 minute. Rinse with potable water. Air dry. (See directions above.)

*For bleach/detergent solution, use hot water if available.


MEASUREMENT TABLE: ppm (parts per million) Available Chlorine

6 oz of this product	-to One Gallon Water	= 2400 ppm
1/2 oz (1 Tablespoon) of this product	-to One Gallon Water	= 200 ppm
1 drop of this product	-to One Gallon Water	= 2/3 ppm

CPPC BLEACH SANITATION IN CARE OF SWINE

HOG HOUSES AND FARROWING HOUSES - to clean and sanitize:

- (1) Remove loose dirt, litter and debris. Dirty or coated surfaces cannot be sanitized.
- (2) Mix 1 oz [powdered] detergent and 6 oz of this product with each gallon of hot* or cold water until detergent is dissolved. Let stand for [at least] 2 minutes.
- (3) Scrub or pressure-spray all surfaces with this solution. Rinse with clear, cold water.
- (4) Allow to dry before housing pigs.



R-10233-1

15 of 26

(57619-00) CPPC BLEACH (14 of 24)

CLEAN AND DISINFECT METAL WATERING TROUGHS AND FEEDERS by pressure-spraying or scrubbing with solution prepared by thoroughly mixing 1 oz [powdered] detergent and 6 oz of this product with each gallon of hot* or cold water. Let stand for [at least] 2 minutes. Rinse thoroughly with clear, cold water; drain dry. (Drinking troughs and feeders should be cleaned and disinfected before housing pigs, and as often as necessary to keep sanitary.)

TO DISINFECT DRINKING WATER: Clean, then rinse. Thoroughly mix 8 drops of this product with each 1 gallon of clean water for 5 ppm available chlorine. (Water containing suspended material is difficult to sanitize.) Let stand for 1 minute.

NOTE: Clean metal surfaces can be sanitized using the above method. Wooden surfaces are difficult to sanitize by any method.

*For bleach/detergent solution, use hot water if available.

MEASUREMENT TABLE: PPM (Parts Per Million Available Chlorine)

6 oz of this product	-to One Gallon Water	= 2400 ppm
1/2 oz (1 Tablespoon) of this product	-to One Gallon Water	= 200 ppm
1 drop of this product	-to One Gallon Water	= 2/3 ppm

CPPC BLEACH IN SANITATION OF RESTAURANTS AND TAVERNS

An unclean kitchen and contaminated food can result in the spread of infectious disease. To help avoid this, it is important to keep all work surfaces, equipment and utensils hygienically clean. This product is a highly effective, economical and convenient germicide for this use in restaurants and taverns, as well as in the home.

TO SANITIZE WORK SURFACES (not utensils): After each use, scrub thoroughly with hot suds; rinse with clear, cold water. Then prepare a sanitizing solution of 1 Tablespoon of this product to each gallon of water. Apply this solution for 1 minute. Air dry.

TO DISINFECT WORK SURFACES (not utensils): After each use, scrub thoroughly with hot suds; rinse with clear, cold water. Then prepare a disinfecting solution of 3/4 cup of this product to each gallon of water. Apply this solution for 2 minutes. Rinse with potable water. Air dry.

TO SANITIZE DISHES, GLASSWARE, UTENSILS: Wash thoroughly; then soak 1 minute in solution of 1 Tablespoon of this product to each gallon of hot water. Use chlorine test strips to adjust to 200 ppm available chlorine. Drain dry. (Do not use on steel, aluminum, silver, or chipped enamel.)

DISINFECTING SINK AND SANITIZING DISHCLOTH[S]: should be a routine follow-up to dishwashing. First wash sink and rinse dishcloth[s] in hot suds. Drain out sudsy water. Then run a gallon of warm water into sink and mix in 3/4 cup of this product. Let stand 2 minutes. Swish dishcloth[s] in this solution; then use it to wipe sides of sink. Soak dishcloth[s] for 1 minute in this solution. Then rinse sink and dishcloth[s] with clear water.

TO DEODORIZE DRAIN PIPES: Flush with very hot water followed by 1 cup of this product. Wait 5 minutes; flush out with clear water.

TO SANITIZE REFRIGERATORS: First wash inside surfaces. Then wipe with solution of 1 Tablespoon of this product to each gallon of warm water. Let stand for [at least] 1 minute. Air dry. (Do not use on steel, aluminum, silver, or chipped enamel.)

ICE CREAM FREEZERS - to clean and sanitize: After using, flush with warm water until water runs clear. Scrub or pressure-spray with solution prepared by thoroughly mixing 2 oz of this product with each 2 gallons of [regular] [powdered] detergent solution. Rinse thoroughly with clean, clear water; drain. Immediately before use, sanitize for 2 minutes with solution prepared by mixing 2 oz of this product with each 4 gallons of water; drain thoroughly.

TO DISINFECT HARD NON-POROUS FLOORS: (plastic or ceramic tile): Mix 3/4 cup of this product per gallon of water. Mop or scrub. (Do not use on cork or linoleum.) Let stand 2 minutes. Rinse.

TO SANITIZE BRUSHES, MOPS & BROOMS: After using brushes, mops and brooms, wash thoroughly; then soak for 2 minutes in solution of 3/4 cup of this product to 1 gallon of warm water. Rinse with clear water; dry. (Not recommended for cellulose sponge mops.)

TO SANITIZE PAILS AND DUSTPANS: Remove heavy dirt prior to cleaning. Wash with solution prepared by thoroughly mixing 3/4 cup of this product with each gallon water. Let stand 2 minutes. Rinse with clear, cold water. Air dry.

TO DEODORIZE AND SANITIZE GARBAGE CANS: Remove heavy dirt with a cleaner. Rinse. Pour in 3/4 cup of this product to each gallon of water; mix. Swab inside surfaces with this solution. Let stand 2 minutes. Rinse with clear water; dry.

ACCEPTED
 APR 28 1998
 Under the Federal Insecticide, Fungicide, and
 Rodenticide Act, as amended, for the
 pesticide registered under
 EPA Reg. No. 67619-1

CPPC BLEACH FOR SANITIZING SOLUTIONS FOR EQUIPMENT AND UTENSILS

This product is authorized by USDA for use in federally inspected meat and poultry plants as a sanitizing solution.

Before using this product, food products and packaging materials must be removed from the room or kept protected.

Before they are treated with a bleach solution, the food processing equipment and utensils must be thoroughly washed and then rinsed with clear, cold water.

The bleach solution used for sanitizing should not exceed 200 ppm (parts per million) available chlorine (one part of this product to 250 parts water, or 1 Tablespoon per gallon). The bleach solution should be applied by spraying, soaking or scrubbing. Treated surfaces should remain wet for at least one minute.

A potable water rinse is not required, provided the equipment and utensils are adequately drained before they come into contact with food. Little or no residue should remain to adulterate or otherwise affect edible products.

CPPC BLEACH FOR SWIMMING POOL DISINFECTION

This product is a 5.25% sodium hypochlorite solution, containing approximately 5% available chlorine by weight. The purity of its ingredients and the carefully supervised process of its manufacture make this product a very stable, uniform source of chlorine for water treatment in swimming and wading pools. This product is especially suitable for use in chlorinators as it is a liquid and has no insoluble particles. This product is widely used as a source of chlorine for swimming pool sanitation and does not have any adverse effects on materials used in pool construction including swimming pool liners.

For each new filling of your pool, use following initial dosages of this product.

Swimming Pool Size in Gallons	Initial Dosage of this product	Swimming Pool Size in Gallons	Initial Dosage of this product
5,000	3 cups	20,000	13 cups
6,000	4 cups	25,000	17 cups
8,000	5 cups	30,000	20 cups
10,000	6 cups	35,000	23 cups
15,000	10 cups		

NOTE: 2 cups = 1 pint; 4 cups = 1 quart; 16 cups = 1 gallon


To determine the volume of water in the pool when filled, figure 7½ gallons of water for each cubic foot of pool capacity. One quart of this product per 6,000 gallons of water will supply approximately 2 ppm (parts per million) available chlorine, but this may dissipate rather rapidly in new water depending on the general sanitation conditions of the pool. Repeat dosage as needed to obtain 0.6 to 1.0 ppm available chlorine.

In chlorinating a swimming pool, mix the required amount of this product with 10 parts water and feed this solution through a chlorinator into the main water supply line to the pool. The feeding rate should be adjusted so the required quantity of this product will be added uniformly throughout the filling of the pool; or, if the water is circulated through a filter, the bleach should be added throughout one complete circulation. If this product cannot be fed into the main water supply line, mix ½ pint of this product with 5 gallons of water and scatter over a portion of the pool surface; repeat until the required amount of this product has been scattered over entire surface of the pool.

Check chlorine level in pool water at least daily with a pool testing set and add this product as needed to maintain 0.6 to 1.0 ppm available chlorine. One pint of this product per 6,000 gallons of water will supply approximately 1.0 ppm available chlorine. Frequency of application of this dosage will vary depending on number of people using the pool, weather conditions (sunlight exposure), and general cleanliness of the pool area. Chlorine level for acid-stabilized pools should be maintained at 1.0 - 1.5 ppm available chlorine.

The effectiveness of the chlorine is best when the pool water has a pH range of 7.2 to 7.6. The pH of the pool water should be checked daily using a pool pH testing set and adjusted as necessary.

The regular use of this product, in the above proportions, in the swimming pool usually prevents the growth of algae in the water; however, if algae growth is causing the pool water to look cloudy and uninviting, it may be corrected by doubling the initial dosage of this product for a few treatments (2 quarts instead of 1 quart per 6,000 gallons of new water). This additional product should be added to the pool in the evening after the pool is out of use so the excess chlorine will be dissipated before the pool is used again.



17 of 26

(57619-03) CPPC BLEACH (16 of 24)

If algae are growing on the bottom or walls of the pool, scrub pool with a solution of 1/2 gallon of this product to 5 gallons of water applying solution with a fiber brush. The pool should be scrubbed while full and rinsed off when algae growth has been removed; all of the growth and dirty solution should be flushed from the pool with clear water before the pool is refilled. Avoid skin contact with undiluted product; if such contact occurs, rinse immediately with water. When added as recommended, this product has no deleterious effect on the eyes, nasal passages, or skin of people using the pool and will have no effect on swimming apparel.

CPPC BLEACH FOR WADING POOL DISINFECTION

This product - a 5.25% sodium hypochlorite solution containing approximately 5% available chlorine by weight - is a very stable, uniform and convenient source of chlorine for water treatment in swimming and wading pools. Also, because this product is a liquid with no soluble particles, it is especially suitable for this use.

In chlorinating wading pools, use 1/8 cup per 100 gallons of new water. Mix required amount of this product with 2 gallons of water and scatter over surface of pool. Mix uniformly with pool water.

Between fillings of pool, add 1 Tablespoon of this product per 100 gallons of water each day. Empty small pools daily. (This product will not harm plastic pools.)

Stabilized pools should maintain a residual of 1.0 to 1.5 ppm available chlorine. Test the pH, available chlorine residual and alkalinity of the water frequently with appropriate test kits. Frequency of water treatment will depend upon temperature and number of swimmers.

The chart below is a guide to the amount of this product required to provide approximately 1/8 cup per 100 gallons of water figured on basis of water depths in pools of various diameters.

Depth of Water	Pool Diameter				
	4 Ft	6 Ft	8 Ft	10 Ft	15 Ft
6 inches	1/16 cup	1/8 cup	1/4 cup	3/8 cup	3/4 cup
1 foot	1/8 cup	1/4 cup	1/2 cup	3/4 cup	1 5/8 cups
2 feet	1/4 cup	1/2 cup	1 cup	1 1/2 cups	3 1/4 cups
3 feet	3/8 cup	3/4 cup	1 1/2 cups	2 1/4 cups	5 cups

TABLE OF LIQUID MEASURES:

3 teaspoons = 1 Tablespoon = 1/2 ounce = 1/16 cup
 1 cup = 16 Tablespoons = 8 ounces = 1/2 pint

CPPC BLEACH BACTERICIDAL EFFICACY

This product, when used as directed below, is effective against the following bacteria:

Gram positive bacteria including:
Staphylococcus aureus
Streptococcus pyogenes

Gram negative bacteria including:
Salmonella choleraesuis
Pseudomonas aeruginosa
Escherichia coli serotype O157:H7
Legionella pneumophila
Shigella dysenteriae

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Prepare a solution of 3/4 cup (6 oz) of this product per gallon of water (2400 ppm available chlorine). -or- [Prepare a dilution of 1 part of this product to 9 parts water (13 oz/gallon) (~5000 ppm available chlorine).] Thoroughly wet surface with the solution and allow it to remain on the surface for 2 minutes. Rinse with clean water and dry.

TO SANITIZE GARBAGE CANS/DIAPER PAILS: Pre-clean garbage can/diaper pail with a cleaning product prior to sanitization. Rinse with water and

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 57619-03

drain. Pour in $\frac{3}{4}$ cup of this product to each gallon of water. Mix. Let stand [at least] 2 minutes. Rinse and air dry.

Toilet Bowls: Flush toilet to remove gross filth. Add $\frac{3}{4}$ -or- 1 cup of this product to the bowl and brush surfaces thoroughly, making sure to get under the rim. Let stand 2 minutes before flushing again.

Use Sites

This product can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms), eating establishments, pet kennels and veterinary premises.

CPPC BLEACH (5.25% Sodium hypochlorite, EPA Reg. No. 67619-xx) TUBERCULOCIDAL EFFICACY

This product, when used as directed below, is effective against *Mycobacterium bovis*.

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Prepare a dilution of 1 part of this product to 9 parts water (13 oz/gallon) (~5000 ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 5 minutes. Rinse with clean water and dry.

Use Sites

This product can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms), eating establishments, pet kennels and veterinary premises.

CPPC BLEACH (5.25% Sodium hypochlorite, EPA Reg. No. 67619-xx) VIRUCIDAL[®] EFFICACY

*This product, when used as directed below, is effective against the following viruses on hard, nonporous, inanimate surfaces:

- | | |
|--|--------------------------|
| ■ Adenovirus Type 2 | ■ Rotavirus |
| ■ Hepatitis A | ■ Cytomegalovirus |
| ■ Human Immunodeficiency Virus Type 1 (HIV-1)* | ■ Influenza A2 |
| ■ Respiratory syncytial virus | ■ Varicella zoster virus |
| ■ Herpes simplex virus 2 | ■ Rhinovirus Type 17 |
| ■ Rubella virus | ■ Canine parvovirus** |
| ■ Feline parvovirus** | |

Directions for use:

Hard Non-porous Surfaces:


To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.) Prepare a solution of $\frac{3}{4}$ cup (6 oz) of this product per gallon water (~2400 ppm available chlorine). -or- Prepare a dilution of 1 part of this product to 9 parts water (13 oz/gallon) (~5000 ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 5 minutes. Rinse with clean water and dry.

*see directions in the Clorox Service Bulletin entitled "Special Instructions for Using CPPC BLEACH to Clean and Decontaminate Against HIV on Surfaces/Objects Soiled with Blood/Body Fluids"

**For Canine and Feline parvovirus use the same instructions as above but keep the solution in contact with the surface for 10 minutes.

TO SANITIZE GARBAGE CANS/DIAPER PAILS: Pre-clean garbage can/diaper pail with a cleaning product prior to sanitization. Rinse with water and drain. Pour in $\frac{3}{4}$ cup of this product to each gallon of water. Mix. Let stand [at least] 1 minute. Rinse and air dry.

Toilet Bowls: Flush toilet to remove gross filth. Add $\frac{3}{4}$ -or- 1 cup of this product to the bowl and brush surfaces thoroughly, making sure to get under the rim. Let stand 2 minutes before flushing again.



19 of 26

(EPA) CPPC BLEACH (15 of 24)

Use Sites

This product can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms), eating establishments, pet kennels and veterinary premises.

CPPC BLEACH FUNGICIDAL ** EFFICACY

This product, when used as directed below, is effective against molds, Athlete's foot fungus** (Trichophyton mentagrophytes), and Mildew** (Aspergillus niger).

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Prepare a solution of 3/4 cup (6 oz) of this product per gallon of water (2400 ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain on the surface for 2 minutes. Rinse with clean water and dry.

TO SANITIZE GARBAGE CANS/DIAPER PAILS: Pre-clean garbage can/diaper pail with a cleaning product prior to sanitization. Rinse with water and drain. Pour in 3/4 cup of this product to each gallon of water. Mix. Let stand [at least] 2 minutes. Rinse and air dry.

Use Sites

This product can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, shower stalls, bathrooms, nurseries, sick rooms, laundry rooms), eating establishments, pet kennels and veterinary premises.

Special Instructions for Using CPPC BLEACH to Clean and Decontaminate Against HIV on Surfaces/Objects Soiled with Blood/Body Fluids

This product kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings (e.g. hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

Personal Protection: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings.



Cleaning procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and other objects before applying this product.

Contact time: Dilute the product to 1:20 (3/4 cup to a gallon) and spray or flood surface; let stand 2 minutes.

Disposal of infectious materials: Use disposable latex gloves, gowns, masks, and eye coverings. Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

FRUIT & VEGETABLE WASHING

Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 1 Tbsp. of this product in 7 1/2 gallons -or- 1 oz of this product in 15 gallons of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.



200926

(67619-xx) CPPC BLEACH (19 of 24)

CPPC BLEACH (5.25% Sodium hypochlorite, EPA Reg. No. 67619-xx) CANDIDA ALBICANS EFFICACY

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.) Prepare a solution of 3/4 cup (6 oz) of this product per gallon water (~2400 ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 2 minutes. Rinse with clean water and dry.

TO SANITIZE DIAPER PAILS: Pre-clean diaper pails with a cleaning product prior to sanitization. Rinse with water and drain. Pour in 3/4 cup to each gallon of water. Mix. Let stand [at least] 2 minutes. Rinse and air dry.

Use Sites

This product can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms), eating establishments, pet kennels and veterinary premises.

CPPC BLEACH FOR DISINFECTION USES (2400 PPM FOR 1 MINUTE) [(SPRAY APPLICATIONS)]

This product can be diluted and spray applied for convenient broad spectrum disinfection of hard, non-porous surfaces in homes, hospitals, commercial, institutional, and eating establishments.

To ensure [sodium] hypochlorite [bleach] stability, prepare solutions daily.

To prepare a 2400 ppm solution, mix:

5 teaspoons of this product	-in-	1 pint water
3 Tablespoons of this product	-in-	1 quart water
3/4 cup of this product	-in-	1 gallon water

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Spray surface using a coarse spray with the [diluted] bleach solution until thoroughly wet. Allow it to remain on the surface for 1 minute. Rinse and dry.

CPPC BLEACH FOR DISINFECTION USES (800 PPM FOR 2 MINUTES) [(SPRAY APPLICATIONS)]

This product can be diluted and spray applied for convenient broad spectrum disinfection of hard, non-porous surfaces in homes, hospitals, commercial, institutional, and eating establishments.

To ensure [sodium] hypochlorite [bleach] stability, prepare solutions daily.

To prepare a 800 ppm solution, mix:

1 1/2 teaspoons of this product	-in-	1 pint water
1 Tablespoon of this product	-in-	1 quart water
1/4 cup of this product	-in-	1 gallon water

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, [first] clean surface by removing gross filth (loose dirt, debris, food materials, etc.) Spray surface using a coarse spray with the [diluted] bleach solution until thoroughly wet. Allow it to remain on the surface for 2 minutes. Rinse and dry.

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

2/pt 26

(57619-xx) CPPC BLEACH (20 of 24)

USE CPPC BLEACH IN SANITIZING CYCLE OF CHEMICAL SANITIZING DISHWASHING MACHINES

This product is an approved sanitizing agent for use in the sanitizing cycle of chemical sanitizing dishwashing machines.
Usage Directions

1. Hook up a bleach bottle to the automatic bleach dispensing system of the chemical sanitizing dishwashing machine. If the bottle is already in place, make sure that sufficient product remains in the bottle to complete the dishwashing job.
2. Wash tableware in the machine following the manufacturer's operating instructions.
3. After the washing/rinsing/sanitizing cycles are completed, remove the dishwashing rack. Let stand 2 minutes. Allow the tableware to air dry.

Caution: Do not sanitize silverware or pewter with this product as these metals may darken.

Bleach Dispensing System Adjustments

The following steps must be followed before using the new chemical sanitizing dishwashing machine, and on a regular basis thereafter:

- a. Start machine and let run until the machine has begun the final rinse cycle.
- b. Take a sample of the rinse water.
- c. Using a special test kit, determine the part per million (ppm) of available chlorine in the sample.
- d. If the ppm of available chlorine is lower than the minimum or higher than the maximum level of available chlorine permitted by local public health authorities, adjust the bleach dispensing system.
- e. Repeat steps "a" through "c" until a correct ppm of available chlorine is achieved.

Your equipment service representative or dishwashing detergent supplier will often make these adjustments for you.

Correct Chlorine Concentration

Local public health codes vary with regard to the parts per million of available chlorine permitted in the final rinse water of chemical sanitizing dishwashing machines. The minimum level is 50 ppm of available chlorine with a maximum level of 200 ppm, although some states require 100 ppm minimum level. Check with your local public health department on the applicable regulations for your area.

PLANT PARASITIC NEMATODES AND PLANT DISEASE-CAUSING FUNGI QUARANTINE USE DIRECTIONS

- Crop/Site/Commodity:**
1. Walks, benches, tools, plant containers in nurseries and other quarantine areas.
 2. Farm equipment and machinery.
 3. Laboratory work areas, equipment and specimens.
 4. Deciduous fruit tree nursery stock (dormant)
- Target Pest/Problem:** Plant parasitic nematodes, plant disease-causing fungi and for general surface disinfection.
- Dosage:** See dilution rate.
- Dilution Rate:** Five parts water with one part this product (equals 0.875% active ingredient).

Laboratory work areas, equipment and specimens: Prepare a solution of four or five parts water to one part product. Scrub areas and implements thoroughly then wipe or allow to dry naturally. It is advisable that workers doing the treatment wear waterproof gloves. Small tools or implements, and other items covered above may be immersed for five to ten minutes in the solution instead of scrubbing manually. Wipe off plant tissue or soak tissue in the solution.

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

R-10233-1

22 of 26

(57619-xx) CPPC BLEACH (21 of 24)

Deciduous Fruit Tree Nursery Stock: Four or five parts water with one part product (equals 0.85% to 1.0% active ingredient)

Method of Application: Drench and dip method.

Deciduous Fruit Tree Nursery Stock:

1. Thoroughly clean all soil from roots.
2. Dip entire tree root system in solution for 30 to 45 seconds.
3. Immediately rinse tree root system with clean water upon removal from dip solution.

Frequency/Timing of Application: As needed.

Deciduous Fruit Tree Nursery Stock:
One application at harvest (tree-digging period)

Field Reentry After Application: Not applicable.

Preharvest Interval: Not applicable.

Other Requirements: Do not apply through any type of irrigation system.

Deciduous Fruit Tree Nursery Stock:
Workers required to wear eye protection and waterproof gloves.

KARNAL BUNT QUARANTINE TREATMENT USE DIRECTIONS

Crop/Site/Commodity: Tools, conveyances, mechanized farm equipment, seed conditioning or milling equipment, soil moving equipment, or grain elevators and structures used for storing and handling grain.

Target Pest/Problem: Karnal bunt (*Tilletia indica*)

Dosage: See dilution rate.

Dilution Rate: Mix 1 part this product to 2.5 parts water.

Method of Application: Before treating remove all soil and plant debris. The dilute solution of sodium hypochlorite will be used to wet the point of runoff surfaces potentially exposed to the pathogen. Saturate any soil removed by the treatment with the solution. It is recommended that the equipment or site be thoroughly washed down with clean water after 15 minutes to minimize corrosion.

Additional Restrictions, User Precautions and Requirements:

Be sure treated surfaces are dry before handling. Protective impervious gloves and safety goggles should be worn when mixing this disinfectant. Protective impervious gloves should be worn while handling or using the dilute mixture. Do not ingest product. Do not breathe vapor or mist. Do not get in eyes, on skin, or on clothing. Store away from flammable materials. Keep containers tightly closed. Wash thoroughly after handling. Use only with adequate ventilation.

EMERGENCY DISINFECTION AFTER MAIN BREAKS

MAINS

Before assembly of the repaired section, flush out mud and soil. Permit water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

R-10233-1

23 of 26

**DISINFECTION OF DRINKING WATER (POTABLE)
(EMERGENCY/PUBLIC/INDIVIDUAL SYSTEMS)**

PUBLIC SYSTEMS

Mix a ratio of this product to water to produce a 10 ppm available chlorine by weight. Begin feeding this solution with a hypochlorinator until a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm is attained throughout the distribution system. Check water frequently with a chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Contact your local Health Department for further details.

EMERGENCY DISINFECTION AFTER DROUGHTS

A. SUPPLEMENTARY WATER SUPPLIES

Gravity or mechanical hypochlorite feeders should be set up on a supplementary line to dose the water to a minimum chlorine residual of 0.2 ppm after a 20 minute contact time. Use a chlorine test kit.

B. WATER SHIPPED IN BY TANKS, TANK CARS, TRUCKS, ETC.

Thoroughly clean all containers and equipment. Spray a 50 ppm available chlorine solution and rinse with potable water after 5 minutes. During the filling of the containers, dose with sufficient amounts of this product to provide at least a 0.22 ppm chlorine residual. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER FIRES

CROSS CONNECTIONS OR EMERGENCY CONNECTIONS

Hypochlorination or gravity feed equipment should be set up near the intake of the untreated water supply. Apply sufficient product to give a chlorine residual of a least 0.1 to 0.2 ppm at the point where the untreated supply enters the regular distribution system. Use a chlorine test kit

EMERGENCY DISINFECTION AFTER FLOODS

WELLS

Thoroughly flush contaminated casing with a 500 ppm available chlorine solution. Backwash the well to increase yield and reduce turbidity, adding sufficient chlorinating solution to the backwash to produce a 10 ppm available chlorine residual, as determined by a chlorine test kit. After the turbidity has been reduced and the casing has been treated, add sufficient chlorinating solution to produce a 50 ppm available chlorine residual. Agitate the well water for several hours and take a representative water sample. Retreat well if water samples are biologically unacceptable.

ASPHALT OR WOOD ROOFS AND SIDINGS

To control fungus and mildew, first remove all physical soil by brushing and hosing with clean water. Apply a 5000 ppm available chlorine solution by brushing or spraying roof or siding. After 30 minutes, rinse by hosing with clean water.

ACCEPTED
APR 28 1998
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 62619-1

24 of 26

SPAS, HOT-TUBS, IMMERSION TANKS, ETC.

A. SPAS/HOT-TUBS

Using a dilution chart or formula, calculate and approximate an amount of product per 1000 gallons of water to obtain a free available chlorine concentration of 5 ppm, as determined by a suitable chlorine test kit. Adjust and maintain pool water pH to between 7.2 and 7.8. Some oils, lotions, fragrances, cleansers, etc. may cause foaming or cloudy water as well as reduce the efficiency of the product.

1. *Maintaining the Water:* To maintain the water, apply the product solution over the surface to maintain a chlorine concentration of 5 ppm.
2. *After Each Use:* Shock treat to control odor and algae, using the product at a rate of 16 ounces to 500 gallons of water.
3. *Periods of Disuse:* During periods of disuse, add product daily to maintain a 3 ppm chlorine concentration.

B. HUBBARD AND IMMERSIONS TANKS

Before patient use, add product to obtain a chlorine residual of 25 ppm, as determined by a suitable test kit. Adjust and maintain the water pH to between 7.2 and 7.6. After each use, drain the tank. Add 10 ounces of product to a bucket of water and circulate this solution through the agitator of the tank for 15 minutes and then rinse out the solution. Clean tank thoroughly and dry with clean cloths.

C. **HYDROTHERAPY TANKS.** Add product to the water to obtain a chlorine residual of 1 ppm as determined by a suitable chlorine test kit. Pool should not be entered until the chlorine residual is below 3 ppm. Adjust and maintain the water pH to between 7.2 and 7.6. Operate pool filter continuously. Drain pool weekly, and clean before refilling.

WATER DISINFECTION

INDIVIDUAL SYSTEMS

1. **DUG WELLS:** Upon completion of the casing (fining), wash the interior of the casing (lining) with a 100 ppm available chlorine solution using a stiff brush. After covering the well, pour the sanitizing solution into the well through both the pipesleeve opening and the pipeline. Wash the exterior of the pump cylinder also with the sanitizing solution. Start pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours flush well until all traces of chlorine have been removed from the water. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS

1. **DRILLED, DRIVEN AND BORED WELLS:** Run pump until water is as free from turbidity as possible. Pour a 100 ppm available chlorine sanitizing solution into the well. Add 5 to 10 gallons of clean, chlorinated water to the well in order to force the sanitizer into the rock formation. Wash the exterior of pump cylinder with the sanitizer. Drop pipeline into well, start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours flush well until all traces of chlorine have been removed from the water. Deep wells with high water levels may necessitate the use of special methods for introduction of the sanitizer into the well. Mix well [(2 drops to 1 quart)] and consult your local Health Department for further details.

2. **FLOWING ARTESIAN WELLS:** Artesian wells generally do not require disinfection. If analysis indicate persistent contamination, the well should be disinfected. Consult your local Health Department for further details.

EMERGENCY DISINFECTION

When boiling water for 1 minute is not practical, water can be made potable by using this product. Prior to addition of the sanitizer, remove all suspended material by filtration or by allowing it to settle to the bottom. Decant the clear or clarified, contaminated water to a clean container and add 8 drops of this product (dependent on product strength) to 1 gallon of water [(2 drops to 1 quart)]. Allow the treated water to stand for 30 minutes. Properly treated water should have a slight chlorine odor. If not, repeat dosage and allow the water to stand an additional 15 minutes. The treated water can then be made palatable by pouring it between clean containers for several times.

For cloudy water, use 16 drops of this product per gallon of water [(4 drops to 1 quart)]. If no chlorine odor is apparent after 30 minutes, repeat dosage and wait an additional 15 minutes.

ACCEPTED

APR 28 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

25 of 24

Crop/Site: Asparagus Seed treatment

Target Pest/Problem: To aid in the prevention of asparagus root rot (*Fusarium oxysporium* and *F. asparagi*)

Dosage: 1 pint plus 8 pints of water.

Dilution or Application Rate: Use 1 gallon of solution per pound of seed.

Method of Application: Wash seed in solution for 40 minutes, providing continuous agitation. After washing seed, spread and air dry.

Frequency/Timing of Applications: 1 application.

Preharvest Interval: Preplant treatment.

Other Requirements: Do not use treated seeds for food or feed. Allow to dry before storing, planting, or treating with other chemicals. Prepare fresh solution for each batch of seed.

Crop/Site: Pepper Seed treatment.

Target Pest/Problem: To aid in the prevention of bacterial spot (*Xanthomonas vesicatoria*)

Dosage: 2 pints plus 8 pints of water.

Dilution or Application Rate: Use 1 gallon of solution per pound of seed.

Method of Application: Wash seed in solution for 40 minutes, providing continuous agitation. After washing seed, spread to air dry.

Frequency/Timing of Application: 1 application.

Preharvest Interval: Preplant treatment.

Other Requirements: Do not use treated seed for food or feed. Allow to dry before storing, planting, or treating with other chemicals. Prepare fresh solution for each batch of seed.

Crop/Site: Tomato Seed treatment.

Target Pest/Problem: To aid in the control of bacterial canker (*Corynebacterium michiganense*) and Tobacco mosaic virus (TMV).

Dosage: 2 pints plus 8 pints of water.

Dilution or Application Rate: Use 1 gallon solution per pound of seed.

Method of Application: Wash seed in solution for 40 minutes, providing continuous agitation. After washing seed, spread to air dry.

Frequency/Timing of Application: 1 application.

Preharvest Interval: Preplant treatment.

Other Requirements: Do not use treated seed for food or feed. Allow to dry before storing, planting, or treating with other chemicals. Prepare fresh solution for each batch of seed.

ACCEPTED
APR 28 1998
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 676197

26026

(67619-xx) CPPC BLEACH (25 of 24)

PRECAUTIONARY STATEMENTS

WARNING:

Hazards to humans and domestic animals:
Causes substantial but temporary eye injury. Do not get in eyes or on clothing.
Harmful if swallowed. May irritate skin. For prolonged use, wear gloves

Statement of Practical Treatment

If in eyes: remove contact lenses and rinse with plenty of water for at least 15 minutes. IF SWALLOWED, drink large amounts of water. DO NOT induce vomiting. [In either case,] call a physician or poison control center immediately.
If in contact with skin: immediately remove contaminated clothing and wash skin thoroughly with water.

Physical and chemical hazards:

This product contains a strong oxidizer. Flush drains before and after use. Do not use or mix with other household chemicals, such as toilet bowl cleaners, rust removers, acid or ammonia containing products. To do so will release hazardous gases. Prolonged contact with metal may cause pitting or discoloration.

Environmental hazards:

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge.

Storage and Disposal:

Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water. Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not reuse empty container but place in trash collection. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

INFORMACIÓN PREVENTIVA

ADVERTENCIA:

Dañino para las personas y los animales domésticos:
Causa lesiones serias pero temporales a los ojos. No permita que entre en los ojos o caiga en la ropa. Es dañino si se ingiere. Puede causar irritaciones a la piel. Para uso prolongado use guantes.

Declaración de Tratamiento Práctico

Si entra en los ojos: quítese los lentes de contacto, enjuáguese los ojos con agua en abundancia durante por lo menos 15 minutos. Si es ingerido, tome bastante cantidad de agua. No induzca vomito. [En cualquier caso,] llame a un medico o al centro de control de veneno. Si entra en contacto con la piel: inmediatamente quítese la ropa contaminada y lávese bien con agua la zona afectada.

Cuidados Químicos y Físicos:

Este producto contiene un fuerte oxidante. Deje correr agua abundantemente en la tubería antes y después del uso. No lo utilice ni mezcle con otros productos químicos del hogar tales como productos para limpiar el inodoro, removedores de óxido, o productos que contengan ácidos o amoníaco. El hacerlo causaría la emisión de gases dañinos. El contacto prolongado con metales puede causar corrosión o decoloración.

Peligros para el medio ambiente

No bote los restos que contienen residuos de este producto en lagos, lagunas, ríos, estuarios, océanos o en cualquier otro cuerpo de agua, a no ser que se haga de acuerdo y con un permiso siguiendo los requerimientos del Sistema Nacional de Eliminación de Descargos y Contaminación. Las autoridades pertinentes tienen que ser notificadas previamente por escrito.

Almacenaje y modo de desechar el envase:

Para que no se deteriore, almacene este producto en áreas secas y frescas manténgalo alejado del calor y del sol. En caso de derrame, lave el área afectada con bastante agua. Para desechar el producto, dilúyalo con agua en la botella antes de botarlo. No use el envase vacío, bótelo en la vase. No contamine el agua, los alimentos o la comida de animales al almacenar, desechar o limpiar equipos.

SATISFACTION GUARANTEED

Satisfaction Guaranteed! At Clorox, we have prided ourselves in making consistent quality bleach for over 80 years, and we are dedicated to ensuring your total satisfaction with our product. If you are not completely satisfied, please call the number below.

QUESTIONS or COMMENTS? -or- SUGGESTIONS?

Please call (800) _____

or write us at: Clorox Consumer Services, P.O. Box 24305, Oakland, CA 94623.

Clorox is a registered trademark of The Clorox Company.
Bottle shape is a registered trademark of The Clorox Company.

CLOROX CUSTOMER ASSISTANCE (800)

Laundry/Household Cleaning Questions? The experts at Clorox are standing by to answer any of your laundry or household cleaning questions.

ENVIRONMENTAL COMMITMENT



This bottle is coded for recyclers. Check to see if recycling facilities in your area accept colored HDPE bottles. Recycle where recycling facilities available.



This product rapidly breaks down almost entirely to salt [and] water.
This product is safe for septic and waste water treatment systems.

This product contains no phosphates -or- phosphorus.

EPA Est. No. 67619

- CA-2
- CA-3
- CA-4
- FL-1
- GA-1
- IL-1
- MO-2
- OH-1
- PR-1
- TX-1

Batch code identifies manufacturing location.


EPA Reg. No. 67619-xx-R
EPA Est. No. 2675-MN-1 -or- EPA Est. No. 5813-
Batch Code Identifies Manufacturing Location

ACCEPTED

APR 28 1998

Under 23 Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67619-1

Mfd. for & © The Clorox Professional Products Company,
Oakland, CA 94612
Made in U.S.A.
Printed in U.S.A.



R-10233-1