



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

April 3, 2020

Joshua Wood  
Regulatory Affairs Specialist  
SePRO Corporation  
11550 North Meridian Street Suite 600  
Carmel, IN 46032

Subject: Registration Review Label Mitigation for Flurprimidol  
Product Name: Cutless 50W Turf Growth Regulator  
EPA Registration Number: 67690-15  
Application Date: December 14, 2018  
Decision Number: 560955

Dear Mr. Wood:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the Flurprimidol Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2  
EPA Reg. No. 67690-15  
Decision No. 560955

If you have any questions about this letter, please contact Srijana Shrestha by phone at 703-305-6471, or via email at [shrestha.srijana@epa.gov](mailto:shrestha.srijana@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read 'Linda Arrington', with a stylized flourish at the end.

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

Enclosure

[Front label for Non-refillable, Non-rigid containers, any size]



# Cutless® 50W

Turf Growth Regulator

For growth management and quality improvement of turfgrasses.

### Active Ingredient

Flurprimidol:  $\alpha$ -(1-methylethyl)- $\alpha$ -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol .....50.0%

Other Ingredients .....50.0%

TOTAL .....100.0%

Contains 0.50 pound active ingredient per one (1) U.S. pound of product.

## Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

Refer to label booklet for additional Precautionary Information and Directions for Use, including First Aid and Storage and Disposal.

**Notice:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet. If terms are unacceptable, return at once unopened.**

EPA Reg. No. 67690-15  
FPL20190724


EPA Est. No. \_\_\_\_\_  
[P/N] \_\_\_\_\_

Cutless is a registered trademark of SePRO Corporation.

SePRO Corporation 11550 N. Meridian Street, Suite 600 Carmel, IN 46032, U.S.A.

[Turf Growth Regulator]

Net Contents \_\_\_\_\_ (Non-refillable)



[Label booklet text, ALL containers]

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b>.</p>	

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Caution. Harmful If Swallowed or Inhaled. Causes moderate eye irritation. Avoid breathing dust or spray mist and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.**

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks; and
- Chemical-resistant gloves (such as natural rubber or barrier laminate).

#### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

### Ground Water Advisory

This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory

This product is classified as having potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs, will reduce the potential loading of this product from runoff and sediment.

---

## DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying this product. Use only according to label directions.

**Note to User: Product packets are water-soluble. Do not allow packet to become wet prior to use. Do not handle with wet hands. Reseal container to protect remaining packets.**

**Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

## POLLINATOR ADVISORY STATEMENT

Protect forage and habitat of pollinators including the monarch butterfly (and its larvae), birds, and bats by following label directions, and making only directed applications.

## INFORMATION FOR GROWTH REGULATION OF PERENNIAL TURFGRASSES

Cutless 50W Turf Growth Regulator is a Class B plant growth regulator (PGR) which reduces leaf blade length and stem internode elongation in turfgrasses resulting in a more compact growth form. Growth regulation results from suppression of gibberellic acid biosynthesis. Under normal growing conditions root growth and lateral expansion of turf are not affected. Follow an appropriate fertility program for the desired turf species in conjunction with applications of this product to provide best turfgrass enhancement and reduce potential for discoloration. Make broadcast treatments on medium to high quality turfgrass areas of uniform species composition. Turf containing significant amounts of coarse textured species such as tall fescue, orchardgrass, timothy, dallisgrass, etc., may respond unevenly to Cutless 50W treatment.

## Benefits of Cutless 50W Applications to Turfgrass

- Shoot growth suppression of warm- and cool-season turfgrasses resulting in decreased mowing frequency and turfgrass clippings.
- Increased turfgrass density, wear resistance, and improved color on warm- and cool-season turfgrass species resulting in improved turf quality.
- Suppressed growth of *Poa annua*, reducing populations and shifting competitive growth advantage towards perennial turfgrasses.
- Improved water use efficiency of warm- and cool-season turfgrass resulting in pre-drought stress conditioning.

**NOTICE TO USER:** Response to Cutless 50W may vary within turfgrass species due to the large number of cultivars and varieties available. Neither the manufacturer nor seller has determined if this product can be used safely or effectively on species not mentioned on this label. For species not listed on this label the user should apply this product to a small test area to determine growth response and desired level of growth regulation prior to large scale applications.

## Use Restrictions

- **DO NOT** apply to putting greens other than those where bentgrass is the desired turf species.
- **DO NOT** apply to bermudagrass putting greens, including those which are overseeded.
- **DO NOT** apply this product to sod farms, turfgrasses grown for seed, including plants or plant materials grown for sale or research purposes.
- **DO NOT** apply to shrubs, bedding plants, and/or food plants.
- **DO NOT** apply during prolonged periods of temperature (heat or cold) or moisture stress. Also avoid applications during periods of extreme disease and insect pressure.
- **DO NOT** apply to saturated soils or when a significant moisture event is anticipated. This product may accumulate in low lying areas and cause prolonged and excessive growth regulation in those areas.
- **DO NOT** apply to areas where *Poa annua* is the desired turfgrass species or areas that contain >80% *Poa annua*.
- **DO NOT** apply to turf used for livestock production.
- The maximum number of annual applications is determined by the sum of the rates applied, not to exceed 3.0 lbs. a.i./A/year or 6.0 lbs. Cutless 50W/A (6-24 applications/year).
- **Chemigation: DO NOT** apply through any type of irrigation system.
- **DO NOT** apply by Aerial application.
- **DO NOT** apply more than 0.275 lb. a.i./A in a single application on golf course roughs.

## Use Precautions for Perennial Turfgrasses

- For best results, delay applications to newly seeded turfgrasses until turf is well established and actively growing.
- For best results, delay application until 6 to 8 weeks after turfgrass sprigging or laying sod. Turfgrass should be well established and actively growing prior to application.
- Additional turfgrass growth regulation may occur when Cutless 50W is tank mixed or used in conjunction with demethylation inhibitor (DMI) or sterol inhibiting fungicides.

## Application Timing

Apply Cutless MEC to actively growing turfgrass. Make spring applications after resumption of active seasonal growth of turfgrass. Scheduler the final application of the season a minimum

of 4 weeks before the onset of inactive grass growth or winter dormancy. To avoid delayed spring transition of bermudagrass, apply to overseeded turfgrasses in dormant bermudagrass stands 4 weeks prior to expected bermudagrass green-up.

### **Irrigation**

Water-in within 24 hours of application to limit surface movement, but not to the point of runoff. To prevent product runoff, time applications to allow for watering-in and maximum absorption into treated turf prior to a rain event. Avoid mowing treated turfgrass areas until after rainfall or irrigation occurs.

### **Turf Color and Post-Treatment Turf Management**

Treated turfgrass may appear darker green in color. This color change, which appears 1 to 2 weeks after treatment, may persist an additional 3 to 6 weeks. Treated areas should be managed to encourage the growth of a healthy vigorous turf.

### ***Poa annua* (Annual Bluegrass) Conversion to Perennial Turfgrasses**

Applications of Cutless 50W followed by management practices designed to encourage vigorous growth of perennial turfgrasses can reduce the *Poa annua* (annual bluegrass) competition in cool-season turf and increase the cover of more desirable perennial species. *Poa annua* is more sensitive to Cutless 50W treatments and is therefore more strongly suppressed than perennial turfgrass species. Discoloration of *Poa annua* can be expected from treatments that provide a desired level of growth regulator activity in perennial grass species. This effect becomes visible 7 to 10 days after treatment and lasts 3 to 6 weeks. The degree of discoloration will be proportional to the *Poa annua* composition of the turf. Application in conjunction with soluble nitrogen fertilizers will also minimize discoloration. Application timing, rates, and precautions for perennial grass conversion through selective reduction of *Poa annua* are provided in the *Poa annua Conversion to Perennial Turfgrasses* section of this label.

## **APPLICATION DIRECTIONS**

### **Mixing Directions**

Add Cutless 50W water-soluble packets to a spray tank half filled with clean water. Begin agitation and allow the packets to dissolve and release their contents into the circulating water. Allow sufficient mixing time to ensure complete dispersion and mixing of Cutless 50W. Finish filling the spray tank. Continue agitation throughout the spraying operation to ensure uniform application. Cutless 50W should be applied using a boom-type sprayer with bypass and/or mechanical agitation calibrated to deliver 40 to 200 gallons/acre of spray solution (1 to 4.6 gallons/1000 ft<sup>2</sup>). In-line strainers and nozzle screens should be 50 mesh or larger. The use of a coloring agent to mark areas already sprayed is suggested for uniform application without skips and overlaps. Performance may be improved with the addition to the spray mix of a readily available nitrogen (N) source at 0.125 to 0.5 lbs N/1000 ft<sup>2</sup> or iron (Fe) at suggested label rates.

### **Cutless 50W + Tank Mixtures**

Cutless 50W can be tank mixed and is compatible with most commonly-used pesticides and foliar nutrient products. However, test compatibility of this product with tank mix partners before use.

**NOTE:** Test the compatibility of this product in any tank-mix combination before use. To determine the physical compatibility with other products, use a jar test as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure sequence for adding required ingredients to the spray tank.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### **SPRAY DRIFT MANAGEMENT**

Applications must be made only when there is no hazard for spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Applicators are required to use a medium or coarser droplet size (according to ASABE standard 572). **When using ground application equipment, apply with nozzle height no more than 2 feet above the target plants. Do not apply when wind speeds exceed 10 miles per hour at the application site. Do not apply during temperature inversions.**

### **SPRAY DRIFT ADVISORIES**

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

#### **Importance of Droplet Size**

An effective way to reduce spray drift is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions. See Wind, Temperature, and Humidity, and Temperature Inversions sections of this label.

#### ***Techniques for Controlling Droplet Size – Ground Boom***

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use a higher-capacity nozzle instead of increasing pressure.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

#### **Boom Height**

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.



**Wind**

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Avoid applications during gusty or windless conditions.

Note: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity**

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

**Temperature Inversions**

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Shielded Sprayers**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**RUNOFF PREVENTION**

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when excessive rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

**GROWTH REDUCTION OF PERENNIAL TURFGRASS SPECIES**

A multiple application program using this product provides growth reduction of perennial turfgrass species resulting in decreased mowing frequency and turfgrass clippings. For cool-season grasses, begin initial applications in early spring following resumption of active growth. For warm-season grasses, begin initial applications when the grass has completely recovered from winter dormancy and is growing vigorously. For both warm- and cool-season grasses, discontinue applications a minimum of 4 weeks before the onset of inactive grass growth or winter dormancy. Use lower rate range in early spring and late fall applications to avoid excessive growth regulation. Refer to Table 1 for rates for growth regulation of perennial turfgrass species.

<b>TABLE 1</b>			
<b>Rate Ranges for Growth Regulation of Perennial Turfgrass Species Using a Multiple Application Program†</b>			
<b>Turfgrass Species</b>	<b>lbs. Cutless</b>	<b>oz. Cutless</b>	<b>Treatment</b>

	<b>50W/A (lbs. a.i./A)</b>	<b>50W/A</b>	<b>Interval</b>
<b>Cool-Season Turfgrasses</b>			
Bentgrass (golf course fairway)	0.25 - 1.00 (0.125-0.5)	4 - 16	2 to 6 weeks
Bentgrass putting greens	0.125 - 0.50 (0.063-0.25)	2 - 8	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass	0.50 - 1.00 (0.25-0.5)	8 - 16	3 to 6 weeks
<b>Warm-Season Turfgrasses</b>			
Seashore Paspalum	0.25 - 1.00 (0.125-0.5)	4 - 16	3 to 6 weeks
Tifway, TifSport, and GN-1 Bermudagrass <sup>††</sup> , Zoysiagrass <sup>†††</sup>	0.25 - 0.75 (0.125-0.375)	4 - 12	3 to 6 weeks

<sup>†</sup> Apply in early spring following resumption of active growth of the grass. Fall applications must be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

<sup>††</sup> California: Apply at 24.6 to 49.2 fl. oz./A

<sup>†††</sup> Not in late summer/fall

### **APPLICATION INFORMATION – CUTLESS 50W TANK MIXES WITH TRINEXAPAC-ETHYL**

Tank mixing Cutless 50W with products containing trinexapac-ethyl can provide enhanced growth suppression and improved turfgrass quality of perennial turfgrass species, versus using either product alone. Plant physiological advantages of tank mixing these two PGRs include:

1. Different site of action within the gibberellic acid (GA) biosynthesis pathway; and
2. Difference in plant site of uptake.

PGR absorption via the foliage (Class A; trinexapac-ethyl) and roots (Class B; Cutless 50W) maximizes plant uptake of each material ensuring sufficient active ingredient is available for GA inhibition.

Blocking GA biosynthesis early and late in the pathway potentially regulates GA more efficiently than higher application rates of individual compounds. Turfgrass response from Cutless 50W tank mixes with reduced rates of products containing trinexapac-ethyl may be greater than with normal use rates of either product applied alone. This combination of plant growth regulators and its use are protected by United States Patent No. 7,135,435 and 9,198,417. Additional patent rights pending.

Applications of Class A plus Class B PGRs can reduce the flush of vegetative growth that occurs at the end of the growth regulation period when only a single PGR class is used.

Desirable growth regulation during early spring or late fall when turfgrass growth and vigor are reduced may be obtained with lower rates of both products.

When tank mixing with products containing trinexapac-ethyl, rainfall or irrigation must be delayed at least 1 hour after application or until product has dried on the leaf surface but should occur within 24 hours after application.

**NOTICE TO USERS:** To the extent consistent with applicable law, this label makes no warranties concerning the performance of Class A PGRs, including trinexapac-ethyl. Read

and follow all label directions including Directions for Use, Precautionary Statements, and Restrictions and Limitations for trinexapac-ethyl.

<b>TABLE 2</b>				
<b>Rate Ranges for Cutless 50W Plus Class A PGR Tank Mixes</b>				
<b>Turfgrass Species</b>		<b>oz. Cutless 50W/A (lb a.i./A)</b>	<b>Trinexapac-ethyl</b>	<b>Treatment Interval</b>
<b>Cool-Season Turfgrasses</b>				
Bentgrass (golf course fairway)		2 – 8 (0.063- 0.25)	½ labeled use rate	2 to 6 weeks
Kentucky Bluegrass/ Perennial Ryegrass		4 – 8 (0.063- 0.25)	½ labeled use rate	3 to 6 weeks
Bentgrass putting greens		2 - 4	½ labeled use rate	2 to 6 weeks
<b>Warm-Season Turfgrasses</b>				
Seashore Paspalum and Zoysiagrass		4 - 8 (0.125- 0.25)	½ labeled use rate	3 to 6 weeks
Tifway, TifSport, and GN-1 Bermudagrass†		2 - 8 (0.063- 0.25)	½ labeled use rate	3 to 6 weeks

† Do not use on bermudagrass putting greens.

### **POA ANNUA (ANNUAL BLUEGRASS) CONVERSION TO PERENNIAL TURFGRASSES**

A growth regulation program using Cutless 50W provides *Poa annua* suppression and cool-season turf conversion. This program provides a gradual perennial grass conversion reducing *Poa annua* populations over one to several years. To maximize seedling establishment, use only a low rate of this product within one (1) week prior to and/or 1 week after interseeding or overseeding of perennial turfgrasses.

Refer to Table 3 for application rates and treatment intervals.

#### ***Bentgrass (Golf Course Fairway)***

Apply Cutless 50W to fairway height bentgrass in early spring following resumption of active growth of the grass. Repeat applications may be made until late summer or early fall. Normal management practices such as fertilization, aeration and interseeding/overseeding will encourage growth of bentgrass.

#### ***Bentgrass Putting Greens***

Annual turfgrass species such as *Poa annua* are more strongly regulated by Cutless 50W. Careful use of this product on putting greens with a high percentage of *Poa annua* can shift the competitive balance between bentgrass and *Poa annua* to favor bentgrass. Follow normal management practices such as fertilization, aeration and interseeding/overseeding that encourages growth of bentgrass. Use of this product on bentgrass greens may increase putting speed without reducing the height of cut.

**Bentgrass Greens with less than 50% *Poa annua***

Apply Cutless 50W to bentgrass as part of an overall greens management program. An initial application may be made in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Repeat applications should be made through early fall.

**Bentgrass Greens with more than 50% *Poa annua***

Apply Cutless 50W to bentgrass greens in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Repeat applications may be made through early fall.

**Kentucky bluegrass, Perennial ryegrass**

Apply Cutless 50W to Kentucky bluegrass and perennial ryegrass in early spring following resumption of active growth of the grass. Repeat applications may be made until late summer or early fall. For bermudagrass fairways overseeded with perennial ryegrass delay applications until perennial ryegrass is well established (4 weeks after germination). To avoid delayed spring green-up of bermudagrass, conduct the final spring application a minimum of 4 weeks prior to expected bermudagrass green-up. Normal management practices such as fertilization, aeration and interseeding/overseeding will encourage growth of bluegrass and/or ryegrass.

<b>TABLE 3</b>				
<b>Rate Ranges for <i>Poa annua</i> (Annual Bluegrass) Conversion to Perennial Turfgrasses with Cutless 50W Using a Multiple Application Program<sup>†</sup></b>				
<b>Turfgrass Species</b>	<b>% <i>Poa annua</i></b>	<b>Initial spring application</b>	<b>Repeat applications</b>	
		<b>oz. Cutless 50W/A (lb. a.i./A)</b>	<b>oz. Cutless 50W/A (lb. a.i./A)</b>	<b>Treatment Interval</b>
Bentgrass (golf course fairway)	0 - 80%	8 – 16 (0.25-0.5)	4 – 16 (0.125-0.5)	2 to 6 weeks
Bentgrass Putting Greens	< 50%	4 – 8 (0.125-0.25)	4 – 8 (0.125-0.25)	2 to 4 weeks
Bentgrass Putting Greens	> 50%	4 (0.125)	4 – 8 (0.125-0.25)	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass Fairways <sup>††</sup>	0-80%	12 – 16 (0.375-0.5)	8 – 16 (0.25-0.5)	3 to 6 weeks

<sup>†</sup> Apply in early spring following resumption of active growth of the grass. Fall applications should be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

<sup>††</sup> For perennial ryegrass overseeded fairways, delay applications until perennial ryegrass is well established (3-4 weeks after germination).

**POA ANNUA (ANNUAL BLUEGRASS) CONVERSION TO PERENNIAL TURFGRASSES - ALTERNATIVE SPRING/FALL APPLICATION PROGRAM**

The alternative spring/fall application program selectively suppresses *Poa annua* growth in perennial turfgrass species. Injury or discoloration of *Poa annua* is expected within 7 to 10 days of application. Cultural practices such as fertilization, aeration and interseeding/overseeding should be performed in conjunction with Cutless 50W applications to encourage growth of the perennial turfgrass species.

Refer to Table 4 for application rates and treatment intervals.

### **Bentgrass or Kentucky bluegrass/perennial ryegrass fairways with less than 50% *Poa annua***

An alternative spring/fall treatment program for management of *Poa annua* in fairways containing less than 50% *Poa annua* is to apply Cutless 50W following resumption of active bentgrass or Kentucky bluegrass/perennial ryegrass growth in the spring. A second application may be made in the early fall. Interseed/overseed with bentgrass or Kentucky bluegrass/perennial ryegrass 3 to 4 weeks following early fall application.

### **Bentgrass or Kentucky bluegrass/perennial ryegrass fairways with more than 50% *Poa annua***

Apply Cutless 50W in late summer or early fall. Interseed/overseed with bentgrass or Kentucky bluegrass/perennial ryegrass 3 to 4 weeks following late summer or early fall application. Make an additional Cutless 50W application the following spring after resumption of active growth of bentgrass or Kentucky bluegrass/perennial ryegrass.

<b>Rate Ranges for <i>Poa annua</i> Conversion to Perennial Turfgrasses Using an Alternative Spring/Fall Application Program</b>		
<b>Turfgrass Species</b>	<b>% <i>Poa annua</i></b>	<b>Spring/Fall application<sup>†</sup> oz. Cutless 50W/A (lb. a.i./A)</b>
Bentgrass or Kentucky bluegrass/perennial ryegrass fairways	< 50%	16 (0.5) in spring; 16 (0.5) in fall
	> 50%	16 (0.5) in late summer or early fall; 16 (0.5) the following spring

<sup>†</sup> Apply in early spring following resumption of active growth of the grass. Fall applications should be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

### **DOLLAR SPOT (*CLARIREEDIA JACKSONII*) SUPPRESSION BY CUTLESS 50W IN CREEPING BENTGRASS**

The active ingredient in Cutless 50W is from the pyrimidine class of chemistry which is structurally similar to pyrimidine fungicides that provide dollar spot control. Programmed applications of this product for turf growth suppression or *Poa annua* conversion have also been shown to suppress dollar spot incidence in creeping bentgrass fairways, greens and tees. Research results have shown Cutless 50W applications at labeled rates and application intervals can significantly reduce dollar spot incidence and populations when compared to untreated control plots. This product should not be used to replace labeled fungicides for the control of dollar spot; rather programmed use may result in longer or improved control of dollar spot in conjunction with conventional fungicides, or delays in the appearance of dollar spot disease, thus leading to the potential for an overall reduction in annual fungicide use.

## EDGING AND BANDING APPLICATIONS FOR GROWTH REGULATION OF PERENNIAL TURFGRASS SPECIES

Cutless 50W can be applied to turfgrass in edging and banding applications along the perimeter of lawns, landscape beds, sidewalks, curbs, parking lots, driveways, posts, mailboxes, building structures, gravestones, fences, roadsides, medians and guardrails to reduce the frequency of trimming and edging. For best results application should occur no more than 3 days after turfgrass has been trimmed to desired height. Apply this product in a 6 inch band width with a single nozzle sprayer. Repeat at 8 to 12 week intervals or as needed for turf growth regulation.

<b>TABLE 5</b>	
<b>Edging/Banding Rates for Growth Regulation of Perennial Turfgrass</b>	
<b>Turfgrass Species</b>	<b>oz. Cutless 50W/A (lb. a.i./A)</b>
<b>Cool-Season Turfgrasses</b>	
Bentgrass	16 – 32 (0.5-1.0)
Kentucky Bluegrass; Perennial Ryegrass; Tall Fescue	24 – 48 (0.75-1.5)
<b>Warm-Season Turfgrasses</b>	
328 Hybrid Bermudagrass	12 – 16 (0.375-0.5)
419 Hybrid Bermudagrass; Seashore Paspalum; St. Augustinegrass; Zoysiagrass	16 – 32 (0.5-1.0)
Common Bermudagrass	24 – 48 (0.75-1.5)

## BACKPACK SPRAYER DOSAGE CHART

For use of Cutless 50W as an edging/banding application in backpack sprayers having tank capacity of 3 to 5 gallons and calibrated to deliver 20 to 40 gallons per acre, refer to Table 6 to determine ounces of Cutless 50W required to achieve desired rate per acre. Use Edging/Banding Rates Table (Table 5) to select Cutless 50W rate for desired turfgrass species.

<b>TABLE 6</b>			
<b>Backpack Sprayer Dosage Directions</b>			
<b>Backpack tank capacity (gallons)</b>	<b>Spray volume (gallons/A)</b>	<b>Linear ft treated †</b>	<b>oz. Cutless 50W added to backpack spray tank</b>
3	20	13,068	2 oz
	30	8,712	
	40	6,534	
4	20	13,068	4 oz
	30	8,712	
	40	6,534	
5	20	17,424	2 oz
	30	11,616	
	40	8,712	

	20	17,424	4 oz
	30	11,616	
	40	8,712	
5	20	21,780	2 oz
	30	14,520	
	40	10,890	
	20	21,780	4 oz
	30	14,520	
	40	10,890	

† Assuming band of spray application is 6-inches wide.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Excess pesticide should be used according to label directions or disposed of at an approved waste disposal facility.

#### Container Handling

#### Nonrefillable Container Disposal (non-rigid, any size):

**(Fiber Carton/Drum and Foil Overpack):** Do not reuse or refill this container. Offer for recycling if available. If recycling not available for fiber carton/drum, then dispose of both foil overpack and drum/carton in a sanitary landfill or incinerator.

**Warranty Disclaimer:** SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

**Misuse:** Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



[Base label for Non-refillable, Non-rigid containers, any size]



# Cutless® 50W

## Turf Growth Regulator

### Active Ingredient

Flurprimidol:  $\alpha$ -(1-methylethyl)- $\alpha$ -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol ..... 50.0%

Other Ingredients ..... 50.0%

**TOTAL** ..... 100.0%

Contains 0.50 pound active ingredient per one (1) U.S. pound of product.

### Keep Out of Reach of Children

## CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS



**Caution. Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid breathing dust or spray mist and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.**

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Excess pesticide should be used according to label directions or disposed of at an approved waste disposal facility.

**Container Handling**

**Nonrefillable Container Disposal (non-rigid, any size):**

**(Fiber Carton/Drum and Foil Overpack):** Do not reuse or refill this container. Offer for recycling if available. If recycling not available for fiber carton/drum, then dispose of both foil overpack and drum/carton in a sanitary landfill or incinerator.

**Refer to label booklet for additional Precautionary Information and Directions for Use.**

**Notice:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer and Misuse statements in label booklet*. If terms are unacceptable, return at once unopened.**

EPA Reg. No. 67690-15  
FPL20190724

EPA Est. No. \_\_\_\_\_  
[P/N] \_\_\_\_\_

Cutless is a registered trademark of SePRO Corporation.

**SePRO Corporation 11550 N. Meridian Street, Suite 600 Carmel, IN 46032, U.S.A.**

**[Turf Growth Regulator]  
refillable)**

**Net Contents \_\_\_\_\_ (Non-**

[Text accessed through the weblink / QR code. This is NOT part of the printed label]

---

## TERMS AND CONDITIONS OF USE

---

If terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label as well as the *Inherent Risks of Use* and *Limitation of Remedies* statements below are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Misuse*, *Inherent Risks of Use*, and *Limitation of Remedies*.

---

## INHERENT RISKS OF USE

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including use under conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), the presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer and/or user of the product.

---

## LIMITATION OF REMEDIES

---

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use*, *Inherent Risks of Use* and *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use*, *Inherent Risks of Use* and *Limitation of Remedies* in any manner.