

67690-20

9/15/2011

1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Laurent Mézin
SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, Indiana 46032-4565

SEP 15 2011

Subject: Topflor Ornamental Plant Growth Regulator
EPA Reg. No. 67690-20
Label Revisions
EPA Decision Number 451701

Dear Mr. Mézin:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment.

If you have any questions, please contact Erin Malone by phone at (703) 347-0253 or via email at malone.erin@epa.gov.

Sincerely,

A handwritten signature in black ink, which appears to read "Shaja B. Joyner", is written over a horizontal line.

Shaja B. Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7504P)

Enclosure: Master Label Stamped "Accepted"

2/19

Topflor®
Ornamental Plant Growth Regulator
EPA Reg. No. 67690-20

S: 903125

Registration Notes:

Final printed labeling based on label submitted to EPA with FPL070111. Update per EPA comments of 07 and 12 September 2011. Other changes as noted below.

FPL20110912 Clean

Label Notes:

General Label changes:

1. Updated FPL date.


EPA requested label changes:

1. Page 9, changed the 5th statement to say, "If you have questions about calibration, you should contact State Extension service specialists, equipment manufacturers, or other experts."
2. Page 10, Added the following statement as the last bulleted point under Chemigation Systems Connected to Public Waters Systems, "Do not apply when wind speed favors drift beyond the area intended for treatment."

3/19

Topflor®, 67690-20

(Front label for all containers):



Topflor®
Ornamental Plant Growth Regulator

A growth regulator for use on ornamental plants grown in containers in commercial nurseries, greenhouses and shadehouses.

IN NEW YORK STATE, TOPFLOR CAN BE USED FOR GREENHOUSE APPLICATIONS ONLY.

Active Ingredient:

flurprimidol: α -(1-methylethyl)- α -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol..... 0.38%

Other Ingredients 99.62%

Total 100.00%

Contains 15 grams of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Refer to inside of label booklet for additional precautionary information and directions for use, including first aid and storage and disposal.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use* and *Limitation of Remedies* inside label booklet.

Shake well before using.

EPA Reg. No. 67690-20
FPL20110912

EPA Est. No. _____
SPC - _____

SePRO Corporation 11550 N. Meridian Street, Suite 600 Carmel, IN 46032, U.S.A.

Ornamental Plant Growth Regulator

Net Contents _____
[Refillable or Nonrefillable container _____]

4/19

(Label Booklet ALL containers):

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye and skin irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call InfoTrac toll free at 1-800-535-5053 .	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coveralls, long-sleeved shirt and long pants;
- Chemical-resistant gloves made of any waterproof material; and
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Shake well before using.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls;
- Chemical-resistant gloves made of any waterproof material;
- Shoes plus socks; and
- Long sleeved shirt and long pants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity \leq 5 gallons) as follows:

Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap.

Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip.

Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity $>$ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

PRODUCT INFORMATION

Topflor Plant Growth Regulator is for use on ornamental plants grown in containers in nurseries, greenhouses and shadehouses. Use of Topflor effectively reduces internode elongation through the inhibition of gibberellin biosynthesis, resulting in a more desirable compact plant. Topflor has been shown to increase the quality of plants even in the absence of growth reduction. Some of these desirable qualities include darker leaf color, higher chlorophyll content, greater leaf thickness, stronger stems, and decreased water loss. When used as directed, Topflor produces no phytotoxic effects.

Do not reuse pots, trays, or other containers that previously were used in the production of a crop that was treated with Topflor.

In New York State, Topflor can be used for greenhouse applications only.

FACTORS AFFECTING PLANT RESPONSE TO TOPFLOR

There are many factors that can affect a plant's response to the application of Topflor. They include cultivar, application technique, environmental conditions, cultural practices and container size. Therefore, the amount of Topflor that is required for the desired plant height may vary.

- **CULTIVARS OR VARIETIES** within a given plant species may respond differently to Topflor. Varieties that are taller or more vigorous generally require more Topflor than naturally short or less vigorous varieties. Growers may consult university research and extension specialists and plant or seed suppliers for vigor and other growth characteristics for newly released varieties.
- **ENVIRONMENTAL CONDITIONS** can also strongly influence the response to Topflor and, therefore, the amount of product applied. Growers in warm climates may need to use higher rates and/or more applications compared to those in cooler climates. The Topflor rate as well as number of applications may also vary depending on the time of year, with higher rates and/or more applications needed during warmer months.
- **CULTURAL PRACTICES** may affect the plant's response to Topflor. Plants that are grown at close spacing or in small pots and using high water and fertility levels may require higher rates of Topflor to achieve the desired response. The effectiveness of a Topflor drench application may be reduced in root media that utilizes a high amount of pine bark.

MIXING INSTRUCTIONS

Shake well before using.

The sprayer must be clean and not contaminated with other chemicals. Use the Dilution Table (Table 1) to determine the amount of Topflor and water needed for the required concentration. Fill the spray tank with half the required amount of water. Measure the desired Topflor volume accurately and add it to the tank. Fill tank with the remaining amount of required water. Agitate the Topflor and water mixture frequently to assure uniform distribution during application.

TABLE 1: TOPFLOR DILUTION RATES				
ppm a.i. desired concentration	ml per gallon solution	fl. oz. per gallon solution	fl. oz. per 10 gallon solution	fl. oz. per 100 gallon solution
0.5	0.48	0.02	0.16	1.61
1	0.96	0.03	0.32	3.23
2	1.91	0.06	0.65	6.46
3	2.87	0.10	0.97	9.69
4	3.82	0.13	1.29	12.92
5	4.78	0.16	1.61	16.15
10	9.55	0.32	3.23	32.30
15	14.33	0.48	4.84	48.44
20	19.10	0.65	6.46	64.59
25	23.88	0.81	8.07	80.74
30	28.65	0.97	9.69	96.89
35	33.43	1.13	11.30	113.04
40	38.20	1.29	12.92	129.18
50	47.75	1.61	16.15	161.48
60	57.30	1.94	19.38	193.77
80	76.41	2.58	25.84	258.37
100	95.51	3.23	32.30	322.96
200	191.02	6.46	64.59	645.91

APPLICATION TECHNIQUES

Plants absorb Topflor through foliage, stems, and roots. **Topflor may be applied as a spray, drench or chemigation to achieve the desired plant height control.** Use industry standard application equipment, which may include backpack sprayers, low-pressure hand wand drench applicators, or other similar equipment. Additionally, standard chemigation equipment and practices may also be used. Multiple or split applications may allow greater treatment flexibility, more uniform growth regulation, and safety from over-application and may be, therefore, desirable.

SPRAY APPLICATIONS

Topflor applied as a foliar spray is absorbed through plant foliage and stems. Additional growth regulation will result from root uptake of Topflor reaching the root medium as runoff from foliar treatments or over-spray.

Dilute Topflor to the desired concentration (a.i.) using Table 1.

When applying as a spray, the following should be noted:

- Do not use additional wetting agents in combination with Topflor as crop injury may occur.
- The spray technique used should provide consistent and uniform coverage over all treated plants. Uneven application or over-application may result in irregular or excessive growth control.
- Adequate spray volume should be used to thoroughly wet the plant foliage. The spray volume that drips down to the stem or media may be desirable as it will be

taken up by the stems and roots increasing the effectiveness of Topflor. However, too much runoff into the media may result in excessive height control.

- Apply uniformly at a rate of 1 gallon of spray per 200 sq. ft. of growing area, regardless of plant spacing. For small plants in small containers or plug trays that are closely spaced, use 0.5 - 1 gallon of spray per 200 sq. ft. of growing area. For larger plants with a well-developed canopy, a spray volume of 1.5 gallons per 200 sq. ft. of growing area is recommended.
- Typical foliar application rate is 0.5 ppm - 80 ppm a.i. (varies by cultivar), applied in 1 gallon of spray mix over 200 square feet.
- Do not allow spray drift to contact non-target plants.

DRENCH APPLICATIONS

Topflor applied as a drench provides treatment accuracy for consistently uniform results. Topflor is readily absorbed by the roots and translocated to the terminals. Root medium should be moist, but not wet at the time of treatment. Best results are obtained when moisture content allows the drench treatment to become well distributed and retained entirely within the pot. This may be achieved by watering the plants the day before treating. Response may be variable if part of the drench solution is lost to flow-through or if root medium is too dry to allow for even distribution of the treatment, especially when multiple cuttings are in the same container. Generally, a volume of 2 fl. oz. (59 ml) is required to treat a 4-inch pot or 4 fl. oz. (118 ml) for treatment of a 6-inch pot (Table 2). The typical application rate is 1 gallon of drench solution (typically 0.25 ppm - 10 ppm as a.i.) (varies by cultivar), per 32 six inch potted plants. Dilute Topflor to the required concentration (a.i.) using the method described in Table 1. When applying as a drench, the use of pine bark in root media may reduce the effectiveness of drench treatments.

TABLE 2: DRENCH VOLUME APPLICATION GUIDELINES							
Pot Diameter (Inches)	Drench Volume		mg a.i./pot from solutions mixed at the following ppm[†]				
	fl oz/pot	ml/pot	0.5 ppm	1 ppm	2 ppm	3 ppm	4 ppm
4	2	59	0.029	0.059	0.118	0.177	0.236
5	3	89	0.044	0.089	0.177	0.266	0.356
6	4	118	0.059	0.118	0.236	0.355	0.473
8	10	296	0.148	0.295	0.590	0.888	1.183
10	25	740	0.370	0.740	1.480	2.220	2.960
12	40	1184	0.592	1.178	2.360	3.543	4.727

[†] Refer to Table 1 for mixing instructions

NOTE: The listed drench volumes were based on the soil capacity of a common 6-inch "azalea-type" pot. Extrapolating the rate for this type pot to smaller or larger containers may not be correct for the total drench volume, but should only be used as a guideline. The user must determine the appropriate rate and drench volume needed to achieve the desired result, based on both pot size and root medium used.

CHEMIGATION

Not for use in California or New York.

Pesticide labels contain directions for use which are necessary for effecting the purpose for which the product is intended and to protect health and the environment. The following information is intended to decrease environmental risks of pesticide contamination of ground water and will decrease direct human exposure to pesticide treated irrigation water by providing appropriate directions for use.

Apply this product only through pressurized drench (flood), sprinkler, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation systems.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment

PRESSURIZED DRENCH (FLOOD) SYSTEM

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlocked to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump. (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

SPRINKLER (SPRAY) CHEMIGATION

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

DRIP (TRICKLE) CHEMIGATION

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

PRODUCT INFORMATION

Pesticide supply tanks are recommended for the application of these products. See label instructions for dilution use rates and timing of applications. Agitate prior to use.

Since the material is used in an injections proportioner the pesticide is to be applied continuously for the duration of the water application.

APPLICATION RATES

The amount of Topflor required for an optimum growth response depends upon several factors: desired height, duration of growth response and degree of control, pot size, stage of growth, method of application, season and cultivar response. Species-specific cultural practices such as watering, potting media, fertilization, temperature and light conditions also affect the growth response to a given dosage. Therefore, growers should establish specific application rates based on small-scale treatments under actual use conditions and keep records as to plant species and cultivar sensitivity before Topflor is applied to a large number of plants. The rates listed on this label are rate ranges and should be used only as a guideline.

For spray, drench and chemigation applications, do not exceed the maximum rate of 0.36 lbs a.i./A for single applications. Do not exceed more than 3.0 lbs a.i./A/year. Rate (lbs a.i./A) will determine the maximum number of seasonal applications allowed not to exceed 3.0 lbs a.i./A/year. If required, repeat applications to the same crop may be applied at 5 to 21 day intervals.

NOTICE TO USER: Plant tolerance to Topflor has been found to be acceptable in research trials for the general plant species listed on this label. However, due to the large number of species of ornamental and nursery plants and their associated varieties and cultivars and due to variable growing conditions, it is impossible to test every plant and variety or cultivar for tolerance to Topflor. The Manufacturer has not determined whether Topflor can be used safely on all ornamental plants. Whereas Topflor has been shown to be safe and effective in a limited number of research trials on certain varieties or cultivars of the plant types listed, the professional user should determine if Topflor can be used safely prior to commercial use. Prior to wide-scale use, users should conduct small-scale tests under local growing conditions using the general guideline rates listed below. For species and their varieties or cultivars not specifically listed on the label, the user assumes all risk from phytotoxicity or unacceptable growth effects.

Topflor is effective in controlling the height of most ornamental crops. The use and rate specifications for the species that follow should be a starting point in determining the best rate for your specific cultural and environmental growing conditions. Before Topflor is applied to a large number of plants, read and understand the section titled *Application Rates*.

Bedding Plants

Topflor is effective on a wide range of bedding plants for height control. See Table 3 for application rate guidelines for a variety of common bedding plants. For specific plants not identified on Table 3, use 0.5 - 80 ppm (a.i.) spray as the recommended general guideline rates for plugs to finished bedding plants.

TABLE 3: SPRAY RATE RANGE TRIAL GUIDELINES FOR SOME BEDDING PLANTS [†]			
Plant	Rate Range (ppm a.i.)	Plant	Rate Range (ppm a.i.)
<i>Ageratum</i>	20 - 60	<i>Nemesia</i> ¹	10 - 15
<i>Celosia</i> , flame	10 - 40	<i>Osteospermum</i>	20 - 60
<i>Coleus</i> , seed	20 - 40	<i>Pansy</i>	2.5 - 7.5
<i>Geranium</i> , zonal	15 - 25	<i>Petunia</i>	20 - 60
<i>Impatiens</i>	20 - 60	<i>Salvia</i>	20 - 80
<i>Marigold</i>	20 - 60	<i>Vinca</i> ²	2.5 - 10

[†] These rate ranges were determined largely under mid-Atlantic conditions using medium-vigor cultivars. Rates should be adjusted to reflect the need for higher rates for vigorous cultivars and in the Sunbelt Region and lower rates in the Northern Belt Region. Topflor is not recommended for use on fibrous begonia. Overly stunted plants can result if they receive spray drift from applications to surrounding crops.

¹ *Nemesia*: A rate of 10 - 15 ppm (a.i.) is recommended for a single spray application. Alternatively, apply twice at 5 ppm (a.i.) with the second application made two weeks after the first.

² Annual vinca (periwinkle): Growers should note that black spotting may result from higher rates of spray application, especially at high temperatures.

Drench: Apply to uniformly moist root media. Apply at a solution concentration at general guideline rates of 0.25 - 4 ppm (a.i.) at the listed volume per pot (see Table 2). Rates for a specific plant species/cultivar and set of use conditions should be

determined in small-scale treatments prior to large-scale applications. The user should determine his/her own optimum rates noting that the above-listed rate range encompasses production in the warmer and cooler climates.

Bedding Plant Plugs

Spray applications of Topflor may be used to control the height of certain aggressive species of bedding plant plugs. Over-regulation and poor performance after transplanting the plug can occur if rates are too high or if used on overly-responsive crops. Due to the responsiveness of bedding plant plugs to Topflor, it should only be used on *Geranium*, *Impatiens*, Marigolds, and *Petunia*. Do not use Topflor on sensitive bedding plant plugs including *Begonia*, Pansy, *Salvia*, and *Vinca*.

Differences in environmental factors and cultural practices during plug production can have a dramatic impact on plant growth regulator rates and results. Growers should conduct trials on a small scale under their growing conditions to determine the optimum rate that will provide proper efficacy while ensuring desirable growth and crop performance in the finishing stage. A trial spray application of 1 - 8 ppm (a.i.) is the suggested general guideline rate, which should be adjusted based upon trial results and user observations. Application timing is suggested after the development of the first 1 - 2 true leaves. Application volume generally should be 0.5 - 1 gallon per 200 sq. ft. of treatment area. Do not apply as a drench to bedding plant plugs.

Bulb or Fibrous Root Crops

Topflor is very effective on most bulb crops. Topflor is more effective when applied as a drench rather than a spray on most bulb crops (see Table 4). For bulb species not listed in Table 4, the grower should determine the optimum rate for the species grown under their cultural and environmental conditions by running trials on a small number of plants. In general, apply a soil drench to uniformly moist rooting media approximately 2 weeks after planting when new growth reaches 1 inch.

TABLE 4: RATE RANGE TRIAL GUIDELINES FOR SOME BULB CROPS [†]		
Plant	Spray Rate Range (ppm a.i.)	Drench Rate Range
		mg a.i./pot
Calla Lily	ND	1 - 2.5
Canna Lily	50 - 80	ND
Dahlia	NR	0.5 - 2
Oriental Hybrid Lily	ND	0.25 - 0.5
"Stargazer"		
Tulip ¹	80 - 100	0.5 - 1
Hyacinth	ND	0.5 - 1
Caladium	NR	0.5 - 2

[†] These rate ranges were determined largely under mid-Atlantic conditions using medium-vigor to vigorous cultivars. Rates should be adjusted to reflect the need for higher rates in the Sunbelt Region and lower rates in the Northern Belt Region or for less vigorous cultivars.

¹ Tulip: spray for control of post-harvest stretch.

ND = Rates for this application technique have not been determined.

NR = Not a recommended use.

Flowering /Foliage Potted Plants

Topflor is effective when applied as a spray or drench on a wide variety of flowering and foliage plants.

TABLE 5: RATE RANGE TRIAL GUIDELINES FOR SOME FLOWERING/FOLIAGE PLANTS[†]

Plant	Spray Rate Range (ppm a.i.)	Drench Rate Range
		mg a.i./pot
<i>Campanula</i>	10 - 30	ND
Cape Primrose (<i>Streptocarpus hybridus</i>)	5 - 20	ND
<i>Chrysanthemum</i> ¹	7.5 - 25	ND
<i>Exacum</i>	25 - 50	0.01 - 0.03
<i>Geranium</i>	10 - 25	ND
New Guinea Impatiens	5 - 10	ND
Poinsettia	2.5 - 80 (see Table 6)	0.03 - 0.5
Sunflower	30 - 50	1 - 2

[†] These rate ranges were determined largely under mid-Atlantic conditions using medium-vigor cultivars. Rates should be adjusted to reflect the need for higher rates for vigorous cultivars and in the Sunbelt Region and lower rates in the Northern Belt Region.

¹ *Chrysanthemum* spray: A rate of 7.5 - 15 ppm (a.i.) is recommended for sensitive varieties and 15 - 25 ppm (a.i.) for others. Spray when the axillary shoots following the pinch are 1.5 inches long or before rapid elongation occurs. If a second application is required, it should be made two weeks after the first.

ND = Rates for this application technique have not been determined.

TABLE 6: POINSETTIA GROWTH VIGOR GROUPS AND SPRAY RATE RANGE TRIAL GUIDELINES[†]

Vigor Groups	Example Cultivars	Weeks of growth after pinch (6.5-inch pot)	Weeks of growth after pinch (6.5-inch pot)	Topflor Spray Rate (ppm a.i.)
		NORTH	SOUTH	
Very Low	'Hollypoint', 'Carousel'	7 - 8	4 - 5	NR
Low	'Maren', 'Sonora Red'	5 - 6	3 - 5	2.5 - 20
Moderate	'Freedom Red', 'Orion'	3 - 4	2 - 3	15 - 40
Substantial	'Red Violet', 'Spotlight Dark Red'	2 - 3	2 - 3	30 - 60
Extreme	'Monet Twilight', 'Snowcap'	2 - 3	1 - 2	60 - 80

[†] These are guidelines to provide relative growth differences among cultivars. They should not be viewed as specific recommendations.

NR = Not a recommended use.

Poinsettia Application Timing

Early Applications: Treat plants at pinch up to 6 weeks after pinch, or 8 to 12 weeks before finishing.

Late Applications: The timing of application should be based upon the height of the poinsettia in relation to height goal. If final plant height goal is 15 inches, then apply Topflor when the plants are 12 - 13 inches in height. To ensure uniformity, any plants

shorter than 12 - 13 inches should not be treated at that time. Like most PGRs, seasonably late applications of Topflor will reduce plant height, with minimal to no effect on bract size.

Perennial Plants, Herbaceous

Topflor is effective in controlling height of a wide variety of herbaceous perennial plants. Rate ranges for different species and cultivars vary greatly. Trials should be conducted using a general guideline rate of 20 - 80 ppm (a.i.) for spray applications. Examples of perennials for which the product has provided optimum height control include:

<i>Acalypha</i>	<i>Arbuton</i>	Butterfly bush (<i>Buddleia</i> spp.)
<i>Coreopsis</i>	<i>Fuchsia</i>	<i>Lantana</i>
<i>Lobelia</i>	<i>Pachystachys</i>	<i>Phlox</i>
<i>Scabiosa</i>	Sage, Mexican bush (<i>Salvia leucantha</i>)	Sage, Russian (<i>Perovskia atriplicifolia</i>)
<i>Verbena</i>	<i>Veronica</i>	

Woody Landscape Plants (Container-grown in greenhouses and shadehouses)

Topflor is effective in controlling the height on a wide variety of woody landscape plants using either spray or drench applications. Rate ranges for different species vary greatly. Trials should be conducted using a general guideline rate of 100 - 200 ppm (a.i.) for spray applications. Typical application rate is 1 gallon of spray mixture (up to 200 ppm) per 200 square feet of potted plants. Examples of woody nursery plants for which the product has provided optimum height control include:

<i>Abelia</i>	Glory bush (<i>Tibouchina</i> spp.)
<i>Azalea</i>	Holly: <i>Ilex</i> X <i>attenuata</i> 'Fosteri' (Foster holly)
<i>Bougainvillea</i>	<i>Ilex</i> X <i>meserveae</i> 'China Girl'
<i>Cotoneaster dammeri</i> 'Coral Beauty'	Honeysuckle 'Goldflame' (<i>Lonicera</i> X <i>heckrottii</i>)
Crape myrtle 'Natchez'	<i>Hydrangea</i>
<i>Dipladenia</i>	<i>Photinia</i> X <i>fraseri</i> (Fraser photinia)
<i>Euonymus kiautschovicus</i> 'Manhattan'	<i>Rhododendron catawbiense</i> 'Nova Zembla'
<i>Gardenia jasminoides</i> 'Mystery'	Rose

TERMS AND CONDITIONS OF USE

If terms of the following *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies*.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer*, *Inherent Risks of Use* and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or this *Limitation of Remedies* in any manner.

© Copyright ____ by SePRO Corporation

(Base label for all containers):



Topflor[®]

Ornamental Plant Growth Regulator

A growth regulator for use on ornamental plants grown in containers in commercial nurseries, greenhouses and shadehouses.

IN NEW YORK STATE, TOPFLOR CAN BE USED FOR GREENHOUSE APPLICATIONS ONLY.

Active Ingredient:

flurprimidol: α -(1-methylethyl)- α -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol..... 0.38%

Other Ingredients 99.62%

Total 100.00%

Contains 15 grams of active ingredient per gallon.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

CAUTION. Causes moderate eye and skin irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet.

Refer to inside of label booklet for additional precautionary information and directions for use, including first aid and storage and disposal.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* inside label booklet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

19/19

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling

Nonrefillable Container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

See attached booklet for complete container disposal directions including triple rinsing and pressure rinsing instructions.

Shake well before using.

EPA Reg. No. 67690-20
FPL20110912

EPA Est. No. _____
SPC - _____

SePRO Corporation 11550 N. Meridian Street, Suite 600 Carmel, IN 46032, U.S.A.

Ornamental Plant Growth Regulator

Net Contents _____
[Refillable or Nonrefillable container _____]