

67690-24

07/05/2011

1/14

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Antimicrobials Division (7510P)
1200 Pennsylvania Avenue NW
Washington, D.C. 20460

EPA Reg. Number: 67690-24

Date of Issuance: JUL - 5 2011

Term of Issuance: Unconditional

Name of Pesticide Product: K-TEA ALGAECIDE

NOTICE OF PESTICIDE:
Registration
[X] Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Ms. Amy Dugger-Ronyak
SEPRO Corp.
11550 N. Meridian St. Suite 600
Carmel, IN 46032

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Antimicrobials Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Copper RED, the EPA has reregistered this product subject to the comments recorded in the succeeding paragraph. This action is taken under the authority of section 4(g) (2) of the Federal Insecticide, Fungicide and Rodenticide Acts, as amended. Registration under this section does not eliminate the need for continual reassessment of pesticides. The EPA may require submission of data at any time to maintain the registration of the subject product.

The following label revisions must be made:

- 1. Delete the following text from page 2 in the "Precautionary Statements" section:

Keep Out of Reach of Children
Si usted no entiende Detail.)
Concentrated Formulation

Signature of Approving Official: Jacqueline Campbell McFarlane
Product Manager, Team 34

Date: JUL - 5 2011

CONCURRENCES

Table with columns for SYMBOL, SURNAME, and DATE, and rows for Regulatory Management Branch II and Antimicrobials Division (7510P).

EPA Form 8570

- 2. Revise the "Precautionary Labeling" on page 2 to agree with the product's acute toxicity profile by stating:

CAUTION. Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

- 3. The statement, "Potable water sources ... treatments" on page 2 must be placed after the "User Safety Instructions" box and updated to be in agreement with the Copper RED by stating:

"For applications in waters destined for use as drinking water, those waters must receive additional and separate potable water treatment. Do not apply more than 1.0 ppm as metallic copper in these waters."

- 4. The "User Safety Recommendations" heading must be brought into compliance with PR Notice 2000-6, Mandatory Labeling, by stating "User Safety *Instructions*." Also, revise the phrase, "User should" to read "Users *must*."
- 5. The Environmental Hazards text currently on the label must be revised to read:

This pesticide is toxic to fish and aquatic invertebrates. Waters treated with this product may be hazardous to aquatic organisms. Treatment of aquatic weeds and algae can result in oxygen loss from decomposition of dead algae and weeds. This oxygen loss can cause fish and invertebrate suffocation. To minimize this hazard, do not treat more than 1/2 of the water body to avoid depletion of oxygen due to decaying vegetation. Wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult with the State or local agency with primary responsibility for regulating pesticides before applying to public waters, to determine if a permit is required.

Certain water conditions including low pH (5.6-5.8), low dissolved organic carbon (DOC) levels (3.0 mg/L or lower), and "soft" waters (i.e., alkalinity less than 50 mg/L), increases the potential acute toxicity to non-target aquatic organisms."

- 6. Revise the first statement in the 1st paragraph under the "Directions for Use" on page 3 by deleting the phrase, "mixing area" and stating "*application area*."
- 7. Revise the "General Precautions and Restrictions" section by adding another bullet with the text, "No more than 1/2 of the water body may be treated at one time."
- 8. The following comments are changes that must be made under the "Directions for Use" to bring them in compliance with PR Notice 2000-6, Mandatory labeling:
 - a. Revise the third statement in the third paragraph under "General Information" on page 4 by deleting the term, recommended.

- b. Revise the 1st statement under “Avoiding Injurious Spray Drift” on page 4 by deleting the term, should, and stating “*must.*”
 - c. Revise the 2nd sentence in first paragraph on page 5 by deleting the phrase, “it is recommended” to read “*use* the highest rate in the rate range *to* be applied.”
 - d. Revise the 1st statement under “For best results” on page 5 by deleting the term, should, and stating “*must.*”
 - e. Revise the 5th statement under “Application methods” on page 6 by deleting the phrase, “it is recommended to.”
 - f. Revise the 2nd statement under “Surface Application” on page 6 by deleting the term, recommended.
 - g. Revise the 2nd statement under “Subsurface Application on page 6 by deleting the phrase, “it is recommended to,” and the term, recommended.
 - h. Revise the 1st and 2nd statements under “Aerial Application on page 6 by deleting the term, recommended.
 - i. Revise the 1st statement under “Drip System Application” on page 7 by deleting the term, should, and stating “*must.*”
 - j. Revise the 2nd statement under “Determining Amount of K-TEA on page 7 by deleting the term, should, and stating “*must.*”
 - k. Revise the last statement in 2nd paragraph under “Submersed Plant Control” on page 8 by deleting the term, should, and stating “*must.*”
 - l. Revise the 3rd statement under “Hydrilla Control” on page 8 by deleting the term, should, and stating “*must.*”
9. Since this product is applied by aerial application, the label must support the following “Spray Drift Management” test as per the Copper RED:

Spray Drift Management

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet size

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on target desposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exists, or b) stable atmosphere conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers and surrogates.

Requirements for Aerial Applications:


- The boom length must not exceed 75% of wingspan or 90% of the rotor blade
- Release spray at the lowest height with efficacy and flight safety. Do not release spray at height greater than 10 ft above the water surface unless a greater height is required for aircraft safety.
- When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

A stamped label with conditions is enclosed for your records. For additional labeling clarity and consistency, please refer to the enclosed EPA reviews. Submit three (3) copies of your final printed labeling before distributing or selling the product bearing the revised labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Should you have any questions regarding this letter, please contact me at (703) 308-6416 or by e-mail at campbell-mcfarlane.jacqueline@epa.gov or Killian Swift by telephone at (703) 308-6346 or by e-mail at swift.killian@epa.gov. When submitting information or data in response to this letter, a copy of this letter should accompany the submission to facilitate processing.

Sincerely,



Jacqueline Campbell-McFarlane
 Product Manager, Team 34
 Regulatory Management Branch II
 Antimicrobials Division (7510P)

[Front of label booklet, ALL containers]



K-Tea*

AQUATIC ALGAECIDE

FOR USE IN SLOW-MOVING OR QUIESCENT BODIES OF WATER INCLUDING: GOLF COURSE, ORNAMENTAL, FISH, IRRIGATION AND FIRE PONDS; FRESH WATER LAKES AND FISH HATCHERIES; POTABLE WATER RESERVOIRS, RIVERS, STREAMS, BAYS AND COVES; AND CROP AND NON-CROP IRRIGATION CONVEYANCE SYSTEMS (CANALS, LATERALS AND DITCHES).

There are no water use restrictions following application of K-Tea.

Active Ingredient

Copper hydroxide† (CAS# 20427-59-2)	12.9%
Other Ingredients	87.1%
TOTAL	100.00%

†Metallic Copper equivalent = 8%

ACCEPTED
with COMMENTS
EPA Letter Dated:

JUL - 5 2011

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

67690-24

KEEP OUT OF THE REACH OF CHILDREN
CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

CONCENTRATED FORMULATION

Refer to inside of label booklet for additional precautionary information and directions for use including storage and disposal.

NOTICE: Read the entire label before using. Use only according to label directions. Before buying or using K-Tea, read *Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* inside label booklet.

For additional information on our products, please visit www.sepro.com.

EPA Reg. No. 67690-24
FPL022509

EPA Est. No. _____
SPC-_____

SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032, U.S.A.

Algaecide

Net contents _____

* Trademark of SePRO Corporation

[Label booklet text, ALL containers]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF THE REACH OF CHILDREN CAUTION / PRECAUCIÓN

(If you do not understand the label, find someone to explain it to you in detail.)

Harmful if inhaled. Causes eye irritation. Harmful if swallowed or absorbed through the skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to K-Tea are barrier laminate, butyl rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils. If you want more options, follow the instructions for categories A on an EPA chemical-resistant category selection chart.

Mixers, loaders, applicators, and other handlers must wear the following:

- Long-sleeve shirt and long pants;
- Shoes plus socks;
- Chemical-resistant gloves (such as nitrile or butyl rubber); and
- Respirator with an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter.

Exception: Aquatic Subsurface Application or Closed Application System

After K-Tea has been diluted or tank mixed with water, users must, at a minimum, wear (NOTE - Mixers and loaders for this application method must still wear the PPE as described in the above section):

- Short-sleeved shirt and long pants;
- Shoes plus socks; and
- Respirator with an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the K-Tea's concentrate. Do not reuse them.

USER SAFETY

Users should:

- Wash the outside of gloves before removing.
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling K-Tea. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

If the following treatment and application guidance is not followed, waters treated with K-Tea may be hazardous to aquatic organisms. Treatment of aquatic weeds and algae can result in oxygen loss from decomposition of dead algae and weeds. Under certain conditions, this oxygen loss can cause fish and invertebrate suffocation. To minimize this potential hazard, in quiescent waters, do not treat more than one-half (1/2) of the water body to avoid depletion of oxygen due to decaying vegetation ~~when susceptible algae mats cover more than one-half (1/2) the water body, when susceptible macrophytic algae or susceptible vascular plants are mature or have grown to the water's surface and cover more than one-half (1/2) of the water body, or when susceptible phytoplanktonic algae blooms are dense and infest greater than one-half (1/2) the water body.~~ Wait at least 14 days between treatments (except for in aquaculture). Consult with the State or local agency with primary responsibility for pesticide regulation before applying to public waters, to determine if a permit or public notification is required.

Certain water conditions including low pH (≤ 6.5), low dissolved organic carbon (DOC) levels (3.0 mg/L or lower), and "soft" waters (i.e. alkalinity less than 50 mg/L), increases the potential acute toxicity to non-target aquatic organisms. Do not use in waters containing trout or other fish species that are highly sensitive to copper if the alkalinity is less than 50 ppm. Fish toxicity generally decreases when the hardness of water increases. K-Tea should not be used in ornamental ponds containing Koi.

DIRECTIONS FOR USE

It is a violation of Federal law to use K-Tea in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply in a way that concentrated K-Tea will contact workers or other persons, either directly or through drift, only protected handlers may be in close proximity to the mixing area or application equipment while in use. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

GENERAL INFORMATION

K-Tea algaecide provides effective control of various green and blue-green (cyanobacteria) algae, including filamentous, planktonic and branched algae which can occur in slow-moving or quiescent bodies of water, including golf course, ornamental, fish, irrigation and fire ponds; fresh water lakes and fish hatcheries; potable water reservoirs and associated waters (rivers, streams, bays and coves); and crop and non-crop irrigation conveyance systems (canals, laterals and ditches). K-Tea is most effective when applied at the first signs of algal bloom.

K-Tea may be applied aerially, through a ground sprayer or spray boat as a surface spray, and as a subsurface application through weighted hoses. K-Tea may also be applied in an invert emulsion or mixed with a polymer (except for CA).

In static or quiescent water such as lakes, reservoirs, ponds, and static canals, the application site is defined by this label as the specific location where K-Tea is applied. In slow-moving and flowing waters such as canals and rivers, the application site is defined by this label as the target location for plant control. Use the lower ~~concentration~~ rate in soft water (less than 50 ppm alkalinity), for light infestations and less mature plants; use the higher concentration in hard water (above 50 ppm alkalinity), for dense infestations and when targeting more mature vegetation.

General Precautions and Restrictions

- Do not enter or allow others to enter until application is complete.
- Do not apply undiluted solution of K-Tea directly to, or otherwise permit it to come into contact with any desirable plants as injury may result. Do not apply in such a way that concentrated K-Tea comes in contact with crops, ornamentals, grass or other desirable plants. Wash spray equipment thoroughly before and after each application.

Avoiding Injurious Spray Drift

Applications ~~should~~ be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward desirable susceptible crops or ornamental plants near enough to be injured. Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

APPLICATION INFORMATION

For aquatic weed control (including vascular plants and algae), do not exceed 1.0 ppm copper during any single application; wait a minimum of 14 days between re-treatments (Exception - In aquaculture do not exceed 0.4 ppm during any single application when targeting nuisance algae; there are no restrictions on retreatment intervals).

Free floating (planktonic) algae, such as *Anabaena*, *Aphanizomenon*, *Chlorella*, *Dictyosphaerium*, *Euglena*, and *Microcystis*, are controlled using 0.2 to 1.0 ppm metallic copper depending upon severity of growth.

Mat-forming (filamentous) algae, such as *Cladophora*, *Oedogonium Lyngbya*, and *Oscillatoria*, may require up to 0.5 to 1.0 ppm metallic copper depending on growth, intensity, and other

certain other conditions. *Chara* and *Phormidium* are difficult to control unless treatment at 0.5 to 1.0 ppm metallic copper is generally initiated at the first signs of algal bloom. For dense infestations of filamentous algae or where the species of *Hydrodictyon*, *Cladophora* or *Pithophora* are present, it is ~~at the~~ the highest rate in the rate range be applied. An adjuvant, such as d-limonene or similar surfactant, may be added for enhanced control of floating mats or difficult to control species of algae. Follow adjuvant product labeling instructions for application rates and use directions.

For best results, K-Tea should be applied when conditions are calm and when algae first appear. Even distribution of K-Tea in the water will improve algae control; therefore, apply in a manner that distributes K-Tea throughout the treated area. Filamentous algae species are easier to control before floating to the water's surface (when they are forming on the pond/lake bottom). The genera of algae listed in Table 1 are commonly found in waters of the United States.

Organism	0.2-1.0 PPM Copper		0.5-1.0 PPM Copper	
Cyanophyceae (Blue-green Algae)	<i>Anabaena</i> <i>Aphanizomenon</i> <i>Cylindrospermum</i> <i>Gloeotrichia</i> <i>Gomphosphaeria</i>	<i>Microcystis</i> <i>Oscillatoria</i> <i>Plectonema</i> <i>Polycystis</i>	<i>Calothrix</i> <i>Nostoc</i> <i>Phormidium</i> <i>Symploca</i>	
Chlorophyceae (Green Algae)	<i>Botryococcus</i> <i>Closterium</i> <i>Coelastrum</i> <i>Draparnaldia</i> <i>Enteromorpha</i> <i>Gloeocystis</i> <i>Microspora</i>	<i>Spirogyra</i> <i>Tribonema</i> <i>Ulothrix</i> <i>Zygnema</i>	<i>Ankistrodesmus</i> <i>Chara</i> <i>Chlorella</i> <i>Cladophora</i> <i>Crucigenia</i> <i>Desmidium</i> <i>Golenkinia</i>	<i>Nitella</i> <i>Oocystis</i> <i>Palmella</i> <i>Pithophora</i> <i>Scenedesmus</i> <i>Staurastrum</i> <i>Tetraedron</i>
Diatomaceae (Diatoms)	<i>Asterionella</i> <i>Fragilaria</i> <i>Gomphonema</i> <i>Melosira</i> <i>Navicula</i>	<i>Nitzschia</i> <i>Stephanodiscus</i> <i>Synedra</i> <i>Tabellaria</i>	<i>Achnanthes</i> <i>Cymbella</i> <i>Neidium</i>	
Protozoa (Flagellates)	<i>Ceramium</i> <i>Cryptomonas</i> <i>Dinobryon</i> <i>Euglena</i> <i>Glenodinium</i>	<i>Mallomonas</i> <i>Synura</i> <i>Uroglena</i> <i>Volvox</i>	<i>Chlamydomonas</i> <i>Eudorina</i> <i>Haematococcus</i> <i>Pandorina</i> <i>Peridinium</i>	

Use Table 2 below to determine the amount of K-Tea required to achieve the desired copper concentration. For optimal algae control, maintain the desired copper concentration for a minimum of three (3) to twenty-four (24) hours. Rates given below represent concentrations for quiescent or slow-moving water. If water flow results in significant dilution of the treated water, it may be necessary to meter K-Tea into the water (refer to Table 3 and the Drip System Application section of this label).

TABLE 2			
APPLICATION RATES FOR QUIESCENT OR SLOW-MOVING WATER			
Average Water Depth of Treatment Site (Feet) [†]	Gallons of K-Tea per Surface Acre to Achieve the Desired Copper Concentration		
	0.2 ppm	0.5 ppm	1.0 ppm
1	0.7	1.7	3.4
2	1.4	3.4	6.8
3	2.0	5.1	10.2

[†] To calculate the application rate in water with an average depth >3 feet, multiply the average depth by the recommended amount of K-Tea in one foot of water.

Application Methods

Apply by handgun, spray boat, helicopter or other method of application which provides uniform coverage of the treated area. K-Tea may be applied diluted or undiluted, whichever is most suitable to ensure uniform coverage of the area to be treated. Dilution with water may be necessary at the lower application rates. Dilute the required amount of K-Tea with enough water to ensure even distribution in the treated area with the type of equipment being used. For best results, ~~do not~~ dilute K-Tea in water to provide a minimum spray mix of 20 to 50 gallons per acre. In areas with heavy infestations of filamentous algae, a total tank mix of 100 gallons per acre may be necessary. To ensure best results, remove large mats of floating algae manually before treatment or break them up before spraying or while application is being made. A second application may be necessary for heavily infested areas. When treating moving water, apply the spray solution counter to the flow of water (unless metering product into flowing water – see recommendations under *Drp System Application* section of this label).

Surface Application

Spray diluted mixture from shore or boat evenly across the surface of the water. Surface applications generally are ~~recommended~~ near shorelines, in shallower waters, and/or for floating algal mats, and may be made from shore into shallow water.

Subsurface Application

In deeper water and/or when filamentous algae is forming on the bottom, ~~it is recommended to~~ make a subsurface application of K-Tea at ~~recommended~~ rates through weighted trailing hoses where the greatest concentration of algae is present. Do not drag hoses on the bottom.

Adjuvants/Surfactants

Adjuvants (polymers, invert emulsions, etc.) or surfactants may be added to K-Tea or to a K-Tea/water premix to improve efficacy. Consult the manufacturer's recommendations regarding the use of these products for improved algae control.

Aerial Application

Apply the ~~recommended~~ rate of K-Tea in a ~~recommended~~ minimum of 10 gallons of total spray solution per surface acre. Add the ~~recommended~~ rates of a drift control or sinking agent to the spray solution. Maintain constant agitation during addition of the polymer (except

* Trademark of SePRO Corporation

California – polymers not approved for use with K-Tea) and continue throughout the application.

Drip System Application

For Use in Irrigation Conveyance Systems and Other Moving Water

For best results, application should be made in anticipation of algae that may interfere with normal flow or delivery of water (obstruction of lateral headgates, screens, pumps, pumping systems and siphon tubes). Delayed treatment may result in matting or compaction of algae mats.

Determine the water flow rate prior to treatment of the water system. If available, use weirs, orifices or similar devices which give accurate water flow measurements. If these devices are not available, volume of flow may be estimated by the following formula:

$$\text{Average width (feet)} \times \text{Average depth (feet)} \times \text{Average velocity (feet/second)} \times 0.9 \\ = \text{Cubic feet per second (cfs)}$$

To determine velocity, measure the time it takes a floating object in the middle of the canal to travel a given distance. Divide the distance traveled (feet) by the time (seconds) for velocity (feet/second). Repeat this procedure at least three times and then calculate the average velocity. Use the average velocity (feet/second) in the formula above to determine the flow rate (cfs).

Once the water flow rate (cfs. or gallons per minute) has been calculated, find the corresponding drip rate for K-Tea in Table 3.

TABLE 3				
APPLICATION RATES FOR MOVING WATER				
WATER FLOW RATE		K-TEA DRIP RATE		
C.F.S.	GAL./MIN.	QTS./HR.	ML./MIN.	FL. OZ./MIN.
1	500	1.25	20	0.7
2	1,000	2.50	40	1.3
3	1,500	3.75	60	2.0
4	2,000	5.00	80	2.3
5	2,500	6.25	100	3.3

Determining Amount of K-TEA

The rates shown above will produce a concentration of 1.0 ppm Cu⁺⁺ in the treated water. This concentration should be maintained for at least 3 hours. Lower concentrations may be used on highly susceptible algae species or if longer exposure times are maintained. To determine the total amount of K-Tea needed to maintain the drip rate for 3 hours, calculate as follows:

$$\begin{aligned} &\text{Quarts / Hour.} \times 3; \text{ or} \\ &\text{Milliliters / Minute} \times 180; \text{ or} \\ &\text{Fluid ounces / Minute} \times 180. \end{aligned}$$

For longer injection periods, multiply dosage rate by desired time in minutes or hours as

appropriate. Thorough mixing is necessary to uniformly disperse K-Tea in the water; therefore, apply K-Tea in the channel at weirs or other structures which create turbulence or at several injection points across the flow. K-Tea may be injected as concentrated K-Tea or diluted with water.

Calibrating For Drip Application-

Pour the amount of K-Tea needed to treat for three hours (calculated above) into a drum or tank equipped with valve or other volume control device that can be calibrated to maintain a constant drip rate. Open the valve and allow K-Tea to drip into a graduated container (measuring cup, graduated cylinder, etc.), using a stopwatch to measure the time required to reach the desired volume. Adjust the valve so that K-Tea is dripping at the desired rate.

NOTE: If the flow rate changes during the treatment period, it may be necessary to recalibrate the delivery system. A small pump or other metering pump may be used to meter K-Tea into the water more accurately.

Distance of algae control from the application point will vary with severity of infestation, water quality and conditions of water conveyance system. Repeat application at a designated point downstream from the previous treatment station. Repeat as necessary to treat entire infested area. It may be necessary to periodically repeat treatments to maintain seasonal control. Consult an Aquatic Specialist to determine optimal use rate, location of treatment stations and duration of treatment period under local conditions.

Submersed Plant Control

K-Tea may be tank mixed with other aquatic herbicides to enhance efficacy, and to control algae in areas where heavy algae growth that may cover target submersed plant species and interfere with herbicide uptake. K-Tea may be applied to control hydrilla (*Hydrilla verticillata*), egeria (*Egeria densa*), and other copper sensitive aquatic weeds. Apply K-Tea at a recommended rate to achieve 0.75 to 1.0 ppm copper. In heavily infested areas, a second application may be necessary.

Do not mix concentrates in tank without first adding water. To ensure compatibility, a jar test is recommended before field application. K-Tea cannot be mixed with any product containing a label prohibition against such mixing and must be used in accordance with the more restrictive of the label limitations and precautions. No label dosage rates should be exceeded.

Hydrilla Control – K-Tea + Diquat Tank Mix

K-Tea can be mixed with diquat (diquat dibromide) at 4 gallons K-Tea and 2 gallons diquat (Littora*, Reward® - 2 lbs a.i./gallon) per acre. Lower rates of K-Tea may also enhance the activity of diquat. K-Tea should be applied at a minimum of 0.1 ppm in combination with diquat. Higher rates may be needed in areas with dense weeds.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in cool dry place.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Nonrefillable Container Disposal (rigid, 5 gallons or less): Do not reuse or refill this

* Trademark of SePRO Corporation

container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container Disposal (rigid, larger than 5 gal): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable Container Disposal (rigid, larger than 5 gal): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment, rinsate collection system, or treatment area. Repeat this rinsing procedure two more times. If returning container, seal all openings which have been opened during use. Return the empty container to a collection site designated by SePRO Corporation. If the container has been damaged and cannot be returned according to the recommended procedures, contact SePRO Corporation at 1-800-419-7779 to obtain proper handling instructions.

WARRANTY DISCLAIMER

SePRO Corporation warrants that K-Tea conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of K-Tea. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of K-Tea contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from K-Tea (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for K-Tea bought, or
- (2) Replacement of amount of K-Tea used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of K-Tea unless SePRO Corporation is promptly notified of such losses or damages in writing. To the extent consistent with applicable law, in no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* above and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or *Limitations of Remedies* in any manner.

©Copyright ____ SePRO Corporation

Littora is a trademark of SePRO Corporation.
Reward® is a trademark of Syngenta Group Company