

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

February 4, 2022

Lei Han, Ph.D Head of Regulatory Affairs SePRO Corporation 11550 N. Meridian Street Suite 600 Carmel, IN 46032

Subject: Registration Review Label Mitigation for Fluridone

Product Name: FLURIDONE SRP EPA Registration Number: 67690-32

Application Date: 05/17/2019 Decision Number: 581582

Dear Dr. Han:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Fluridone Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2 EPA Reg. No. 67690-32 Decision No. 581582

If you have any questions about this letter, please contact Srijana Shrestha by phone at 202-566-2329, or via email at <a hrestha.srijana@epa.gov.</a>

Sincerely,

Linda Arrington, Branch Chief

Risk Management and Implementation Branch 4

Pesticide Re-Evaluation Division

Office of Pesticide Programs

Enclosure: Stamped Label



FLURIDONE

GROUP

12

**HERBICIDE** 

# **Avast! WDG**

**AQUATIC HERBICIDE** 

An herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, potable water sources, drainage canals, irrigation canals and rivers.

Active Ingredient	
fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone	5.0%
Other Ingredients	95.0%
TOTAL	100.0%
Contains 2 pounds of fluridone per 40-pound container.	

# Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

Refer to label booklet for additional Precautionary Information and Directions for Use including First Aid and Storage and Disposal.

**NOTICE:** Read the entire label before using. Use only according to label directions. **Before** buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet. If terms are unacceptable, return at once, unopened.

EPA Reg. No. 67690-32 FPL20210720	EPA Est. No [P/N]	
SePRO Corporation • 11550 N. Meridian	Street, Suite 600 • Car	mel, IN 46032, U.S.A.
Aquatic Herbicide	Net Weight	(Non-refillable)

ACCEPTED

Feb 04, 2022

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under

EPA Reg. No. 67690-32

#### [Label booklet text]

	FIRST AID			
If in eyes	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.			
	Remove contact lenses, if present, after the first 5 minutes, then continue			
	rinsing eye.			
	Call a poison control center or doctor for treatment advice.			
If on skin or	Take off contaminated clothing.			
clothing	Rinse skin immediately with plenty of water for 15 - 20 minutes.			
	Call a poison control center or doctor for treatment advice.			
If swallowed	Call a poison control center or doctor immediately for treatment advice.			
	Have person sip a glass of water if able to swallow.			
	<ul> <li>Do not induce vomiting unless told to do so by a poison control center or</li> </ul>			
	doctor.			
	Do not give anything by mouth to an unconscious person.			
If inhaled	Move person to fresh air.			
	If person is not breathing, call 911 or an ambulance, then give artificial			
	respiration, preferably mouth-to-mouth if possible.			
	Call a poison control center or doctor for further treatment advice.			
HOTLINE NUMBER				

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if Swallowed, Absorbed Through Skin or Inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### **ENGINEERING CONTROLS (AIRCRAFT)**

Aircraft pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides [40 CFR 170.305].

#### **ENVIRONMENTAL HAZARDS**

Follow use directions carefully so as to minimize adverse effects on non-target organisms. Do not contaminate untreated water when disposing of equipment washwaters. Trees and shrubs growing in water treated with Avast! WDG may occasionally develop chlorosis. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas.

# **Non-Target Organisms Advisory Statement**

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

#### PRODUCT INFORMATION

Avast! WDG is a selective systemic aquatic herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, drainage canals, irrigation canals and rivers. Avast! WDG is a pelleted formulation containing 5 percent fluridone. Avast! WDG is absorbed from water by plant shoots and from hydrosoil by the roots of aquatic vascular plants. It is important to maintain Avast! WDG in contact with the target plants for as long as possible. Rapid water movement or any condition which results in rapid dilution of Avast! WDG in treated water will reduce its effectiveness.

In susceptible plants, Avast! WDG inhibits the formation of carotene. In the absence of carotene, chlorophyll is rapidly degraded by sunlight. Herbicidal symptoms of Avast! WDG appear in seven to ten days and appear as white (chlorotic) or pink growing points. Under optimum conditions, 30 to 90 days are required before the desired level of aquatic weed management is achieved with Avast! WDG. Species susceptibility to Avast! WDG may vary, depending on time of year, stage of growth, and water movement. For best results, apply Avast! WDG prior to initiation of weed growth or when weeds begin active growth. Application to mature target plants may require an application rate at the higher end of the specified rate range and may take longer to control.

Avast! WDG is not corrosive to application equipment.

The label provides recommendations on the use of a chemical analysis for the active ingredient.

SePRO Corporation recommends the use of a High-Performance Liquid Chromatography (HPLC) for the determination of the active ingredient concentration in the water. Contact SePRO Corporation to incorporate this test, known as a FasTEST®, into your treatment program. Other proven chemical analyses for the active ingredient may also be used. The FasTEST is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredient in the water.

#### **FASTEST**

Application rates are provided in pounds of Avast! WDG to achieve a desired concentration of the active ingredient in parts per billion (ppb). The maximum application rate or sum of all application rates is 90 ppb in ponds and 150 ppb in lakes and reservoirs per annual

**growth cycle.** This maximum concentration is the amount of product calculated as the target application rate, NOT determined by testing the residues of the active ingredient in the treated water.

# **Weed Resistance Management**

For resistance management, Avast! WDG is a Group 12 herbicide. Any weed population may contain or develop plants naturally resistant to Avast! WDG and other Group 12 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Avast! WDG or other Group 12 herbicides within a growing season or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where
  information on resistance in target weed species is available, use the less resistanceprone partner at a rate that will control the target weed(s) equally as well as the more
  resistance-prone partner. Consult your local extension service or pest control advisor if
  you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the desirable plants and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of
  resistance development. Indicators of possible herbicide resistance include: (1) failure
  to control a weed species normally controlled by the herbicide at the dose applied,
  especially if control is achieved on adjacent weeds; (2) a spreading patch of noncontrolled plants of a particular weed species; (3) surviving plants mixed with controlled
  individuals of the same species. If resistance is suspected, prevent weed seed
  production in the affected area by an alternative herbicide from a different group or by a
  mechanical method. Prevent movement of resistant weed seeds to other areas by
  cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of plants and weed biotypes.

#### **Use Restrictions**

- **Obtain required permits:** Permits may be required by state or local agencies. Consult with appropriate State or local water authorities before applying this product.
- **New York State:** Application of Avast! WDG is not permitted in waters less than two (w) feet deep.
- **Hydroponic Farming:** Do not use water treated with Avast! WDG for hydroponic farming.
- **Greenhouse and Nursery Plants:** Do not use water treated with Avast! WDG for irrigating greenhouse or nursery plants unless use of an approved assay confirms that residues are

less than 1 ppb.

Water Use Restrictions Following Applications of Avast! WDG (Days)

Application Rate	Drinking <sup>†</sup>	Fishing	Swimming	Livestock/Pet Consumption	Irrigation <sup>††</sup>
Maximum Rate (150 ppb) or less	0	0	0	0	See irrigation instructions below

Note below, under *Potable Water Intakes*, the information for application of Avast! WDG within ¼ mile (1,320 feet) of a functional potable water intake.

- Potable Water Intakes: Concentrations of the active ingredient, fluridone, up to 150 ppb are allowed in potable water sources; however, in lakes and reservoirs or other sources of potable water, DO NOT apply Avast! WDG at application rates greater than 20 ppb within one-fourth (1/4) mile (1,320 feet) of any functioning potable water intake. At application rates of 8 to 20 ppb, Avast! WDG may be applied where functioning potable water intakes are present. NOTE: Existing potable water intakes that are no longer in use, such as those that have been replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water intakes.
- Aircraft pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides [40 CFR 170.305].

#### **Use Precautions**

• Irrigation: Irrigation with water treated with Avast! WDG may result in injury to the irrigated vegetation. Follow these precautions and inform those who irrigate from areas treated with Avast! WDG of the irrigation time frames or water assay requirements presented in the table below. Follow the following time frames and assay directions to reduce the potential for injury to vegetation irrigated with water treated with Avast! WDG. There is a greater potential for crop injury when water treated with Avast! WDG is applied to crops grown in low organic and sandy soils.

WAITING PERIODS BEFORE IRRIGATING WITH WATER TREATED WITH AVAST! WDG			
		Days Af	ter Application
Application Site	Tree Crops   Row Crops/   Areas to be Planted Including		Newly Seeded Crops/Seedbeds or Areas to be Planted Including Overseeded Golf Course Greens
Ponds and Static Canals †	7	30	Assay required
Canals	7	7	Assay required
Rivers	7	7	Assay required
Lakes and Reservoirs <sup>††</sup>	7	7	Assay required

For purposes of Avast! WDG labeling, a pond is defined as a body of water 10 acres or less

Note below, under *Irrigation*, specific time frames or fluridone residues that provide the widest margin of safety for irrigating with water treated with Avast! WDG.

in size. A lake or reservoir is greater than 10 acres.

<sup>††</sup> In lakes and reservoirs where one-half or greater of the body of water is treated, use the pond and static canal irrigation precautions.

Where the use of Avast! WDG treated water is desired for irrigating crops prior to the time frames established above, use aFASTESTFasTEST or other appropriate means of analysis is recommended to measure the concentration in the treated water. Where FASTESTFasTEST has determined that concentrations are less than 10 parts per billion (ppb), there are no irrigation precautions for irrigating established tree crops, established row crops or turf. For tobacco, tomatoes, peppers or other plants within the Solanaceae family and for newly seeded crops or newly seeded grasses, such as overseeded golf course greens, do not use Avast! WDG treated water if concentrations are greater than 5 parts per billion. Furthermore, when rotating crops, do not plant members of the Solanaceae family in land that has been previously irrigated with fluridone concentrations in excess of 5 ppb. It is recommended that an aquatic specialist be consulted prior to commencing irrigation of these sites.

# WEED CONTROL INFORMATION

Avast! WDG selectivity is dependent upon dosage, time of year, stage of growth, method of application and water movement. The following categories, Controlled, Partially Controlled and Not Controlled, are provided to describe expected efficacy under ideal treatment conditions, using higher to maximum application rates. Use of lower rates will increase selectivity of some species listed as Controlled or Partially Controlled. Additional aquatic plants may be controlled, partially controlled or tolerant to Avast! WDG. Consult an aquatic specialist prior to application to determine a plant's susceptibility to Avast! WDG.

# Vascular Aquatic Plants Controlled by Avast! WDG

#### Submersed Plants

Bladderwort (*Utricularia* spp.)

Coontail, Common (Ceratophyllum demersum)†

Elodea, Common (Elodea canadensis)†

Egeria: Brazilian Elodea (Egeria densa)

Fanwort; Cabomba (Cabomba caroliniana)

Hydrilla (*Hydrilla verticillata*)

Naiad (Najas spp.)†

Pondweed (*Potamogeton* spp.), except Illinois Pondweed<sup>†</sup>

Watermilfoil (Myriophyllum spp.), except Variable-Leaf Milfoil

#### Shoreline Grasses

Paragrass (*Urochloa mutica*)

<sup>†</sup> Native plants that are often tolerant to fluridone at lower use rates. Please consult an aquatic specialist for recommended Avast! WDG use rates (not to exceed maximum labeled rates) when selective control of exotic species is desired.

# Vascular Aquatic Plants Partially Controlled by Avast! WDG

# Floating Plants

Salvinia (Salvinia spp.)

#### Emersed Plants

Alligatorweed (*Alternanthera philoxeroides*)

Cattail (*Typha* spp.)

Lotus, American (Nelumbo lutea)

Parrotfeather (*Myriophyllum aquaticum*)

Smartweed (*Polygonum* spp.)

Spatterdock (Nuphar luteum)

Spikerush (*Eleocharis* spp.)

Waterlily (*Nymphaea* spp.)

Waterprimrose, Creeping (Ludwigia peploides)

Waterpurslane (*Ludwigia palustris*)

Watershield (Brasenia schreberi)

#### Submersed Plants

Limnophila (*Limnophila sessiliflora*)

Pondweed, Illinois (*Potamogeton illinoensis*)

Tapegrass; American Eelgrass (Vallisneria americana)

Watermilfoil, Variable-Leaf (*Myriophyllum heterophyllum*)

#### Shoreline Grasses

Barnyardgrass (Echinochloa crusgalli)

Canarygrass, Reed (*Philaris arundinaceae*)

Cutgrass, Giant (*Zizaniopsis miliacea*)

Torpedograss (Panicum repens)

Watergrass, Southern (*Hydrochloa caroliniensis*)

# Vascular Aquatic Plants Not Controlled by Avast! WDG

#### Floating Plants

Waterhyacinth, Floating (*Eichornia crassipes*)

Water Lettuce (*Pistia stratiotes*)

#### Emersed Plants

Arrowhead (Sagittaria spp.)

Bacopa (Bacopa spp.)

Big Floatingheart; Banana Lily (Nymphoides aquatica)

Bulrush (Scirpus spp.)

Frogbit, American (*Limnobium spongia*)

Pickerelweed; Lanceleaf (*Pontederia* spp.)

Rush (Juncus spp.)

Water Pennywort (Hydrocotyle spp.)

#### Shoreline Grasses

Maidencane (*Panicum hemitomon*)

NOTE: Algae (Chara, Nitella and filimentous species) are not controlled by Avast! WDG.

#### APPLICATION DIRECTIONS

The aquatic plants present in the treatment site should be identified prior to application to determine their susceptibility to Avast! WDG. It is also important to determine the area (acres) to be treated and the average depth in order to select the proper application rate. Do not exceed the maximum labeled rate for a given treatment site per annual growth cycle.

# **Application to Ponds**

Avast! WDG may be applied to the entire surface area of a pond. For single applications, rates may be selected to provide 45 to 90 ppb in the treated water, although actual concentrations in treated water may be substantially lower at any point in time due to the slow-release formulation of this product. When treating for optimum selective control, lower rates may be applied for sensitive target species. Use the higher rate within the rate range where there is a dense weed mass, when treating more difficult to control species, and for ponds less than 5 acres in size with an average depth of less than 4 feet. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the section of this label entitled *Application Rate Calculation - Ponds, Lakes and Reservoirs*. Split or multiple applications may be used where dilution of treated water is anticipated; however, the sum of all applications should total 45 to 90 ppb and must not exceed a total of 90 ppb per annual growth cycle.

Average Water Depth		vast! WDG per urface Acre
of Treatment Site (feet)	45 ppb	90 ppb
1	2.5	5.0
2	5.0	10.0
3	7.5	15.0
4	10.0	20.0
5	12.5	25.0
6	15.0	30.0
7	17.0	34.0
8	19.5	39.0
9	22.0	44.0
10	24.5	49.0

#### **Application to Lakes and Reservoirs**

The following treatments may be used for treating both whole lakes or reservoirs and partial areas of lakes or reservoirs (bays, etc.). For best results in treating partial lakes and reservoirs. Avast! WDG treatment areas should be a minimum of 5 acres in size. Treatment of

areas smaller than 5 acres or treatment of narrow strips, such as boat lanes or shorelines, may not produce satisfactory results due to dilution by untreated water. Rate ranges are provided as a guide to include a wide range of environmental factors, such as target species, plant susceptibility, selectivity and other aquatic plant management objectives. Application rates and methods should be selected to meet the specific lake/reservoir aquatic plant management goals.

# Whole Lake or Reservoir Treatments (Limited or No Water Discharge)

# Single Application to Whole Lakes or Reservoirs

Where single applications to whole lakes or reservoirs are desired, Avast! WDG may be applied at an application rate of 16 to 90 ppb. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the section of this label entitled Application Rate Calculation -Ponds, Lakes and Reservoirs. Choose an application rate from the following table to meet the aquatic plant management objective. Where greater plant selectivity is desired, such as when controlling Eurasian watermilfoil and curlyleaf pondweed, an application rate lower in the rate range may be chosen. For other plant species, an aquatic specialist should be contacted to determine when to choose application rates lower in the rate range to meet specific plant management goals. Use the higher rate within the rate range where there is a dense weed mass or when treating more difficult to control plant species. Retreatments may be required to control more difficult to control species or in the event of a heavy rainfall event where dilution of the treatment concentration has occurred. In these cases, a second application or more may be required; however, the sum of all applications may not exceed 150 ppb per annual growth cycle. Refer to the following section, Split or Multiple Applications to Whole Lakes or Reservoirs, for guidelines and maximum rate allowed.

Average Water Depth	Pounds of Avast! WDG per Treated Surface Acre		
of Treatment Site (feet)	16 ppb	90 ppb	
1	0.9	5.0	
2	1.7	10.0	
3	2.6	15.0	
4	3.5	20.0	
5	4.3	25.0	
6	5.2	30.0	
7	6.0	34.0	
8	6.9	39.0	
9	7.8	44.0	
10	8.6	49.0	
11	9.5	54.0	
12	10.4	59.0	
13	11.2	64.0	
14	12.1	68.0	
15	13.0	73.0	

16	13.8	78.0
17	14.7	83.0
18	15.6	88.0
19	16.4	93.0
20	17.3	98.0

# Split or Multiple Applications to Whole Lakes or Reservoirs

To meet certain plant management objectives, split or multiple applications may be desired in making whole lake treatments. Split or multiple application programs are desirable when the objective is to use the minimum effective dose and to maintain this lower dose for sufficient time to ensure efficacy and enhance selectivity. In these situations, the lower rates (16 to 75 ppb) within the rate range may be used. In controlling Eurasian watermilfoil and curlyleaf pondweed and where greater plant selectivity is desired, an application rate lower in the rate range may be chosen. For other plant species, an aquatic specialist should be contacted to determine when to choose application rates lower in the rate range to meet specific plant management goals. For split or repeated applications, the sum of all applications must not exceed 150 ppb per annual growth cycle.

**Note**: In treating lakes or reservoirs that contain potable water intakes and the application requires treating within ½ mile of a potable water intake, no single application can exceed 20 ppb. Additionally, the sum of all applications must not exceed 150 ppb per annual growth cycle.

#### Partial Lake or Reservoir Treatments

Where dilution of Avast! WDG with untreated water is anticipated, such as in partial lake or reservoir treatments, split or multiple applications may be used to extend the contact time with the target plants. The application rate and use frequency of Avast! WDG in a partial lake is highly dependent upon the treatment area. An application rate at the higher end of the specified rate range may be required and frequency of applications will vary depending upon the potential for untreated water to dilute the Avast! WDG concentration in the treatment area. Use a rate at the higher end of the rate range where greater dilution with untreated water is anticipated.

#### Application Sites Greater than ¼ Mile from a Functioning Potable Water Intake

For single applications, Avast! WDG may be applied at application rates from 45 to 150 ppb. Split or multiple applications may be made, however, the sum of all applications must not exceed 150 ppb per annual growth cycle. Split applications should be conducted to maintain a sufficient concentration in the target area for a period of 45 days or longer. The use of FASTEST or other appropriate means of analysis is recommended to maintain the desired concentration in the target area over time.

#### Application Sites Within ¼ Mile of a Functioning Potable Water Intake

In treatment areas that are within ¼ mile of a potable water intake, no single application can exceed 20 ppb. When utilizing split or repeated applications of Avast! WDG for sites which contain a potable water intake, FASTEST or other appropriate means of analysis is required to determine the actual concentration in the water. Additionally, the sum of all

applications must not exceed 150 ppb per annual growth cycle.

# Application Rate Calculation — Ponds, Lakes and Reservoirs

The amount of Avast! WDG to be applied to provide the desired ppb concentration of active ingredient equivalents in treated water may be calculated as follows:

Pounds of Avast! WDG required per treated surface acre = Average water depth of treatment site (feet) **x** Desired ppb concentration of active ingredient equivalents **x** 0.054

For example, the pounds per acre of Avast! WDG required to provide a concentration of 25 ppb of active ingredient equivalents in water with an average depth of 5 feet is calculated as follows:

 $5 \times 25 \times 0.054 = 6.75$  pounds per treated surface acre.

**Note:** Calculated rates may not exceed the maximum allowable rate in pounds per treated surface acre for the water depth listed in the application rate table for the site to be treated.

# **Application to Drainage Canals, Irrigation Canals and Rivers**

#### Static Canals

In static drainage and irrigation canals, apply Avast! WDG at the rate of 20 to 40 pounds per surface acre.

### Moving Water Canals and Rivers

The performance of Avast! WDG will be enhanced by restricting or reducing water flow. In slow moving bodies of water, use an application technique that maintains a concentration of 10 to 40 ppb in the target area for a minimum of 45 days. Avast! WDG can be applied by split or multiple broadcast applications or by metering in the product to provide a uniform concentration of the herbicide based upon the flow pattern. The use of FASTEST or other appropriate means of analysis is recommended to maintain the desired concentration in the target area over time.

# Static or Moving Canals or Rivers Containing a Functioning Potable Water Intake

In treating a static or moving water canal or river which contains a functioning potable water intake, applications of Avast! WDG greater than 20 ppb must be made more than ½ mile from a functioning potable water intake. Applications of less than 20 ppb may be applied within ¼ mile from a functioning potable water intake; however, if applications of Avast! WDG are made within ¼ mile from a functioning potable water intake, the FASTEST or other appropriate means of analysis must be utilized to demonstrate that concentrations do not exceed 150 ppb at the potable water intake.

# Application Rate Calculation — Drainage Canals, Irrigation Canals and Rivers

The amount of Avast! WDG to be applied through a metering system to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

- 1. Average flow rate (feet per second) **x** Average width (feet) **x** Average depth (feet) **x** 0.9 = Cubic feet per second (CFS)
- 2. CFS x 1.98 = acre-feet per day (water movement)
- 3. Acre-feet per day **x** desired ppb **x** 0.054 = Pounds of Avast! WDG required per day

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

# **Container Handling:**

**Non-refillable, rigid container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Non-refillable, non-rigid container. DO NOT reuse or refill this container. Completely empty liner into application equipment by shaking and tapping sides and bottom to loosen clinging particles. If not emptied in this manner, the bag may be considered an acute hazardous waste and must be disposed of in accordance with local, state and federal regulations. When completely empty, offer for recycling if available or dispose of in a sanitary landfill or by incineration or by other procedures approved by state and local authorities. If outer packaging is contaminated and cannot be reused, dispose of it in the manner required for its liner.

**Non-refillable container. DO NOT reuse or refill this container.** Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or by other procedures approved by state and local authorities.

<u>Warranty Disclaimer</u>: SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

<u>Misuse</u>: Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <a href="http://seprolabels.com/terms">http://seprolabels.com/terms</a> or scan the image below.



© Copyright \_\_\_\_ SePRO Corporation

Avast! is a registered trademark of SePRO Corporation.



FLURIDONE GROUP 12 HERBICIDE

# **Avast! WDG**

**AQUATIC HERBICIDE** 

# **Active Ingredient**

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1 <i>H</i> )-pyridinone	5.0%
Other Ingredients	<u>95.0%</u>
TOTAL	100.0%

Contains 2 pounds of fluridone per 40-pound container.

# Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

	FIRST AID	
If in eyes	Hold eye open and rinse slowly and gently with water for 15 - 20	
	minutes. Remove contact lenses, if present, after the first 5 minutes,	
	then continue rinsing eye.	
	Call a poison control center or doctor for treatment advice.	
If on skin or	Take off contaminated clothing.	
clothing	Rinse skin immediately with plenty of water for 15 - 20 minutes.	
	Call a poison control center or doctor for treatment advice.	
If swallowed	Call a poison control center or doctor immediately for treatment advice.	
	Have person sip a glass of water if able to swallow.	
	Do not induce vomiting unless told to do so by a poison control center or	
	doctor.	
	Do not give anything by mouth to an unconscious person.	
If inhaled	Move person to fresh air.	
	If person is not breathing, call 911 or an ambulance, then give artificial	
	respiration, preferably mouth-to-mouth if possible.	
	Call a poison control center or doctor for further treatment advice.	
HOTLINE NUMBER		
	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if Swallowed, Absorbed Through Skin or Inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

# **Container Handling**

**Non-refillable Container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

See attached booklet for complete container disposal directions including triple rinsing and pressure rinsing instructions.

Refer to the label booklet for additional Precautionary Statements and the Directions for Use.

**NOTICE:** Read the entire label before using. Use only according to label directions. **Before** buying or using this product, read *Warranty Disclaimer* and *Misuse* statements inside label booklet. If terms are unacceptable, return at once unopened.

EPA Reg. No. 67690-32 FPL20210720	EPA Est. No [P/N]
SePRO Corporation • 11550 N. Meridi	an Street, Suite 600 • Carmel, IN 46032, U.S.A.
Aquatic Herbicide	Net Weight (Non-refillable)

#### TERMS AND CONDITIONS OF USE

If terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label as well as the *Inherent Risks of Use* and *Limitation of Remedies* statements below are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer, Misuse, Inherent Risks of Use,* and *Limitation of Remedies.* 

#### INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including use under conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), the presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer and/or user of the product.

#### LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use, Inherent Risks of Use* and *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use, Inherent Risks of Use* and *Limitation of Remedies* in any manner.