

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460 EPA Reg. Number:

67690-45

Date of Issuance:

1 0 MAR 2008

Date of Expiration:

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

SonarOne

Name and Address of Registrant (include ZIP Code):

SePRO Corporation

11550 N. Meridian St., Ste. 600

Carmel, IN 46032

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you make the following revisions:

- 1. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar product to submit data.
- 2. Add the statement, "EPA Reg. No. 67690-45"
- 3. On page 1, modify the Hazards to Humans and Domestic Animals statement beginning, "Harmful if swallowed..." to read as follows: "Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. [Wear protective eyewear.]*

*[Protective eyewear may be specified, if appropriate]

- 4. On page 1, move the "If swallowed" box to the top of the First Aid statement. Add, "immediately" to the first bullet, after the word, "doctor."
- 5. On page 14, under the Warranty Disclaimer section, add the words, "To the extent permitted by law," ahead of the word "SePRO" at the beginning of the second sentence. Likewise, on page 15, add the words, "To the extent permitted by law," ahead of the word SePRO at the beginning of the second paragraph.
- 6. Submit one copy of Final Printed Label before the product is released for shipment.

vanne S. Miller

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Joanne I. Miller

Product Manager 23

Herbicide Branch

Registration Division (7505P)

Date:

1 0 MAR 2008

page 2 EPA Reg. No. 67690-45

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

ith COMMENTS in EPA Letter Deted



registered under EPA Reg. No.



SonarOne* 67690-45

AN HERBICIDE FOR MANAGEMENT OF AQUATIC VEGETATION IN FRESH WATER PONDS, LAKES, RESERVOIRS, POTABLE WATER SOURCES, DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS.

Active Ingredient

| fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone | 5.0% |
|---|-------|
| Other Ingredients | 95.0% |
| TOTAL | |
| Contains 0.05 pound active ingredient per pound of product | |

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children **CAUTION / PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

Harmful If Swallowed, Absorbed Through Skin, Or If Inhaled. Avoid breathing of dust or contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

| William Strain and Control Control | militari mani mani na na mani alia 1920. — 20. Emili mani man da aliang maning ang maning ang maning ang maning |
|------------------------------------|--|
| (1911年) 在1911年 (1911年) | FIRSTAID |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin | Take off contaminated clothing. |
| or | Rinse skin immediately with plenty of water for 15 - 20 minutes. |
| clothing | Call a poison control center or doctor for treatment advice. |
| If swallowed | Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. |
| | Do not induce vomiting unless told to do so by a poison control center or doctor. |
| | Do not give anything by mouth to an unconscious person. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. |

4317

Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

ENVIRONMENTAL HAZARDS

Do not apply to water except as specified on the label. Do not contaminate water outside the intended treatment area by disposal of equipment washwaters. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas. Trees and shrubs growing in water treated with **SonarOne** herbicide may occasionally develop chlorosis.

Follow use directions carefully so as to minimize adverse effects on non-target organisms. In order to avoid impact on threatened or endangered aquatic plant or animal species, users must consult their State Fish and Game Agency or the U.S. Fish and Wildlife Service before making applications.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous-Waste Representative at the nearest EPA Regional Office for guidance.

Nonrefillable Container Disposal: Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) after emptying. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Refillable Container Disposal: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. If returning the container for refilling, seal all openings which have been opened during use and return the empty container to a collection site designated by SePRO Corporation. If the container has been damaged and cannot be returned according to the recommended procedures <u>OR</u> to clean the container before final disposal, empty the remaining contents from this container into application equipment ensuring no product remains in the container.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

Refer to the label booklet for the Directions for Use.

NOTICE: Read the entire label before using. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies inside label booklet.

^{*}Trademark of SePRO Corporation

EPA Reg. No. 67690-XX FPL100207

SonarOne*, 67690-XX EPA Est. No. XXXXX-XX-XXX5 7 7 17 SPC-XXXXXX

SePRO Corporation • 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Aquatic Herbicide

Net Contents ____ pounds



SonarOne*

AN HERBICIDE FOR MANAGEMENT OF AQUATIC VEGETATION IN FRESH WATER PONDS, LAKES, RESERVOIRS, POTABLE WATER SOURCES, DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS.

Active Ingredient

| fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone | 5.0% |
|---|-------|
| Other Ingredients | 95.0% |
| TOTAL | |
| Contains 0.05 pound active ingredient per pound of product. | |

Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

| COLUMN TO THE CARD CONTRACTOR | STANDARD CONTRACTOR TO CONTRACTOR TO A CONTRACTOR AND A C | | | | |
|--|--|--|--|--|--|
| | FIRST AID | | | | |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15 - 20 | | | | |
| | minutes. Remove contact lenses, if present, after the first 5 | | | | |
| minutes, then continue rinsing eye. | | | | | |
| Call a poison control center or doctor for treatment advice | | | | | |
| If on skin | Take off contaminated clothing. | | | | |
| or | Rinse skin immediately with plenty of water for 15 - 20 minutes. | | | | |
| clothing | Call a poison control center or doctor for treatment advice. | | | | |
| lf | Call a poison control center or doctor for treatment advice. | | | | |
| swallowed | Have person sip a glass of water if able to swallow. | | | | |
|) | Do not induce vomiting unless told to do so by a poison control | | | | |
| center or doctor. | | | | | |
| Do not give anything by mouth to an unconscious person | | | | | |
| If inhaled Move person to fresh air. | | | | | |
| | If person is not breathing, call 911 or an ambulance, then give | | | | |
| | artificial respiration, preferably mouth-to-mouth if possible. | | | | |
| | Call a poison control center or doctor for further treatment advice. | | | | |
| | duct container or label with you when calling a poison control center or | | | | |
| | ing for treatment. In case of emergency endangering health or the | | | | |
| environment involving this product, call INFOTRAC at 1-800-535-5053. | | | | | |

Refer to the inside of the label booklet for additional precautionary information and Directions for Use including Storage and Disposal.

3-17

NOTICE: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitation of Remedies* inside label booklet.

EPA Reg. No. 67690-XX FPL100207 EPA Est. No. XXXXX-XX-XXX SPC-XXXXXX

SePRO Corporation • 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Aquatic Herbicide

Net Contents ____ pounds

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Harmful If Swallowed, Absorbed Through Skin, Or If Inhaled. Avoid breathing of dust or contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply to water except as specified on the label. Do not contaminate water outside the intended treatment area by disposal of equipment washwaters. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas. Trees and shrubs growing in water treated with **SonarOne** herbicide may occasionally develop chlorosis.

Follow use directions carefully so as to minimize adverse effects on non-target organisms. In order to avoid impact on threatened or endangered aquatic plant or animal species, users must consult their State Fish and Game Agency or the U.S. Fish and Wildlife Service before making applications.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

GENERAL INFORMATION

SonarOne herbicide is a selective systemic aquatic herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, drainage canals, irrigation canals, and rivers. **SonarOne** is a pelleted formulation containing 5% fluridone. **SonarOne** is absorbed from water by plant shoots and from hydrosoil by the roots of aquatic vascular plants. It is important to maintain **SonarOne** in contact with the target plants for as long as possible. Rapid water movement or any condition which results in rapid dilution of **SonarOne** in treated water will reduce its effectiveness. In susceptible plants, **SonarOne** inhibits the formation of carotene. In the absence of carotene, chlorophyll is rapidly degraded by sunlight.

Herbicidal symptoms of **SonarOne** appear in seven to ten days and appear as white (chlorotic) or pink growing points. Under optimum conditions 30 - 90 days are required before the desired level of aquatic weed management is achieved with **SonarOne**. Species susceptibility to **SonarOne** may vary depending on time of year, stage of growth and water movement. For best results, apply **SonarOne** prior to initiation of weed growth or when weeds begin active growth. Application to mature target plants may require higher application rates and may take longer to control.

SonarOne is not corrosive to application equipment.

This label provides recommendations on the use of a chemical analysis for the active ingredient. SePRO Corporation recommends the use of an Enzyme-Linked Immunoassay (ELISA Test) for the determination of the active ingredient concentration in the water. Contact SePRO Corporation to incorporate this test, known as a FasTEST, into your treatment program. Other proven chemical analyses for the active ingredient may also be used. The chemical analysis, FasTEST, is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredient in the water.

Application rates are provided in pounds of **SonarOne** to achieve a desired concentration of the active ingredient in part per billion (ppb). **The maximum application rate or sum of all application rates is 90 ppb in ponds and 150 ppb in lakes and reservoirs per annual growth cycle.** This maximum concentration is the amount of product calculated as the target application rate, NOT determined by testing the residues of the active ingredient in the treated water.

GENERAL USE PRECAUTIONS

- Obtain Required Permits: Consult with appropriate state or local water authorities before applying this product to public waters. Permits and/or posting treatment notification may be required by state or local public agencies.
- New York State: Application of SonarOne is not permitted in waters less than two
 (2) feet deep.
- Hydroponic Farming: Do not use SonarOne treated water for hydroponic farming.
- Greenhouse and Nursery Plants: Do not use SonarOne treated water for irrigating greenhouse or nursery plants unless a FasTEST assay has been run and confirmed that residues are less than 1 ppb.
- Water Use Restrictions Following Application with SonarOne (Days)

| | Application Rate | | Fishing | The second secon | Livestock/Pet Consumption | Irrigation th |
|---|------------------|---|----------|--|------------------------------|--------------------------|
| | Maximum | | | | | See irrigation |
| | Rate (150 | 0 | . 0 | . 0 | 0 | instructions |
| Ì | ppb) or less | | <u> </u> | | | below |

Note below, under *Potable Water Intakes*, the information for application of **SonarOne** within ¼ miles (1,320) feet of a functioning potable water intake.

^{††} Note below, under *Irrigation*, specific time frames or fluridone residues that provide the widest safety margin for irrigating with fluridone treated water.

• Potable Water Intakes: Concentrations of the active ingredient fluridone up to 150 ppb are allowed in potable water sources; however, in lakes and reservoirs or other sources of potable water, <u>DO NOT APPLY</u> SonarOne at application rates greater than 20 ppb within one-fourth (¹/₄) mile (1,320 feet) of any functioning potable water intake. At application rates of 8 - 20 ppb, SonarOne <u>MAY BE APPLIED</u> within ¼ mile where functioning potable water intakes are present. NOTE: Existing potable water intakes which are no longer in use, such as those replaced by connections to potable water wells or a municipal water system, are not considered to be functioning potable water intakes.

^{*}Trademark of SePRO Corporation

Irrigation: Irrigation with SonarOne treated water may result in injury to the
irrigated vegetation. SePRO Corporation recommends following these precautions
and informing those who irrigate from areas treated with SonarOne of the irrigation
time frames or water FasTEST assay requirements presented in the table below.
These time frames and FasTEST assay recommendations are suggestions which
should be followed to reduce the potential for injury to vegetation irrigated with water
treated with SonarOne. Greater potential for crop injury occurs where SonarOne
treated water is applied to crops grown on low organic and sandy soils.

| | ilden sinden s | Days After A | pplication |
|---|---------------------------|--|--|
| Application Site | Established Tree Crops | Established Row Crops/ Turf/Plants | Newly Seeded Crops/Seedbeds or Areas to be Planted Including Overseeded Golf Course Greens |
| | | | • • • • • • • • • • • • • • • • • • • |
| Ponds and Static Canals † | 7 | 30 | FasTEST assay required |
| Ponds and Static Canals [†] Canals | 7 | 30 7 | FasTEST assay required FasTEST assay required |
| | 7 7 7 | 30 7 7 | |

^T For purposes of **SonarOne** labeling, a pond is defined as a body of water 10 acres or less in size. A lake or reservoir is greater than 10 acres.

^{††} In lakes and reservoirs where one-half or greater of the body of water is treated, use the pond and static canal irrigation precautions.

Where the use of SonarOne treated water is desired for irrigating crops prior to the time frames established above, the use of FasTEST assay is recommended to measure the concentration in the treated water. Where FasTEST has determined that concentrations are less than 10 parts per billion, there are no irrigation precautions for irrigating established tree crops, established row crops or turf. For tobacco, tomatoes, peppers or other plants within the Solanaceae Family and newly seeded crops or newly seeded grasses such as overseeded golf course greens, do not use SonarOne treated water if concentration are greater than 5 ppb; furthermore, when rotating crops, do not plant members of the Solanaceae family in land that has been previously irrigated with fluridone concentrations in excess of 5 ppb. It is recommended that an aquatic specialist be consulted prior to commencing irrigation of these sites.

PLANT CONTROL INFORMATION

SonarOne selectivity is dependent upon dosage, time of year, stage of growth, method of application, and water movement. The following categories: controlled, partially controlled, and not controlled, are provided to describe expected efficacy under ideal treatment conditions using higher to maximum label rates. Use of lower rates will increase selectivity of some species listed as controlled or partially controlled. Additional aquatic plants may be controlled, partially controlled, or tolerant to SonarOne. Consult an aquatic specialist prior to application of SonarOne to determine a plant's susceptibility to SonarOne. NOTE: algae (chara, nitella, and filamentous species are not controlled by SonarOne)

VASCULAR AQUATIC PLANTS CONTROLLED BY SonarOne †

Submersed Plants:

bladderwort (*Utricularia* spp.)
common coontail (*Ceratophyllum demersum*)[†]
common Elodea (*Elodea canadensis*)[†]
egeria, Brazilian Elodea (*Egeria densa*)
fanwort, Cabomba (*Cabomba caroliniana*)
hydrilla (*Hydrilla verticillata*)
naiad (*Najas* spp.)[†]
pondweed (*Potamogeton* spp., except Illinois
pondweed)[†]
watermilfoil (*Myriophyllum* spp. except variable-leaf milfoil)

Shoreline Grasses:

paragrass (Urochloa mutica)

Floating Plants:

salvinia (Salvinia spp.) duckweed (*Lemna*[†], *Spirodela*[†], and *Landoltia* spp.) mosquito fern (*Azolla caroliniana*) [†]

VASCULAR AQUATIC PLANTS PARTIALLY CONTROLLED BY SonarOne:

Submersed Plants:

Illinois pondweed (*Potamogeton illinoensis*) limnophila (*Limnophila sessiliflora*) tapegrass, American eelgrass (*Vallisneria americana*) watermilfoil--variable-leaf (*Myriophyllum heterophyllum*)

Shoreline Grasses:

barnyardgrass (*Echinochloa crusgalli*) giant cutgrass (*Zizaniopsis miliacea*) reed canarygrass (*Philaris arundinaceae*) southern watergrass (*Hydrochloa caroliniensis*) torpedograss (*Panicum repens*)

Emersed Plants:

alligatorweed (Alternanthera philoxeroides)
American lotus (Nelumbo lutea) cattail (Typha spp.)
creeping waterprimrose (Ludwigia peploides) parrotfeather (Myriophyllum aquaticum) smartweed (Polygonum spp.) spatterdock (Nuphar luteum) spikerush (Eleocharis spp.) waterlily (Nymphaea spp.) waterpurslane (Ludwigia palustris) watershield (Brasenia schreberi)

VASCULAR AQUATIC PLANTS NOT CONTROLLED BY SonarOne:

Emersed Plants:

American frogbit (Limnobium spongia) arrowhead (Sagittaria spp.) bacopa (Bacopa spp.) big floatingheart, banana lily (Nymphoides aquatica) bulrush (Scirpus spp.) pickerelweed, lanceleaf (Pontederia spp.) rush (Juncus spp.) water pennywort (Hydrocotyle spp.)

Floating Plants:

floating waterhyacinth (Eichhornia crassipes)
waterlettuce (Pistia stratiotes)

Shoreline Grasses:

maidencane (Panicum hemitomon)

[†] Species denoted are native plants that are often tolerant to **SonarOne** at lower use rates. Please consult an aquatic specialist for recommended **SonarOne** use rates when selective control of exotic species is desired.

APPLICATION DIRECTIONS

The aquatic plants present in the treatment site should be identified prior to application to determine their susceptibility to **SonarOne**. It is important to determine the area (acres) to be treated and the average depth in order to select the proper application rate. Do not exceed the maximum labeled rate for a given treatment site per annual growth cycle.

APPLICATION TO PONDS

SonarOne may be applied to the entire surface area of a pond. For single applications, rates may be selected to provide 30 - 90 ppb to the treated water, although actual concentrations in treated water may be substantially lower at any point in time due to the slow-release formulation of this product. When treating for optimum selective control, lower rates may be applied for sensitive target species. Use the higher rate within the rate range where there is a dense weed mass, when treating more difficult to control species, and for ponds less than 5 acres in size with an average depth less than 4 feet. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the *Application Rate Calculation—Ponds, Lakes and Reservoirs* section of this label. Split or multiple applications are recommended where dilution of treated water is anticipated; however, the sum of all applications should total 30 - 90 ppb and must not exceed a total of 90 ppb per annual growth cycle.

| Average Water Depth of Treatment Site | Pounds of SonarOne per | |
|--|------------------------|--------|
| (feet) | 45 ppb | 90 ppb |
| 1 | 2.5 | 5 |
| 2 | 5 | 10 |
| 3 | 7.5 | 15 |
| 4 | 10 | 20 |
| 5 | 12.5 | 25 |
| 6 | 15 | 30 |
| 7 | 17 · | 34 |
| 8 | 19.5 | 39 |
| 9 | 22 | 44 |
| 10 | 24.5 | 49 |

APPLICATION TO LAKES AND RESERVOIRS

The following treatments are recommended for treating both whole lakes or reservoirs and partial areas of lakes or reservoirs (bays, etc.). For best results in treating partial lakes and reservoirs, **SonarOne** treatment areas should be a minimum of 5 acres in size. Treatment of areas smaller than 5 acres or treatment of narrow strips such as boat lanes or shorelines may not produce satisfactory results due to dilution by untreated water. Rate ranges are provided as a guide to include a wide range of environmental factors, such as target species, plant susceptibility, selectivity and other aquatic plant management objectives. Application rates and methods should be selected to meet the specific lake/reservoir aquatic plant management goals.

NOTE: In treating lakes or reservoirs that contain potable water intakes and where the application requires treating within one-fourth (¼) mile of a potable water intake, no single application can exceed 20 ppb. Additionally, the sum of all applications cannot exceed 150 ppb per annual growth cycle.

Whole Lake or Reservoir Treatments (Limited or No Water Discharge)

Single Application to Whole Lakes or Reservoirs - Where single applications to
whole lakes or reservoirs are desired, apply SonarOne at an application rate of 16 90 ppb. Application rates necessary to obtain these concentrations in treated water
are shown in the following table. For additional application rate calculations, refer to
the Application Rate Calculation—Ponds, Lakes and Reservoirs section of this label.

Choose an application rate to meet the aquatic plant management objective. Where greater plant selectivity is desired such as when controlling Eurasian watermilfoil and curlyleaf pondweed, choose an application rate lower in the rate range. For other plant species, SePRO recommends contacting an aquatic specialist in determining when to choose application rates lower in the rate range to meet specific plant management goals. Use the higher rate within the rate range where there is a dense weed mass or when treating more difficult to control plant species or in the event of a heavy rainfall event where dilution has occurred. In these cases, a second application or more may be required; however, the sum of all applications cannot exceed 150 ppb per annual growth cycle. Refer to the next section of this label, *Split or Multiple Applications to Whole Lakes or Reservoirs*, for guidelines and maximum rate allowed.

| Average Water Depth | Pounds of SonarOne Per Treated Surface Acre | |
|---------------------|---|--------|
| (feet) | 16 ppb | 90 ppb |
| 1 | 0.9 | 5 |
| 2 | 1.7 | 10 |
| 3 | 2.6 | 15 |
| 4 | 3.5 | 20 |
| 5 | 4.3 | 25 |
| 6 | 5.2 | 30 |
| 7 | 6.0 | 34 |
| 8 | 6.9 | 39 |
| 9 | 7.8 | 44 |
| 10 | 8.6 | 49 |
| 11 | 9.5 | 54 |
| 12 | 10.4 | 59 |
| 13 | 11.2 | 64 |
| 14 | 12.1 | 68 |
| 15 | 13.0 | 73 |
| 16 | 13.8 | 78 |
| 17 | 14.7 | 83 |
| 18 | 15.6 | 88 |
| 19 | 16.4 | 93 |
| 20 | 17.3 | 98 |

^{*}Trademark of SePRO Corporation

• Split or Multiple Applications to Whole Lakes or Reservoirs - To meet certain plant management objectives, split or multiple applications may be desired in making whole lake treatments. Split or multiple application programs are desirable when the objective is to use the minimum effective dose and to maintain this lower dose for the sufficient time to ensure efficacy and enhance selectivity. Under these situations, use the lower rates (16 - 75 ppb) within the rate range. In controlling Eurasian watermilfoil and curlyleaf pondweed and where greater plant selectivity is desired, choose an application rate lower in the rate range. For other plant species, SePRO recommends contacting an aquatic specialist in determining when to choose application rates lower in the rate range to meet specific plant management goals. For split or repeated applications, the sum of all applications must not exceed 150 ppb per annual growth cycle.

Partial Lake or Reservoir Treatments

Where dilution of **SonarOne** with untreated water is anticipated, such as in partial lake or reservoir treatments, split or multiple applications may be used to extend the contact time to the target plants. The application rate and use frequency of **SonarOne** in a partial lake is highly dependent upon the treatment area. Higher application rates may be required and frequency of applications will vary depending upon the potential of untreated water diluting the **SonarOne** concentration in the treatment area. Use higher rates where greater dilution with untreated water is anticipated.

- Application Sites Greater Than ¼ Mile from a Functioning Potable Water Intake For single applications, apply SonarOne at application rates from 45 150 ppb. Split or multiple applications may be made, however, the sum of all applications cannot exceed 150 ppb per annual growth cycle. Split applications should be conducted to maintain a sufficient concentration in the target area for a period of 45 days or longer. The use of FasTEST is recommended to maintain the desired concentration in the target area over time.
- Application Sites Within ¼ Mile of a Functioning Potable Water Intake In treatment areas that are within ¼ mile of a potable water intake, no single application can exceed 20 ppb. When utilizing split or repeated applications of SonarOne for sites which contain a potable water intake, FasTEST is required to determine the actual concentration in the water. Additionally, the sum of all applications cannot exceed 150 ppb per annual growth cycle.

<u>APPLICATION RATE CALCULATION — PONDS, LAKES AND RESERVOIRS</u>

The amount of **SonarOne** to be applied to provide the desired ppb concentration of active ingredient equivalents in treated water may be calculated as follows:

Pounds of **SonarOne** required per treated acre = Avg. water depth of treatment site **x** Desired ppb concentration of active ingredient equivalents **x** 0.054

For example: the pounds per acre of **SonarOne** required to provide a concentration of 25 ppb of active ingredient equivalents in water with an average depth of 5 feet is calculated as follows:

 $5 \times 25 \times 0.054 = 6.75$ pounds per treated surface acre.

NOTE: Calculated rates should not exceed the maximum allowable rate in pounds per treated surface acre for the water depth listed in the application rate table for the site to be treated.

APPLICATION TO DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS

- Static Canals In static drainage and irrigation canals, SonarOne should be applied at the rate of 20 40 pounds per surface acre.
- Moving Water Canals and Rivers The performance of SonarOne will be enhanced by restricting or reducing water flow. In slow moving bodies of water use an application technique that maintains a concentration of 10 40 ppb in the applied area for a minimum of 45 days. SonarOne can be applied by split or multiple broadcast applications or by metering in the product to provide a uniform concentration of the herbicide based upon the flow pattern. The use of FasTEST is recommended to maintain the desired concentration in the target area over time.
- Static or Moving Water Canals or Rivers Containing a Functioning Potable Water Intake In treating a static or moving water canal or river which contains a functioning potable water intake, applications of SonarOne greater than 20 ppb must be made more than ¼ mile from a functioning potable water intake. Applications less than 20 ppb may be applied within ¼ mile from a functioning potable water intake; however, if applications of SonarOne are made within ¼ mile from a functioning water intake, the FasTEST must be utilized to demonstrate that concentrations do not exceed 150 ppb at the potable water intake.

APPLICATION RATE CALCULATION-DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS

The amount of **SonarOne** to be applied through a metering system to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

- 1. Average flow rate (ft. per second) **x** average width (ft.) **x** average depth (ft.) **x** 0.9 = CFS (cubic feet per second)
- 2. CFS x 1.98 = acre feet per day (water movement)
- 3. Acre feet per day x desired ppb x 0.054 = pounds SonarOne required per day.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Nonrefillable Container Disposal: Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) after emptying. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Refillable Container Disposal: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. If returning the container for refilling, seal all openings which have been opened during use and return the empty container to a collection site designated by SePRO Corporation. If the container has been damaged and cannot be returned according to the recommended procedures <u>OR</u> to clean the container before final disposal, empty the remaining contents from this container into application equipment ensuring no product remains in the container.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner or application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. All such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

^{*}Trademark of SePRO Corporation

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* above and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or *Limitations of Remedies* in any manner.

© Copyright 2007 SePRO Corporation