

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

L4 JUN 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Amy Dugger-Ronyak SePRO Corporation 11550 N. Meridian Street, Suite 600 Carmel, IN 46032-4565

Subject:

SonarOne

EPA Registration Number 67690-45 Submission dated April 14, 2008

Response to Agency email dated March 27, 2008

Dear Ms. Dugger-Ronyak:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

- 1) Add an appropriate EPA Establishment Number to the label
- 2) Add appropriate Net Contents information to the label
- 3) On page 8, revise the statement "Application to mature target plants may require higher application rates and may take longer to control" to "Application to mature target plants may require an application rate at the higher end of the specified rate range and may take longer to control"
- 4) On page 9, in the subsection Irrigation, revise "SePRO Corporation recommends following these precautions and informing....." to "Follow these precautions and inform those who irrigate.... Also revise "These time frames and assay recommendations are suggestions which should be followed to reduce the potential for injury..." To "Follow the following time frames and assay directions to reduce the potential for injury..."
- 5) On page 9, revise "the use of a FasTest assay is recommended to measure the concentration in the treated water" to "use the FasTest assay to measure the concentration in the treated water"
- 6) On page 10, revise "Please consult an aquatic specialist for recommended SonarOne use rates when selective control of exotic species is desired" to "Please consult an aquatic specialist for recommended SonarOne use rates (not to exceed maximum labeled rates) when selective control of exotic species is desired"
- 7) On page 12, revise "Split or multiple applications are recommended where dilution...." to "Split or multiple applications may be used where dilution...."
- 8) On page 12, revise "The following treatments are recommended for treating both whole lakes..." to "The following treatments may be used for treating both whole lakes..."

Page 2 EPA Registration Number 67690-45

- 9) On page 13, revise "Choose an application rate to meet the aquatic plant management objective" to "Choose an application rate from the table below to meet the aquatic plant management objective"
- 10) On page 14, revise "Higher application rates may be required and frequency of applications...." to "An application rate at the higher end of the specified rate range may be required and frequency of applications...."
- 11) On page 14, revise "Use higher rates where greater dilution with untreated water is anticipated" to "Use a rate at the higher end of the rate range where greater dilution with untreated water is anticipated"
- 12) On page 14, revise "Calculated rates should not exceed the maximum allowable rate in pounds per" to "Calculated rates may not exceed the maximum allowable rate in pounds per"
- 13) On page 15, revise "In static drainage and irrigation canals, SonarOne should be applied at the rate of 20 to 40 pounds per surface acre" to "In static drainage and irrigation canals, apply SonarOne at the rate of 20 to 40 pounds per surface acre"

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records. If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

Joanne I Miller

Product Manager 23


Herbicide Branch

Registration Division (7505P)

vanne A. Miller

SonarOne* 67690-45

(Base label for Nonrefillable, Rigid containers 5 gal or less)



SonarOne*

AQUATIC HERBICIDE

In EPA Letter Dated:

4 JUN 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

67690-45

AN HERBICIDE FOR MANAGEMENT OF AQUATIC VEGETATION IN FRESH WATER PONDS, LAKES, RESERVOIRS, POTABLE WATER SOURCES, DRAINAGE CANALS. IRRIGATION CANALS AND RIVERS.

Active Ingredient

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone	5.0%
Other Ingredients	. <u>95.0%</u>
TOTAL	100.0%
Contains 0.05 pound active ingredient per pound of product.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

Harmful If Swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. Wear protective eyewear.

	FIRST AID
lf	Call a poison control center or doctor immediately for treatment advice.
swallowed	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by a poison control center or
	doctor.
l	Do not give anything by mouth to an unconscious person.
If in eyes	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue
	rinsing eye.
L	Call a poison control center or doctor for treatment advice.
If on skin	Take off contaminated clothing.
or	Rinse skin immediately with plenty of water for 15 - 20 minutes.
clothing	Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air.
	If person is not breathing, call 911 or an ambulance, then give artificial
	respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment advice.
Have the pro	duct container or label with you when calling a poison control contar or doctor

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053.

^{*}Trademark of SePRO Corporation

ENVIRONMENTAL HAZARDS

Do not apply to water except as specified on the label. Do not contaminate water outside the intended treatment area by disposal of equipment washwaters. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas. Trees and shrubs growing in water treated with SonarOne herbicide may occasionally develop chlorosis. Follow use directions carefully so as to minimize adverse effects on non-target organisms.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Nonrefillable Container Disposal (rigid, 5 gallons or less): Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times.

Refer to the label booklet for the Directions for Use.

NOTICE: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitation of Remedies* inside label booklet.

For additional information on our products, please visit www.sepro.com.

, , , , , , , , , , , , , , , , , , , ,	
EPA Reg. No. 67690-45 FPL041008	EPA Est. No
SePRO Corporation • 11550 N. Meridian	Street, Suite 600 • Carmel, IN 46032, U.S.A.
Aquatic Herbicide	Net contents

^{*}Trademark of SePRO Corporation



SonarOne*

AQUATIC HERBICIDE

AN HERBICIDE FOR MANAGEMENT OF AQUATIC VEGETATION IN FRESH WATER PONDS, LAKES, RESERVOIRS, POTABLE WATER SOURCES, DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS.

Active Ingredient

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone	5.0%
Other Ingredients	95.0%
TOTAL	
Contains 0.05 pound active ingredient per pound of product	•

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

Harmful If Swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. Wear protective eyewear.

The state of the s	
	FIRST AID
If	Call a poison control center or doctor immediately for treatment advice.
swallowed	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by a poison control center or
	doctor.
	Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue
	rinsing eye.
	Call a poison control center or doctor for treatment advice.
If on skin	Take off contaminated clothing.
or	Rinse skin immediately with plenty of water for 15 - 20 minutes.
clothing	Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air.
	If person is not breathing, call 911 or an ambulance, then give artificial
	respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment advice.
Have the are	dust container or lebel with you when calling a poison central center or dector

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

^{*}Trademark of SePRO Corporation

ENVIRONMENTAL HAZARDS

Do not apply to water except as specified on the label. Do not contaminate water outside the intended treatment area by disposal of equipment washwaters. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas. Trees and shrubs growing in water treated with SonarOne herbicide may occasionally develop chlorosis. Follow use directions carefully so as to minimize adverse effects on non-target organisms.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Refillable Container Disposal (rigid, greater than 5 gallons): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times: If returning the container for refilling, seal all openings which have been opened during use and return the empty container to a collection site designated by SePRO Corporation. If the container has been damaged and cannot be returned according to the recommended procedures, contact SePRO Corporation at 1-800-419-7779 to obtain proper handling instructions.

Refer to the label booklet for the Directions for Use.

NOTICE: Read the entire label before using. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies inside label booklet.

For additional information on our products, please visit www.sepro.com.

EPA Reg. No. 67690-45

EPA Est. No. _____

FPL041008

SPC-_____

SePRO Corporation • 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Aquatic Herbicide

Net contents _____



SonarOne*

AQUATIC HERBICIDE

AN HERBICIDE FOR MANAGEMENT OF AQUATIC VEGETATION IN FRESH WATER PONDS, LAKES, RESERVOIRS, POTABLE WATER SOURCES, DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS.

Active Ingredient fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl] Other Ingredients TOTAL Contains 0.05 pound active ingredient per pound of prod	<u>95.0%</u> 100.0%
Keep Out of Reach of Children CAUTION / PRECAUCIÓN Si usted no entiende la etiqueta, busque a alguien para detalle. (If you do not understand this label, find someon	
Refer to the inside of the label booklet for additional Directions for Use including Storage and Disposal.	precautionary information and
NOTICE: Read the entire label before using. Use only a buying or using this product, read Warranty Disclain Limitation of Remedies inside label booklet.	
For additional information on our products, please visit w	ww.sepro.com.
EPA Reg. No. 67690-45 FPL041008	EPA Est. No
SePRO Corporation • 11550 N. Meridian Street, Suite	600 • Carmel, IN 46032, U.S.A.

Aquatic Herbicide

Net contents

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Harmful If Swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. Wear protective eyewear.

	FIRST AID
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
or going for t	duct container or label with you when calling a poison control center or doctor, reatment. In case of emergency endangering health or the environment product, call INFOTRAC at 1-800-535-5053 .

ENVIRONMENTAL HAZARDS

Do not apply to water except as specified on the label. Do not contaminate water outside the intended treatment area by disposal of equipment washwaters. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas. Trees and shrubs growing in water treated with SonarOne herbicide may occasionally develop chlorosis. Follow use directions carefully so as to minimize adverse effects on non-target organisms.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

GENERAL INFORMATION

SonarOne herbicide is a selective systemic aquatic herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, drainage canals, irrigation canals, and rivers. SonarOne is a pelleted formulation containing 5% fluridone. SonarOne is absorbed from water by plant shoots and from hydrosoil by the roots of aquatic vascular plants. It is important to maintain SonarOne in contact with the target plants for as long as possible. Rapid water movement or any condition which results in rapid dilution of SonarOne in treated water will reduce its effectiveness. In susceptible plants, SonarOne inhibits the formation of carotene. In the absence of carotene, chlorophyll is rapidly degraded by sunlight.

Herbicidal symptoms of SonarOne appear in 7 - 10 days and appear as white (chlorotic) or pink growing points. Under optimum conditions 30 - 90 days are required before the desired level of aquatic weed management is achieved with SonarOne. Species susceptibility to SonarOne may vary depending on time of year, stage of growth and water movement. For best results, apply SonarOne prior to initiation of weed growth or when weeds begin active growth. Application to mature target plants may require higher application rates and may take longer to control.

SonarOne is not corrosive to application equipment.

This label provides recommendations on the use of a chemical analysis for the active ingredient. SePRO Corporation recommends the use of an Enzyme-Linked Immunoassay (ELISA Test) for the determination of the active ingredient concentration in the water. Contact SePRO Corporation to incorporate this test, known as a FasTEST, into your treatment program. Other proven chemical analyses for the active ingredient may also be used. The chemical analysis, FasTEST, is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredient in the water.

Application rates are provided in pounds of SonarOne to achieve a desired concentration of the active ingredient in part per billion (ppb). The maximum application rate or sum of all application rates is 90 ppb in ponds and 150 ppb in lakes and reservoirs per annual growth cycle. This maximum concentration is the amount of product calculated as the target application rate, NOT determined by testing the residues of the active ingredient in the treated water.

General Use Precautions

- Obtain Required Permits: Consult with appropriate state or local water authorities before applying this product to public waters. Permits and/or posting treatment notification may be required by state or local public agencies.
- New York State: Application of SonarOne is not permitted in waters less than two (2) feet deep.
- Hydroponic Farming: Do not use SonarOne treated water for hydroponic farming.
- Greenhouse and Nursery Plants: Do not use SonarOne treated water for irrigating greenhouse or nursery plants unless a FasTEST assay has been run and confirmed that residues are less than 1 ppb.
- Water Use Restrictions Following Application with SonarOne (Days)

Application Rate	Drinking [†]	Fishing		Livestock/Pet Consumption	Irrigation ^{††}
Maximum					See irrigation
Rate (150	0	0	0	0	instructions
ppb) or less				·	below

[†] Note below, under *Potable Water Intakes*, the information for application of SonarOne within ¼ miles (1,320) feet of a functioning potable water intake.

^{††} Note below, under *Irrigation*, specific time frames or fluridone residues that provide the widest safety margin for irrigating with fluridone treated water.

- Potable Water Intakes: Concentrations of the active ingredient fluridone up to 150 ppb are allowed in potable water sources; however, in lakes and reservoirs or other sources of potable water, do not apply SonarOne at application rates greater than 20 ppb within one-fourth (¹/₄) mile (1,320 feet) of any functioning potable water intake. At application rates of 8 20 ppb, SonarOne may be applied within ¼ mile where functioning potable water intakes are present. NOTE: Existing potable water intakes which are no longer in use, such as those replaced by connections to potable water wells or a municipal water system, are not considered to be functioning potable water intakes.
- Irrigation: Irrigation with SonarOne treated water may result in injury to the irrigated vegetation. SePRO Corporation recommends following these precautions and informing those who irrigate from areas treated with SonarOne of the irrigation time frames or water FasTEST assay requirements presented in the table below. These time frames and FasTEST assay recommendations are suggestions which should be followed to reduce the potential for injury to vegetation irrigated with water treated with SonarOne. Greater potential for crop injury occurs where SonarOne treated water is applied to crops grown on low organic and sandy soils.

		Days After A	pplication
		British period with the	Newly Seeded
		Established	Crops/Seedbeds or Areas to be Planted
Application Site	Established Tree Crops	Row Crops/ Turf/Plants	Including Overseeded Golf Course Greens
Ponds and Static Canals †	7	30	FasTEST assay required
Canals	7	7	FasTEST assay required
Rivers	7	. 7	FasTEST assay required
Lakes and Reservoirs ††	7	7	FasTEST assay required

[†] For purposes of SonarOne labeling, a pond is defined as a body of water 10 acres or less in size. A lake or reservoir is greater than 10 acres.

^{††} In lakes and reservoirs where one-half or greater of the body of water is treated, use the pond and static canal irrigation precautions.

Where the use of SonarOne treated water is desired for irrigating crops prior to the time frames established above, the use of FasTEST assay is recommended to measure the concentration in the treated water. Where FasTEST has determined that concentrations are less than 10 parts per billion, there are no irrigation precautions for irrigating established tree crops, established row crops or turf. For tobacco, tomatoes, peppers or other plants within the Solanaceae Family and newly seeded crops or newly seeded grasses such as overseeded golf course greens, do not use SonarOne

treated water if concentration are greater than 5 ppb; furthermore, when rotating crops, do not plant members of the Solanaceae family in land that has been previously irrigated with fluridone concentrations in excess of 5 ppb. It is recommended that an aquatic specialist be consulted prior to commencing irrigation of these sites.

PLANT CONTROL INFORMATION

SonarOne selectivity is dependent upon dosage, time of year, stage of growth, method of application, and water movement. The following categories: controlled, partially controlled, and not controlled, are provided to describe expected efficacy under ideal treatment conditions using higher to maximum label rates. Use of lower rates will increase selectivity of some species listed as controlled or partially controlled. Additional aquatic plants may be controlled, partially controlled, or tolerant to SonarOne. Consult an aquatic specialist prior to application of SonarOne to determine a plant's susceptibility to SonarOne. NOTE: algae (chara, nitella, and filamentous species) are not controlled by SonarOne.

Vascular Aquatic Plants Controlled By SonarOne: 1

Submersed Plants:

bladderwort (*Utricularia* spp.)
common coontail (*Ceratophyllum demersum*)[†]
common Elodea (*Elodea canadensis*)[†]
egeria, Brazilian Elodea (*Egeria densa*)
fanwort, Cabomba (*Cabomba caroliniana*)
hydrilla (*Hydrilla verticillata*)
naiad (*Najas* spp.)[†]
pondweed (*Potamogeton* spp., except Illinois pondweed)[†]
watermilfoil (*Myriophyllum* spp. except variable-leaf milfoil)

Floating Plants:

salvinia (*Salvinia* spp.) duckweed (*Lemna*[†], *Spirodela*[†], and *Landoltia* spp.) mosquito fern (*Azolla caroliniana*) [†]

Shoreline Grasses:

paragrass (*Urochloa mutica*)

¹Species denoted by a dagger (†) are native plants that are often tolerant to fluridone at lower use rates. Please consult an aquatic specialist for recommended SonarOne use rates when selective control of exotic species is desired.

Vascular Aquatic Plants Partially Controlled By SonarOne:

Submersed Plants:

Illinois pondweed (*Potamogeton illinoensis*) limnophila (*Limnophila sessiliflora*) tapegrass, American eelgrass (*Vallisneria americana*) watermilfoil--variable-leaf (*Myriophyllum heterophyllum*)

Emersed Plants:

alligatorweed (Alternanthera philoxeroides)

^{*}Trademark of SePRO Corporation

American lotus (Nelumbo lutea)
cattail (Typha spp.)
creeping waterprimrose (Ludwigia peploides)
parrotfeather (Myriophyllum aquaticum)
smartweed (Polygonum spp.)
spatterdock (Nuphar luteum)
spikerush (Eleocharis spp.)
waterlily (Nymphaea spp.)
waterpurslane (Ludwigia palustris)
watershield (Brasenia schreberi)

Shoreline Grasses:

barnyardgrass (*Echinochloa crusgalli*) giant cutgrass (*Zizaniopsis miliacea*) reed canarygrass (*Philaris arundinaceae*) southern watergrass (*Hydrochloa caroliniensis*) torpedograss (*Panicum repens*)

Vascular Aquatic Plants Not Controlled By SonarOne:

Emersed Plants:

American frogbit (Limnobium spongia)
arrowhead (Sagittaria spp.)
bacopa (Bacopa spp.)
big floatingheart, banana lily (Nymphoides aquatica)
bulrush (Scirpus spp.)
pickerelweed, lanceleaf (Pontederia spp.)
rush (Juncus spp.)
water pennywort (Hydrocotyle spp.)

Floating Plants:

floating waterhyacinth (Eichhornia crassipes) waterlettuce (Pistia stratiotes)

Shoreline Grasses:

maidencane (Panicum hemitomon)

NOTE: Algae (chara, nitella, and filamentous species) are not controlled by SonarOne.

APPLICATION DIRECTIONS

The aquatic plants present in the treatment site should be identified prior to application to determine their susceptibility to SonarOne. It is important to determine the area (acres) to be treated and the average depth in order to select the proper application rate. Do not exceed the maximum labeled rate for a given treatment site per annual growth cycle.

Application to Ponds

SonarOne may be applied to the entire surface area of a pond. For single applications, rates may be selected to provide 30 - 90 ppb to the treated water, although actual

^{*}Trademark of SePRO Corporation

concentrations in treated water may be substantially lower at any point in time due to the slow-release formulation of this product. When treating for optimum selective control, lower rates may be applied for sensitive target species. Use the higher rate within the rate range where there is a dense weed mass, when treating more difficult to control species, and for ponds less than 5 acres in size with an average depth less than 4 feet. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the *Application Rate Calculation—Ponds, Lakes and Reservoirs* section of this label. Split or multiple applications are recommended where dilution of treated water is anticipated; however, the sum of all applications should total 30 - 90 ppb and must not exceed a total of 90 ppb per annual growth cycle.

Average Water Depth of Treatment Site	Pounds of SonarOne per Treated Surface Acre		
(feet)	45 ppb	90 ppb	
1	2.5	5.0	
2	5.0	10.0	
3	7.5	15.0	
4	10.0	20.0	
5	12.5	25.0	
6	15.0	30.0	
7	17.0	34.0	
8	19.5	39.0	
9	22.0	44.0	
10	24.5	49.0	

Application to Lakes and Reservoirs

The following treatments are recommended for treating both whole lakes or reservoirs and partial areas of lakes or reservoirs (bays, etc.). For best results in treating partial lakes and reservoirs, SonarOne treatment areas should be a minimum of 5 acres in size. Treatment of areas smaller than 5 acres or treatment of narrow strips such as boat lanes or shorelines may not produce satisfactory results due to dilution by untreated water. Rate ranges are provided as a guide to include a wide range of environmental factors, such as target species, plant susceptibility, selectivity and other aquatic plant management objectives. Application rates and methods should be selected to meet the specific lake/reservoir aquatic plant management goals.

NOTE: In treating lakes or reservoirs that contain potable water intakes and where the application requires treating within one-fourth ($\frac{1}{4}$) mile of a potable water intake, no single application can exceed 20 ppb. Additionally, the sum of all applications cannot exceed 150 ppb per annual growth cycle.

A. Whole Lake or Reservoir Treatments (Limited or No Water Discharge)

Single Application to Whole Lakes or Reservoirs

Where single applications to whole lakes or reservoirs are desired, apply SonarOne at an application rate of 16 - 90 ppb. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional

^{*}Trademark of SePRO Corporation

application rate calculations, refer to the Application Rate Calculation—Ponds, Lakes and Reservoirs section of this label. Choose an application rate to meet the aquatic plant management objective. Where greater plant selectivity is desired such as when controlling Eurasian watermilfoil and curlyleaf pondweed, choose an application rate lower in the rate range. For other plant species, SePRO recommends contacting an aquatic specialist in determining when to choose application rates lower in the rate range to meet specific plant management goals. Use the higher rate within the rate range where there is a dense weed mass or when treating more difficult to control plant species or in the event of a heavy rainfall event where dilution has occurred. In these cases, a second application or more may be required; however, the sum of all applications cannot exceed 150 ppb per annual growth cycle. Refer to the section of this label entitled, Split or Multiple Applications to Whole Lakes or Reservoirs, for guidelines and maximum rate allowed.

Average Water Depth : of Treatment Site		of SonarOne
(feet)		90 ppb
1	0.9	5.0
2	1.7	10.0
3	2.6	15.0
4	3.5	20.0
5	4.3	25.0
6	5.2	30.0
7	6.0	34.0
8	6.9	39.0
9	7.8	44.0
10	8.6	49.0
11	9.5	54.0
12	10.4	59.0
13	11.2	64.0
14	12.1	68.0
15	13.0	73.0
16	13.8	78.0
17	.14.7	83.0
18	15.6	88.0
19	16.4	93.0
20	17.3	98.0

Split or Multiple Applications to Whole Lakes or Reservoirs

To meet certain plant management objectives, split or multiple applications may be desired in making whole lake treatments. Split or multiple application programs are desirable when the objective is to use the minimum effective dose and to maintain this lower dose for the sufficient time to ensure efficacy and enhance selectivity. Under these situations, use the lower rates (16 - 75 ppb) within the rate range. In controlling Eurasian watermilfoil and curlyleaf pondweed and where greater plant selectivity is desired, choose an application rate lower in the rate range. For other plant species, SePRO recommends contacting an aquatic specialist in determining when to choose application rates lower in the rate range to meet

specific plant management goals. For split or repeated applications, the sum of all applications must not exceed 150 ppb per annual growth cycle.

B. Partial Lake or Reservoir Treatments

Where dilution of SonarOne with untreated water is anticipated, such as in partial lake or reservoir treatments, split or multiple applications may be used to extend the contact time to the target plants. The application rate and use frequency of SonarOne in a partial lake is highly dependent upon the treatment area. Higher application rates may be required and frequency of applications will vary depending upon the potential of untreated water diluting the SonarOne concentration in the treatment area. Use higher rates where greater dilution with untreated water is anticipated.

Application Sites Greater Than ¼ Mile from a Functioning Potable Water Intake For single applications, apply SonarOne at application rates from 45 - 150 ppb. Split or multiple applications may be made, however, the sum of all applications cannot exceed 150 ppb per annual growth cycle. Split applications should be conducted to maintain a sufficient concentration in the target area for a period of 45 days or longer. The use of FasTEST is recommended to maintain the desired concentration in the target area over time.

Application Sites within ¼ Mile of a Functioning Potable Water Intake In treatment areas that are within ¼ mile of a potable water intake, no single application can exceed 20 ppb. When utilizing split or repeated applications of SonarOne for sites which contain a potable water intake, FasTEST is required to determine the actual concentration in the water. Additionally, the sum of all applications cannot exceed 150 ppb per annual growth cycle.

<u>Application Rate Calculation — Ponds, Lakes and Reservoirs</u>

The amount of SonarOne to be applied to provide the desired ppb concentration of active ingredient equivalents in treated water may be calculated as follows:

Pounds of SonarOne required per treated acre = Avg. water depth of treatment site **x** Desired ppb concentration of active ingredient equivalents **x** 0.054

For example: the pounds per acre of SonarOne required to provide a concentration of 25 ppb of active ingredient equivalents in water with an average depth of 5 feet is calculated as follows:

 $5 \times 25 \times 0.054 = 6.75$ pounds per treated surface acre.

NOTE: Calculated rates should not exceed the maximum allowable rate in pounds per treated surface acre for the water depth listed in the application rate table for the site to be treated.

Application to Drainage Canals, Irrigation Canals and Rivers

Static Canals

In static drainage and irrigation canals, SonarOne should be applied at the rate of 20 - 40 pounds per surface acre.

Moving Water Canals and Rivers

The performance of SonarOne will be enhanced by restricting or reducing water flow. In slow moving bodies of water use an application technique that maintains a concentration of 10 - 40 ppb in the applied area for a minimum of 45 days. SonarOne can be applied by split or multiple broadcast applications or by metering in the product to provide a uniform concentration of the herbicide based upon the flow pattern. The use of FasTEST is recommended to maintain the desired concentration in the target area over time.

Static or Moving Water Canals or Rivers Containing a Functioning Potable Water Intake In treating a static or moving water canal or river which contains a functioning potable water intake, applications of SonarOne greater than 20 ppb must be made more than ¼ mile from a functioning potable water intake. Applications less than 20 ppb may be applied within ¼ mile from a functioning potable water intake; however, if applications of SonarOne are made within ¼ mile from a functioning water intake, the FasTEST must be utilized to demonstrate that concentrations do not exceed 150 ppb at the potable water intake.

<u>Application Rate Calculation — Drainage Canals, Irrigation Canals and Rivers</u>

The amount of SonarOne to be applied through a metering system to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

- 1. Average flow rate (ft. per second) **x** average width (ft.) **x** average depth (ft.) **x** 0.9 = CFS (cubic feet per second)
- 2. CFS x 1.98 = acre feet per day (water movement)
- 3. Acre feet per day \mathbf{x} desired ppb \mathbf{x} 0.054 = pounds SonarOne required per day.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Nonrefillable Container Disposal (rigid, 5 gallons or less): Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times.

Refillable Container Disposal (rigid, greater than 5 gallons): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. If returning the container for refilling, seal all openings which have been opened during use and return the empty container to a collection site designated by SePRO Corporation. If the container has been damaged and cannot be returned according to the recommended procedures, contact SePRO Corporation at 1,800-419-7779 to obtain proper handling instructions.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent

SonarOne*, 67690-45

of SePRO Corporation or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitations of Remedies in any manner.

© Copyright ____ SePRO Corporation