



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

67690-46

Date of Issuance:

MAY 9 2008

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance: **Conditional**

Name of Pesticide Product:

SP5075 Turf Growth Regulator

Name and Address of Registrant (include ZIP Code):

SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, IN 46032-4565

Attn. Amy Dugger-Ronyak

Note: Changes in labeling, differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7) (A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA section 4.
2. You must submit two copies of a final printed label within 45 days from the date of this notice which makes the following changes:
 - A. EPA registration number must read 67690-46.

Signature of Approving Official:

Tony Kish
Product Manager Team 22
Fungicide Branch
Registration Division (7505P)

Date:

MAY 9 2008

Please make the following labeling changes:

1. On page 3 and 12, in the Hazards to Humans and Domestic Animals section, the first paragraph, second sentence should read as follows: "or on skin or clothing. Harmful if swallowed.....Avoid."
2. On page 4, delete "Enhanced" because exaggerated claims are not allowed.
3. On page 11, add "To the extent consistent with applicable law" in front of the first sentence.

The six acute studies have been reviewed and are classified as acceptable.

The proposed basic and alternate formula CSFs for the end-use product, SP5075 Turf Growth Regulator both dated October 24, 2007 are acceptable.

You must submit a one year GLP storage stability/corrosion characteristic study on or before May 6, 2009.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with comments" is enclosed for your records.

[Booklet Front Label]



ACCEPTED
with COMMENTS
In EPA Letter Dated
MAY 9 2008

SP5075
Turf Growth Regulator

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

67690-46

FOR GROWTH MANAGEMENT AND QUALITY IMPROVEMENT OF PERENNIAL COOL AND WARM SEASON TURFGRASSES ON GOLF COURSES.

Active Ingredients:

- flurprimidol: α -(1-methylethyl)- α -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol 13.26%
- trinexapac-ethyl: 4-(cyclopropyl-alpha-hydroxymethylene)-3,5-dioxo-cyclohexanecarboxylic acid ethyl ester 5.00%

Other Ingredients 81.74%

Total 100.00%

Contains 1.10 pounds of flurprimidol per gallon of product.

Contains 0.41 pounds of trinexapac-ethyl per gallon of product.

Keep Out of Reach of Children

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

* Trademark of SePRO Corporation

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

SPECIALTY CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

Refer to inside of label booklet for additional precautionary information and Directions for Use including Storage and Disposal.

Notice: Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* inside label booklet.**

For additional product information, visit us at www.sepro.com

EPA Reg. No. 67690-XX
FPL102407

EPA Est. No. XXXXX-XX-XXX
SPC-XXXXXX

SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.

Turf Growth Regulator

Net Contents _____

[Label Booklet]

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals**Keep Out of Reach of Children****WARNING / AVISO**

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. DO NOT GET IN EYES OR ON SKIN OR CLOTHING. HARMFUL IF SWALLOWED OR INHALED. AVOID BREATHING SPRAY MIST. WEAR PROTECTIVE EYEWEAR AND CHEMICAL-RESISTANT GLOVES.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Chemical-resistant gloves (such as nitrile, butyl rubber, neoprene, or barrier laminate); and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all directions for use before applying this product.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170).

The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated area without footwear until sprays have dried.

GENERAL INFORMATION

SP5075 integrates patented[†] synergy of turf growth regulator technology for use on both warm and cool-season perennial turfgrasses on golf courses. SP5075 reduces stem elongation and leaf blade length in perennial turfgrasses resulting in a more compact and dense growth form. Growth regulation from SP5075 results from suppression of the plant hormone, gibberellic acid (GA), responsible for cell elongation in most plants.

SP5075's patented site of action plant growth regulator (PGR) synergy results in enhanced growth suppression, improved turfgrass color and quality, extended spray intervals, and suppression of *Poa annua*. Plant physiological advantages to applications of SP5075 include:

1. Multiple plant sites of uptake: SP5075 is absorbed by plants via roots, stems, and leaves; and
2. Multi-site activity within gibberellic acid (GA) biosynthesis pathway: SP5075 inhibits GA production at both early and late stages in the pathway.

Turf growth regulator absorption via the roots and foliage allows for more efficient uptake by the plant ensuring an optimal amount of active ingredient is available for GA inhibition. Additionally, blocking GA biosynthesis early and late in the biological pathway regulates GA more efficiently than at a single site within this cycle.

[†]The synergy derived from the combination of Type II Class A and Type II Class B PGRs is protected by United States Patent No. 7,135,435. Additional patent rights pending.

Broadcast treatments should be made on medium to high quality turfgrass areas. An appropriate fertility program for the desired turf species should be followed in conjunction with SP5075 applications to provide the best turfgrass enhancement and reduce potential for discoloration.

Benefits of SP5075 Applications to Turfgrass:

- Shoot growth suppression of warm and cool season turfgrasses resulting in decreased mowing frequency and turfgrass clippings.
- Increased turfgrass density, wear resistance, and improved color on warm and cool season turfgrass species resulting in improved turf quality.
- Suppression of *Poa annua* (Annual Bluegrass) in cool-season turfgrasses.
- Improved water use efficiency of warm and cool season turfgrass resulting in pre-drought stress conditioning.

NOTICE TO USER: Turfgrass responses to SP5075 may vary within turfgrass species due to the large number of cultivars and varieties available. The rates recommended on this label are rate ranges and should be used only as guidelines. Neither the manufacturer nor seller has determined if SP5075 can be used safely or effectively on turfgrass species not mentioned on this label. For turfgrass species not listed on this label the user should apply SP5075 to a small test area to determine growth response and desired level of growth regulation prior to large scale applications.

General Use Precautions

- **SP5075 is not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.**
- Do not apply to bermudagrass putting greens or overseeded bermudagrass putting greens.
- Do not apply to shrubs, bedding plants, and/or food plants.
- Not recommended for turfgrasses under stress due to temperature and moisture extremes and disease, and/or insect pressures.
- Do not seed turfgrass species within 3 weeks prior to or 3 weeks after a SP5075 application.
- SP5075 should not be applied until 6 to 8 weeks after turfgrass sprigging or laying sod. Turfgrass should be well established and actively growing prior to application.
- Do not apply to saturated soils or when a significant moisture event is anticipated.
- Additional turfgrass growth regulation may occur when SP5075 is tank mixed or used in conjunction with demethylation inhibitor (DMI) or sterol inhibiting fungicides.
- Do not apply to turf used for livestock production.
- Multiple applications of SP5075 can be made yearly, but do not apply more than 350 fl oz/A of SP5075 per acre per year.
- **Chemigation:** Do not apply SP5075 through any type of irrigation system.

Application Timing:

Applications of SP5075 should be made to turfgrass actively growing. Spring applications should be made after resumption of active seasonal growth of turfgrass. The final application of the season should be timed a minimum of 4 weeks before the onset of inactive grass growth or winter dormancy. Applications to overseeded turfgrasses in dormant bermudagrass stands should be made 4 weeks prior to expected bermudagrass green-up.

Irrigation:

Rainfall or irrigation should be delayed at least 2 hours after application or until product has dried on the leaf surface but should occur within 24 hours after application. Avoid mowing turfgrass areas treated with SP5075 until after rainfall or irrigation occurs.

Turf Color and Post-Treatment Turf Management:

Turfgrass treated with SP5075 may appear darker green in color. This color change, which appears 1 to 2 weeks after treatment, may persist an additional 3 to 6 weeks. Treated areas should be managed to encourage the growth of a healthy vigorous turf.

APPLICATION DIRECTIONS**Mixing Directions:**

Add SP5075 to a spray tank half filled with clean water while agitating. Allow sufficient mixing time to ensure consistent mixing of SP5075. Finish filling the spray tank. Continue agitation throughout the spraying operation to ensure uniform application. SP5075 should be applied using a boom-type sprayer with bypass and/or mechanical agitation calibrated to deliver 20 to 100 gallons/acre of spray solution (0.5 to 2.5 gallons/1000 ft²). In-line strainers and nozzle screens should be 50 mesh or larger. The use of a coloring agent or foam to mark areas already sprayed is suggested for uniform application without skips and overlaps. Performance may be improved with the addition of a readily available nitrogen (N) source at 0.125 to 0.5 lbs N/1000 ft² or iron (Fe) at suggested label rates to the spray mix.

SP5075 + Tank Mixtures:

SP5075 can be tank mixed and is compatible with most commonly-used pesticides and foliar nutrient products. However, compatibility of SP5075 with tank mix partners should be tested before use.

NOTE: It is recommended that the compatibility of SP5075 in any tank-mix combination be tested before use. To determine the physical compatibility with other products, use a jar test as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure sequence for adding required ingredients to the spray tank.

Read and follow all label directions for each tank-mix product.

SP5075 FOR GROWTH REDUCTION OF PERENNIAL TURFGRASS SPECIES

A multiple application program using SP5075 provides growth reduction of perennial turfgrass species resulting in decreased mowing frequency and turfgrass clippings. For cool-season grasses, initial applications should begin in early spring following resumption of active growth. For warm-season grasses, initial applications should

* Trademark of SePRO Corporation

begin when the grass has completely recovered from winter dormancy and is growing vigorously. For both warm and cool-season grasses, applications should be discontinued a minimum of 4 weeks before the onset of inactive grass growth or winter dormancy. Use lower rate range in early spring and late fall applications to avoid excessive growth regulation. Refer to Table 1 for rate recommendations for growth regulation of perennial turfgrass species.

TABLE 1			
RECOMMENDED RATE RANGES FOR GROWTH REGULATION OF PERENNIAL TURFGRASS SPECIES WITH SP5075 USING A MULTIPLE APPLICATION PROGRAM.			
Turfgrass Species	Early Spring/Late Fall Applications	Repeat Applications	
	Rate of SP5075/A (fl oz/A)^{2,3}	Rate of SP5075/A (fl oz/A)²	Treatment Interval
Cool-Season Turfgrasses			
Bentgrass (golf course fairway)	10 to 15 fl oz/A	10 to 20 fl oz/A	2 to 6 weeks
Bentgrass putting greens	5 to 8 fl oz/A	5 to 10 fl oz/A	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass Mixture	15 to 22 fl oz/A	15 to 30 fl oz/A	2 to 6 weeks
Perennial Ryegrass ¹	15 to 22 fl oz/A	15 to 30 fl oz/A	2 to 6 weeks
Warm-Season Turfgrasses			
Seashore Paspalum	10 to 15 fl oz/A	10 to 20 fl oz/A	2 to 6 weeks
Tifway Bermudagrass	10 to 15 fl oz/A	10 to 20 fl oz/A	2 to 6 weeks
TifSport Bermudagrass	8 to 12 fl oz/A	8 to 16 fl oz/A	2 to 6 weeks
Zoysiagrass	8 to 12 fl oz/A	8 to 16 fl oz/A	2 to 6 weeks

¹ For perennial ryegrass overseeded fairways, applications should be delayed until perennial ryegrass is well established (3 to 4 weeks after germination). Final spring application should be made a minimum of 4 weeks prior to expected bermudagrass green-up.

² Higher rate ranges should be utilized for perennial turfgrass species maintained at higher mowing heights.

³ Use lower rate ranges during early spring or late fall when turfgrass growth and vigor are reduced.

SP5075 FOR POA ANNUA (ANNUAL BLUEGRASS) SUPPRESSION IN COOL SEASON TURF

A growth regulation program using SP5075 suppresses *Poa annua* in cool season turfgrasses. Research has demonstrated that programmed applications of SP5075 are effective at suppressing *Poa annua* populations in cool-season turfgrasses when compared to formulations of trinexapac-ethyl which have shown to have minimal impact on reducing *Poa annua* populations. This program provides a gradual increase in cool-season turfgrass populations while reducing *Poa annua* populations over one to several growing seasons. Temporary initial discoloration of *Poa annua* is possible with aggressive rates of SP5075, especially during temperature extremes. Overseeding of perennial turfgrass species should occur a minimum of 3 weeks following the last application of SP5075 for the season. For more aggressive *Poa annua* conversion,

users should consider programmed applications of Cutless 50W at labeled rates and timings for the desired turfgrass species. Refer to Table 2 for rate recommendations for *Poa annua* suppression in cool-season turfgrasses using SP5075.

- **Bentgrass (golf course fairway)**

Apply SP5075 for *Poa annua* suppression in fairway height bentgrass at a rate of 8 to 15 fl oz/A in early spring following resumption of active growth of the grass. Repeat applications of 8 to 20 fl oz/A of SP5075 should be made at 3 to 6 week intervals until late summer or early fall. Reduce rates of SP5075 should be considered in bentgrass fairways with high populations of *Poa annua* or when temporary *Poa annua* discoloration cannot be tolerated. Normal management practices such as fertilization, aeration and overseeding will encourage growth of bentgrass.

- **Bentgrass Putting Greens**

Annual turfgrass species, such as *Poa annua* are more strongly regulated by applications of SP5075 often resulting in transitory yellowing or temporary discoloration. Therefore, reduced rates of SP5075 should be used on bentgrass putting greens with high populations of *Poa annua* (>50%) or when temporary *Poa annua* discoloration cannot be tolerated. Over time, programmed applications of SP5075 throughout the growing season will suppress *Poa annua* populations, resulting in increased populations of bentgrass. Follow normal management practices such as fertilization, aeration and overseeding that encourage growth of bentgrass. Do not overseed within 3 weeks following the final application of SP5075 in the season. Use of SP5075 on bentgrass greens may increase putting speed without reducing the height of cut.

- **Bentgrass Greens with less than 50% *Poa annua*:** Apply SP5075 to bentgrass as part of an overall greens management program both for growth regulation and *Poa annua* suppression. An initial application of SP5075 at 5 to 10 fl oz/A should be made in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Repeat applications of 5 to 15 fl oz/A of SP5075 should be made at 2 to 4 week intervals through early fall.
- **Bentgrass Greens with more than 50% *Poa annua*:** Apply SP5075 to bentgrass as part of an overall greens management program both for growth regulation and *Poa annua* suppression. An initial application of SP5075 at 5 fl oz/A should be made in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Repeat applications of 5 to 10 fl oz/A of SP5075 should be made at 2 to 4 week intervals through early fall.

- **Kentucky bluegrass, Perennial ryegrass Mixtures**

Apply SP5075 for *Poa annua* suppression in fairway height Kentucky bluegrass and perennial ryegrass mixtures at a rate of 15 to 25 fl oz/A in early spring following resumption of active growth of the grass. Repeat applications of 15 to 30 fl oz/A of SP5075 should be made at 2 to 6 week intervals until late summer or early fall. Reduce rates of SP5075 should be considered in Kentucky bluegrass, perennial ryegrass mixtures with high populations of *Poa annua* or when temporary *Poa annua* discoloration cannot be tolerated. Normal management practices such as fertilization, aeration and overseeding will encourage growth of bluegrass and/or ryegrass.

- **Perennial ryegrass**

Apply SP5075 for *Poa annua* suppression in fairway height perennial ryegrass at a rate of 15 to 22 fl oz/A in early spring following resumption of active growth of the grass. Repeat applications of 15 to 30 fl oz/A of SP5075 should be made at 2 to 6 week intervals until late summer or early fall. Reduce rates of SP5075 should be considered in perennial ryegrass with high populations of *Poa annua* or when temporary *Poa annua* discoloration cannot be tolerated. For bermudagrass fairways overseeded with perennial ryegrass applications of SP5075 should be delayed until perennial ryegrass is well established (3 to 4 weeks after germination). Final spring application should be made a minimum of 4 weeks prior to expected bermudagrass green-up. Normal management practices such as fertilization, aeration and overseeding will encourage growth of perennial ryegrass.

Normal management practices such as fertilization, aeration and overseeding will encourage growth of bluegrass and/or ryegrass.

TABLE 2				
RECOMMENDED RATE RANGES FOR POA ANNUA SUPPRESSION IN COOL-SEASON PERENNIAL TURFGRASS WITH SP5075 USING A MULTIPLE APPLICATION PROGRAM.				
Turfgrass Species	% <i>Poa annua</i>	Initial spring application	Repeat applications ¹	
		Rate of SP5075/A (fl oz/A)	Rate of SP5075/A (fl oz/A)	Treatment Interval
Bentgrass (golf course fairway)	0-80%	8 to 15 fl oz/A	8 to 20 fl oz/A	2 to 6 weeks
Bentgrass Putting Greens	Less than 50%	5 to 10 fl oz/A	5 to 15 fl oz/A	2 to 4 weeks
Bentgrass Putting Greens	More than 50%	5 fl oz/A	5 to 10 fl oz/A	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass	0-80%	15 to 25 fl oz/A	15 to 30 fl oz/A	2 to 6 weeks

* Trademark of SePRO Corporation

Fairways Mixture				
Perennial Ryegrass ²	0-80%	15 to 30 fl oz/A	15 to 30 fl oz/A	2 to 6 weeks

¹ Apply in early spring following resumption of active growth of the grass. Fall applications should be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

² For perennial ryegrass overseeded fairways, applications should be delayed until perennial ryegrass is well established (3 to 4 weeks after germination). Final spring application should be made a minimum of 4 weeks prior to expected bermudagrass green-up.

STORAGE AND DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used on site according to use directions or disposed of at an approved waste disposal facility.

Container Disposal: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent consistent with applicable law, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is notified in writing within twenty-one (21) days from the date of application of such losses or damages. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies can not be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitations of Remedies in any manner.

© Copyright 2007 by SePRO Corporation

[Base Label]



SP5075
Turf Growth Regulator

FOR GROWTH MANAGEMENT AND QUALITY IMPROVEMENT OF PERENNIAL COOL AND WARM SEASON TURFGRASSES ON GOLF COURSES.

Active Ingredients:

- flurprimidol: α -(1-methylethyl)- α -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol 13.35%
- trinexapac-ethyl: 4-(cyclopropyl-alpha-hydroxymethylene)-3,5-dioxo-cyclohexanecarboxylic acid ethyl ester 5.12%

Other Ingredients 81.53%
Total 100%

Contains 1.10 pounds of flurprimidol per gallon of product.
 Contains 0.42 pounds of trinexapac-ethyl per gallon of product.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING / AVISO

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. DO NOT GET IN EYES OR ON SKIN OR CLOTHING. HARMFUL IF SWALLOWED OR INHALED. AVOID BREATHING SPRAY MIST. WEAR PROTECTIVE EYEWEAR AND CHEMICAL-RESISTANT GLOVES.

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

* Trademark of SePRO Corporation

If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053.</p>	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

SP5075 is not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

STORAGE AND DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used on site according to use directions or disposed of at an approved waste disposal facility.

Container Disposal: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SPECIALTY CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

Refer to inside of label booklet for additional precautionary information and Directions for Use.

Notice: Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* inside label booklet.**

For additional product information, visit us at www.sepro.com

EPA Reg. No. 67690-XX
FPL102407

EPA Est. No. XXXXX-XX-XXX
SPC-XXXXXX

SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.

Turf Growth Regulator

Net Contents _____