

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460

EPA Reg. Number: Da	ite of Issuance:
---------------------	------------------

67690-49

2 2 APR 2009

Term of Issuance:

conditional

Name of Pesticide Product:

SP1304

NOTICE OF PESTICIDE:

<u>x</u> Registration <u>Reregistration</u> (under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Sepro Corporation 11550 N. Meridian Street Suite 600 Carmel, IN 46032

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2. Submit the data listed below:
 - a. Within one year of the date on this registration notice, submit Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies.
- 3. Make the following label changes:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 67690-49."

Signature of Approving Official:

oanne J. Miller

Date:

Joanne I. Miller

Product Manager 23

Herbicide Branch

Registration Division (7505P)

22 APR 2000

EPA Form 8570-6

Pertains to Professional Uses Label:

- b. Add Note to Physician including the following suggested types of information that may be included, if applicable: "- technical information on symptomatology; use of supportive treatments to maintain life functions; medicine that will counteract the specific physiological effects of the pesticide; company telephone number to specific medical personnel who can provide specialized medical advice."
- c. Immediately following PPE and User Safety Requirements, add "Engineering Controls: Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticide [40 CFR 170.240(d)(6)]".
- d. Under the Agricultural Use Requirement Box add the statement "Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours."
- e. In the Agricultural Use Requirement Box add "PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves made of any water-proof material, shoes plus socks, protective eyewear."
- f. Place appropriate statement in the Directions for Use, under General Precautions and Restrictions, "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application."
- g. Change "Drift Management" to Spray Drift Management. Change the following to read "A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product."

Droplet Size

"When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles."

Wind Speed

"Do not apply at wind speeds greater than 15mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field."

Pertains to Consumer/Homeowner Uses Label:

- h. Under Hazard to Humans and Domestic Animals (for both labels) add the statement "Causes substantial, but temporary eye injury. Harmful if swallowed. Avoid contact with contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes, and gloves." Delete the language under "Concentrated Formulation".
- i. Under Important General Use Information and Precautions, add the statement "Do not allow people or pets to enter the treated area until dusts have settled."
- j. Under Storage and Disposal, change the bullets to read "Pesticide Storage" and "Pesticide Disposal". Move the first statement "Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available." under "Pesticide Disposal" bullet.
- k. Change Environmental Hazard statement for Residential Use label to read "This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater."

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)

SP1304

Aquatic Herbicide

EPA Reg. No. 67690-XX

Master Label (main brand name) – split label containing professional and consumer uses.

Registration Notes: Draft label submitted to EPA on 10/10/2008.

Table of Contents for Master Label

1.	Main Label - Professional Uses	Pages 2 - 13
11.	Main Label – Consumer/Homeowner Uses	Pages 14 - 23

[Editor's note: This portion of the master label contains Professional Uses.]

[Base label for Non-dilutable, Non-Rigid containers, any size]

ACCEPTED with COMMENTS in EPA Letter Dated

2 2 APR 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

67690-49

Sepro

SP1304

AQUATIC HERBICIDE

A SELECTIVE HERBICIDE FOR CONTROLLING CERTAIN UNWANTED AQUATIC PLANTS

Active Ingredient:

2,4-dichlorophenoxyacetic acid, dimethylamine salt:	20.0%
Other Ingredients	80.0%
Total	
2.4-d acid equivalence (a.e.): 16.5%	

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

CONCENTRATED FORMULATION

May cause substantial but temporary eye injury. May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

	\cdot		
TENERAL PROPERTY	FIRST AID		
If in eyes	 Hold eye open and rinse slowly and gently with water for 15 - 20 		
	minutes. Remove contact lenses, if present, after the first 5 minutes,		
	then continue rinsing eye.		
	Call a poison control center or doctor for treatment advice:		
If	Call a poison control center or doctor immediately for treatment advice.		
swallowed	Have person sip a glass of water if able to swallow.		
	Do not induce vomiting unless told to do so by a poison control conter-		
	or doctor.		
	Do not give anything by mouth to an unconscious person.		
If inhaled	Move person to fresh air.		
	If person is not breathing, call 911 or an ambulance, then give artificial		
	respiration, preferably mouth-to-mouth if possible.		
	 Call a poison control center or doctor for further treatment advice. 		

^{*}Trademark of SePRO Corporation

If on skin
or`
clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

ENVIRONMENTAL HAZARDS

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, in quiescent waters, apply the product to areas separated by untreated sections that can be treated after vegetation in treated areas has disintegrated. During the growing season, weeds decompose in a 2 to 4 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Nonrefillable Container Disposal (non-rigid, any size): Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refer to label booklet for additional precautionary information and directions for use.

NOTICE: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

Aquatic Herbicide	Net Contents:		
SePRO Corporation 11550 N. Meri	dian Street, Suite 600, Carmel, IN	46032 U.	S.A.
EPA Reg. No. 67690-XX FPL100908	EPA Est. No	· · · · · · · · · · · · · · · · · · ·	
For additional information on our pro	ducts, please visit www.sepro.com.		i f

[Front label for Non-dilutable, Non-Rigid containers, any size]



SP1304

AQUATIC HERBICIDE

A SELECTIVE HERBICIDE FOR CONTROLLING CERTAIN UNWANTED AQUATIC PLANTS

LANIO	
Active Ingredient: 2,4-dichlorophenoxyacetic acid, dimethylami Other Ingredients Total	ne salt:20.0% 80.0% 100.00%
2,4-d acid equivalence (a.e.): 16.5%	
KEEP OUT OF REACH OF CHILD WARNING / AVISO Si usted no entiende la etiqueta, busque a a (If you do not understand the label, find som	lguien para que se la explique a usted en detalle.
CONCENTRATED FORMULATION	
Refer to inside of label booklet for additional for use including storage and disposal.	onal precautionary information and directions
	ccording to label directions. Before using this ent Risks of Use, and Limitation of Remedies aptable, return at once unopened.
For additional information on our products, p	lease visit www.sepro.com.
EPA Reg. No. 67690-XX FPL100908	EPA Est. No
SePRO Corporation 11550 N. Meridian St	
Aquatic Herbicide	Net Contents:

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

CONCENTRATED FORMULATION

May cause substantial but temporary eye injury. May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

	FIRSTAID
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	 Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
If inhaled	 Do not give anything by mouth to an unconscious person. Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeve shirt and long pants;
- · Shoes and socks; and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

^{*}Trademark of SePRO Corporation

26

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, in quiescent waters, apply the product to areas separated by untreated sections that can be treated after vegetation in treated areas has disintegrated. During the growing season, weeds decompose in a 2 to 4 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

GENERAL INFORMATION

Use SP1304 herbicide for control of certain aquatic weeds in: ponds; lakes; reservoirs; marshes; bayous drainage ditches; non-irrigation canals; and rivers and streams that are quiescent or slow-flowing. This product is designed to selectively control the weeds listed on this label. While certain other weeds may be suppressed, control may be incomplete. Reduced control may occur in lakes or ponds where water replacement comes from bottom springs.

SP1304 is formulated on biodegradable granules that, when applied to water bodies, immediately delivers the herbicide down to the critical areas for controlling target weeds.

When applying SP1304 follow all applicable use directions, precautions and limitations. All SP1304 concentrations referred to in this label are based on acid equivalence (a.e.).

Obtain Required Permits: Consult with the State or local agency with primary responsibility for pesticide regulation before applying to public waters, to determine if a permit or public notification is required.

Recreational Use of Water in Treatment Area: There are no restrictions on the use, of treated water for recreational purposes, including swimming, fishing and domestic purposes.

Livestock Use of Water from Treatment Area: There are no restrictions on consumption of treated water for potable use by livestock, pets, or other animals.

^{*}Trademark of SePRO Corporation

GENERAL PRECAUTIONS AND RESTRICTIONS

- <u>Do not</u> apply SP1304 in a way that will contact workers or other persons, either directly or through drift; only protected handlers may be in close proximity to the mixing area or application equipment while in use; otherwise, do not enter or allow people (or pets) to enter the treatment area while SP1304 is being applied. For requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.
- Chemigation: Do not apply SP1304 through any type of irrigation system.
- Applications to a target area are limited to two (2) per season.
- Apply a maximum of 65.4 pounds of SP1304 (10.8 lbs a.e.)/acre-foot per application.
 <u>Do not exceed 4.0 ppm during any single application.</u>
- <u>Do not</u> apply within 21 days of previous application except when conducting split treatments. Split treatments, over relatively short periods of time (e.g. 1 to 4 days), may be effective in some areas to maintain adequate exposure with target plants, such as small sites or sites with higher dilution potential.
- When treating moving bodies of water, applications must be made while traveling upstream to prevent concentration of herbicide downstream from the application.

APPLICATION TO WATERS USED FOR IRRIGATION Irrigation Restrictions

- <u>Do not</u> use treated water for irrigating greenhouse or nursery plants unless 2,4-D residues are confirmed to be less than 1 ppb by laboratory analysis.
- <u>Do not</u> use water treated with SP1304 for hydroponic farming.
- <u>Do not</u> apply SP1304 directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers, or other desirable susceptible broadleaf plants, and <u>do not</u> permit dust to drift into these areas.
- This label describes both required and recommended uses of a chemical analysis for the active ingredient 2,4-D. SePRO Corporation recommends the use of an Enzyme-Linked Immunoassay (ELISA) test for the determination of SP1304 concentration in water. Contact SePRO Corporation for the incorporation of this analysis into your treatment program. Other proven chemical analysis for the active ingredient may also be used. The ELISA analysis is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredients in the water.
- If SP1304 treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with 2,4-D such as pastures, turf or established grasses, cereal grains, the treated water may be used to irrigate and/or mix sprays for these sites at any time during and after the application.
- Due to potential phytotoxicity and/or residue considerations, the following restrictions are applicable to other uses of irrigation water:
 - o If treated water is intended to be used to irrigate or mix sprays for crops not labeled for direct treatment with 2,4-D, the water must not be used unless **one** of the following restrictions has been observed:
 - A setback distance described in Table 1 (DRINKING WATER SETBACK CONTINUES) was used for the application;
 - A waiting period of 21 days from the time of application has elapsed; or
 - An approved assay indicates that the 2,4-D concentration is 100 ppb or less at the water intake. See Table 2 (SAMPLING FOR DRINKING WATER ANALYSIS) for the recommended waiting periods after application but before taking the initial water sample at water intake.

o If treated water is intended to be used to irrigate non-crop areas not labeled for direct treatment with 2,4-D (e.g. landscape ornamentals) or for other irrigation uses not described, consult with SePRO Corporation prior to commencing irrigation if 2,4-D concentrations are greater than 100 ppb.

APPLICATIONS TO POTABLE WATER SOURCES Potable Water Restrictions

- The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of 2,4-D in water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that 2,4-D concentrations in potable water do not exceed 70 ppb, respectively, at the time of consumption.
- The drinking water setback distances from functioning potable water intakes are provided in Table 1.

TABLES.						
DRINKING WATER SETBACK DISTANCES						
Application concentration and minimum setback distance (ft) from functioning potable						
	water intake					
≤1 ppm [†] 1.1 to 2.0 ppm [†] 2.1 to 3.0 ppm [†] 3.1 to 4.0 ppm [†]						
600	1,200	1,800	2,400			

[†] ppm acid equivalent target water concentration

- Following each application of SP1304, treated water must not be used for potable water unless <u>one</u> of the following restrictions has been observed:
 - o A setback distance described in Table 1 was used for the application;
 - A waiting period of at least 21 days from the time of application has elapsed; or
 - O An approved assay indicates that the 2,4-D concentration is 70 ppb or less at the water intake. Sampling for drinking water analysis should occur no sooner than stated in Table 2. NOTE: Sampling for drinking water analysis should occur no sooner than 3 days after SP1304 application. Analysis of 2,4-D in drinking water samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

	TABLE2:					
SAMPLING FOR DRINKING WATER ANALYSIS						
Minimum days after application before initial water sampling at the functioning potable water intake						
≤1 ppm ^{††} 1.1 to 3.0 ppm ^{††} 3.1 50 4.0 ppm ^{††}						
5	10	14				

These are general guidelines; the amount of time required for residues to reach concentrations acceptable for drinking or irrigation will depend on the total acres treated relative to water body size, application rates, water exchange rates, weed density, and various other factors. Consult a SePRO Aquatic Specialist for site specific recommendations.

 If no setback distance from Table 1 is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an

^{††} ppm acid equivalent target water concentration

^{*}Trademark of SePRO Corporation

76

advisory to shut off all potable water intakes inside the setback zone prior to SP1304 application. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water. The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

Example:

- Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting should include the day and time of application. Posting may be removed if analysis of a sample collected at the intake no sooner than stated in Table 2 shows that the 2,4-D concentration is 70 ppb or less, or after 21 days following application, whichever occurs first.
 - Text of notification: Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water unless water at functioning drinking water intakes is tested no sooner than [insert days from Table 2] and is demonstrated by assay to contain no more than 70 ppb 2,4-D. Application Date: ______ Time: _____.
- NOTE: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.
- Drinking water setback distances do not apply to terrestrial applications of 2,4-D adjacent to water bodies with potable water intakes.

DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity), method of application (e.g., ground, aerial, airblast), can influence the potential for drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Applying SP1304 through an enclosed eductor or slurry injection system via a continuous stream of water and/or injected under the water surface further minimizes drift potential.

Wind Speed

Do not apply at wind speeds greater than 15 mph. SP1304 may be applied using an enclosed eductor or slurry injection system. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) near enough to be injured.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below application height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants: Do not apply under circumstances where drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers,

^{*}Trademark of SePRO Corporation

tomatoes, beans, and other vegetables, or tobacco. Small amounts of pesticide drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements: Applicators must follow all state and local pesticide drift requirements regarding application of triclopyr or 2,4-D herbicides to aquatic sites. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Aerial applications

- Apply SP1304 at the lowest height consistent with efficacy and flight safety. Do not apply
 at a height greater than 10 feet above the water surface or plant canopy unless a greater
 height is required for aircraft safety.
- When applications are made with a crosswind, the swath will be displaced downwind. The
 applicator must compensate for this by adjusting the path of the aircraft upwind.

APPLICATION INFORMATION

When to Apply:

For best results, apply SP1304 in the spring and early summer, during the time weeds start to grow. If desired, this timing can be checked by sampling the lake bottom in areas heavily infested with weeds the year before. If treatments are delayed until weeds form a dense mat or reach the surface, two treatments may be necessary. Make the second treatment when weeds show signs of recovery. Treatments made after September may be less effective depending upon water temperatures, region of the country, and weed growth. Occasionally, a second application will be necessary if heavy re-growth occurs or weeds re-infest from untreated areas.

How to Apply:

For Large Areas:

Use a fertilizer spreader or mechanical seeder such as the Gerber or Gandy or other equipment capable of uniformly applying this product. Before applying any chemical, calibrate your method of application to be sure of spreading the proper amount. When using boats and power equipment, you must determine the proper combination of (1) boat speed (2) rate of delivery from the spreader, and (3) width of swath covered by the granties.

For Small Areas: (Around Docks or Isolated Patches of Weeds)
Use a portable spreader such as the Cyclone seeder or other equipment capable of uniformly applying this product. Estimate or measure out the area you want to treat. Weigh out the amount of material needed and spread uniformly over the area. More uniform coverage is obtained by dividing the required amount in two and covering the area twice, applying the second half at right angles to the first.

Use the following formula to calibrate your spreader's delivery in pounds of SP1304 per minute:

 $\frac{\text{Miles per hour x spreader swath width (ft) x pounds per acre}}{495} = \textbf{Pounds per Minute}$

^{*}Trademark of SePRO Corporation

Example Calculation: To apply 100 pounds of SP1304 per acre using a spreader that covers a 20 foot swath from a boat traveling at 4 miles per hour, set the spreader to deliver 16.1 pounds of SP1304 granules per minute:

 $\frac{4 \text{ mph x 20 feet x 100 Lbs./A}}{495}$ = 16.1 lbs/min.

Amounts to Use:

Rates of application vary with susceptibility of weed species to SP1304 (refer to Table 3, *AQUATIC PLANT SUSCEPTIBILITY TO SP1304*), density of weed mass at time of treatment, stage of growth, water depth, and rate of water exchange through the treated area. Consult with a SePRO Aquatic Specialist for site specific recommendations. Rates should be selected according to rates found in Table 4 (*POUNDS OF SP1304 TO APPLY FOR TARGET CONCENTRATION OF 2 AND 4 PPM 2.4-D*).

AQUATIC PLANT SUSCEPTIBILI PLANT SPECIES	SP1304 PPM [†]
usceptible Weeds	
watermilfoil (Myriophyllum spp.)	up to 1.0 to 4.0
bladderwort (Utricularia spp.) coontail ^{††} (Ceratophyllym Demersum) water chestnut (Trapa natans) water stargrass ^{††} (Heteranthera dubia) watershield (Brasenia spp.) white water lily (Nymphaea spp.) yellow water lily (Nuphar spp.) or spatterdock ^{††}	up to 2.0 to 4.0

[†]Lower concentrations may be effective under certain conditions, such as when treating larger/continuous areas or areas with less water exchange, etc.

^{††} Repeat applications may be necessary.

POUNDS OF SP1304 TO APPLY FOR A TARGET CONGENTRATION OF 2 AND 4 PPM 2,4-D [†]				
	Average Depth	For typical	For difficult.	
Surface Area	(feet)	conditions – 2 ppm	conditions – 4 ppm	
		(pounds)	(pounds)	
	1	32.7	(,,,) 65.4	
	2	65.4		
1 acre	3	98.2	196.3 '' ' '	
	4	130.9	261.8	
	5	163.6	327.2	

Higher rates in the rate range are recommended: in areas of greater water exchange, when treating more mature plants, when targeting more difficult to control aquatic species, and when treating small areas in larger bodies of water (spot treatments). Lower concentrations are generally used when: conducting early season large-scale

^{*}Trademark of SePRO Corporation

treatments; and treating larger areas, more immature plants, and in areas with less potential for rapid water exchange. Some areas may require a repeat application to control re-growth.

NOTE: The following formula can be used to calculate applications rates based on depths exceeding 4 feet deep or when using a concentration not in the table above:

Pounds of SP1304 per Acre = average depth x target ppm x 16.36

Example Calculation:

6 foot average depth x 2.25 ppm x 16.36 = 221 pounds of SP1304 per acre

NOTE: apply a maximum of 65.4 pounds of SP1304 (10.8 lbs a.e.)/acre-foot per application. Do not exceed 4.0 ppm during any single application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Nonrefillable Container Disposal (non-rigid, any size): Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plantinjury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. To the extent permitted by applicable law in no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* above and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or *Limitations of Remedies* in any manner.

©Copyright ____ SePRO Corporation

[Editor's note: This portion of the master label contains Consumer/Homeowner Uses.]

[Front label for Household/Residential container]



SP1304

AQUATIC HERBICIDE

READY-TO-USE

A SELECTIVE HERBICIDE FOR CONTROLLING CERTAIN UNWANTED AQUATIC PLANTS

Active Ingredient:

2,4-dichlorophenoxyacetic acid, dimethylamine salt:	20.0%
Other Ingredients	<u>80.0%</u>
Total	100.00%
2.4-d acid equivalence (a.e.): 16.5%	

KEEP OUT OF REACH OF CHILDREN WARNING

Specialty Chemical: Do not ship or store with food, feeds, drugs or clothing.

See back panel booklet for additional precautionary statements, directions for use and warranty information.

Net []

[Editorial Note: Bracketed [text] indicates marketing claims; text in curly brackets {text} show place holders for graphics.]



SP1304

AQUATIC HERBICIDE

READY-TO-USE

PRODUCT FACTS


This product will provide herbicidal control of certain unwanted aquatic plants species.

AQUATIC WEEDS CONTROLLED

Controls Various Types of Nuisance Aquatic Weeds







Milfoil

ontail Li

(See complete list under the directions for use section of this label)

How SP1304 Works

SP1304 is applied directly to water in a pond or water front shoreline. The active ingredient, 2,4-D, is quickly absorbed by aquatic plants. SP1304 is formulated on biodegradable granules that that when applied to water bodies immediately sink and deliver the product to the critical areas for controlling target weeds. During the growing season, typically weeds decompose in a 2 to 4 week period following treatment.

Where to Apply Apply SP1304 herbicide to ponds and water front shorelines.	
Questions, Comments or Information {graphic of phone} Call	
{graphic of internet/computer} www	
	•

Open booklet for details. Made in U.S.A.

SePRO Corporation 11550 N. Meridian Street, Suite 600, Carmel, IN 46032 U.S.A

EPA Reg. No. 67690-XX FPL100908 EPA Est. No. _____

© Copyright _____ SePRO Corporation.

[Label booklet text for Household/Residential containers]



SP1304

AQUATIC HERBICIDE

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN WARNING

May cause substantial but temporary eye injury. May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

	FIRST AID
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
•	duct container or label with you when calling a poison control certer or doctor,

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

ENVIRONMENTAL HAZARDS

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. Which treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, in quiescent waters, apply the product to areas separated by untreated sections that can be treated after vegetation in treated areas has disintegrated. During the growing season, weeds decompose in a 2 to 4 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into

^{*}Trademark of SePRO Corporation

untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

GENERAL INFORMATION

Use SP1304 herbicide for control of certain aquatic weeds in ponds and on water front shorelines. This product is designed to selectively control the weeds listed on this label. While certain other weeds may be suppressed, control may be incomplete. Reduced control may occur in lakes or ponds where water replacement comes from bottom springs.

SP1304 is formulated on biodegradable granules that, when applied to water bodies, immediately delivers the herbicide down to the critical areas for controlling target weeds.

When applying SP1304 follow all applicable use directions, precautions and limitations. All concentrations referred to in this label are based on acid equivalence (a.e.).

IMPORTANT GENERAL USE INFORMATION AND PRECAUTIONS

- There are no restrictions on fishing, swimming, recreational use, or consumption of treated water for potable use by livestock, pet, or other animals following application of SP1304.
- <u>Do not</u> allow people or pets to enter the treatment area while SP1304 is being applied.
- <u>Do not</u> apply this product through any type of irrigation system.
- Applications to target areas are limited to two (2) per season.
- Apply a maximum of 65.4 pounds of SP1304 (10.8 lbs a.e.)/acre-foot per application. <u>Do not</u> exceed 4.0 ppm during any single application.
- <u>Do not</u> apply within 21 days of previous application except when conducting split treatments. Split treatments, over relatively short periods of time (e.g. 1 to 4 days), may be effective in some areas to maintain adequate exposure with target plants, such as small sites or sites with higher dilution potential.
- Obtain Required Permits: Consult with appropriate state or local water authorities before applying SP1304. If the pond/lake is not privately owned by you be sure to check with state or local water authorities who have regulatory authority over public water as permits may be required by state or local public agencies.

APPLICATION TO WATERS USED FOR IRRIGATION Irrigation Restrictions

- Do not use treated water for irrigating greenhouse or nursery plants unless 2,4-D residues are confirmed to be less than 1 ppb by laboratory analysis.
- <u>Do not</u> use water treated with SP1304 for hydroponic farming.
- <u>Do not</u> apply SP1304 directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers, or other desirable susceptible broadleaf plants, and <u>do not</u> permit dust to drift into these areas.
- This label describes both required and recommended uses of a chemical analysis for the active ingredient 2,4-D. SePRO Corporation recommends the use of an Enzyme-Linked Immunoassay (ELISA) test for the determination of SP1304 concentration in water. Contact SePRO Corporation for the incorporation of this analysis into your treatment program. Other proven chemical analysis for the active ingredient may also be used. The

^{*}Trademark of SePRO Corporation

ELISA analysis is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredients in the water.

- If SP1304 treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with 2,4-D such as pastures, turf or established grasses, cereal grains, the treated water may be used to irrigate and/or mix sprays for these sites at any time during and after the application.
- Due to potential phytotoxicity and/or residue considerations, the following restrictions are applicable to other uses of irrigation water:
 - If treated water is intended to be used to irrigate or mix sprays for crops not labeled for direct treatment with 2,4-D, the water must not be used unless <u>one</u> of the following restrictions has been observed:
 - A setback distance described in Table 1 (DRINKING WATER SETBACK DISTANCES) was used for the application;
 - A waiting period of 21 days from the time of application has elapsed; or
 - An approved assay indicates that the 2,4-D concentration is 100 ppb or less at the water intake. See Table 2 (SAMPLING FOR DRINKING WATER ANALYSIS) for the recommended waiting periods after application but before taking the initial water sample at water intake.
 - o If treated water is intended to be used to irrigate non-crop areas not labeled for direct treatment with 2,4-D (e.g. landscape ornamentals) or for other irrigation uses not described, consult with SePRO Corporation prior to commencing irrigation if 2,4-D concentrations are greater than 100 ppb.

APPLICATIONS TO POTABLE WATER SOURCES Potable Water Restrictions

- The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of 2,4-D in water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that 2,4-D concentrations in potable water do not exceed 70 ppb, respectively, at the time of consumption.
- The drinking water setback distances from functioning potable water intakes are provided in Table 1.

	DRINKING WATE	R SETBACK DISTA	NCES
Application concentration and minimum setback distance (ft) from functioning potable			
water intake			
<1 ppm [†]	1.1 to 2.0 ppm [†]	2.1 to 3.0 ppm [†]	3.1 to 4.0 ppm ^f .
600	1,200	1,800	2,400

[†] ppm acid equivalent target water concentration

- Following each application of SP1304, treated water must not be used for potable water unless one of the following restrictions has been observed:
 - o A setback distance described in Table 1 was used for the application;
 - o A waiting period of at least 21 days from the time of application has elapsed; of the
 - o An approved assay indicates that the 2,4-D concentration is 70 ppb or less at the water intake. Sampling for drinking water analysis should occur no sooner than stated in Table 2. **NOTE**: Sampling for drinking water analysis should occur no sooner than 3 days after SP1304 application. Analysis of 2,4-D drinking water samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method

^{*}Trademark of SePRO Corporation

Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

The second second second second second second is a second	AND THE STATE OF T		
SAMPLIN	G FOR DRINKING WATER A	ANALYSIS T	
Minimum days after application before initial water sampling			
at the functioning potable water intake			
<1 ppm ^{††}	1.1 to 3.0 ppm ^{††}	3.1 to 4.0 ppm ^{††}	
5	10	14	

These are general guidelines; the amount of time required for residues to reach concentrations acceptable for drinking or irrigation will depend on the total acres treated relative to water body size, application rates, water exchange rates, weed density, and various other factors. Consult a SePRO Aquatic Specialist for site specific recommendations.

• If no setback distance from Table 1 is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an advisory to shut off all potable water intakes inside the setback zone prior to SP1304 application. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water. The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

o Example:

- Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting should include the day and time of application. Posting may be removed if analysis of a sample collected at the intake no sooner than stated in Table 2 shows that the 2,4-D concentration is 70 ppb or less, or after 21 days following application, whichever occurs first.
 - **Text of notification:** Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water unless water at functioning drinking water intakes is tested no sooner than [insert days from Table 2] and is demonstrated by assay to contain no more than 70 ppb 2,4-D. Application Date:
- **NOTE**: Existing potable water intakes that are no longer in use, such a sitrose replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

WHEN & HOW TO APPLY

- For best results, apply during sunny, dry, non-windy conditions.
- Apply directly into the water where weeds are present.
- For best results, apply in the spring and early summer, during the time weeds start to grow.
- If treatments are delayed until weeds form a dense mat or reach the surface, two treatments may be necessary. Make the second treatment when weeds show signs of recovery.

th ppm acid equivalent target water concentration

^{*}Trademark of SePRO Corporation

 Treatments made after September may be less effective depending upon water temperatures, region of the country, and weed growth.

- Use a portable spreader such as the Cyclone seeder or other equipment capable of uniformly applying this product. Estimate or measure out the area you want to treat.
 Weigh out the amount of material needed and spread uniformly over the area. More uniform coverage is obtained by dividing the required amount in two and covering the area twice, applying the second half at right angles to the first.
- Use the following formula to calibrate your spreader's delivery in pounds of SP1304 per minute:

Miles per hour x spreader swath width (ft) x pounds per acre = Pounds per Minute 495

Example Calculation: To apply 100 pounds of SP1304 per acre using a spreader that covers a 20 foot swath from a boat traveling at 4 miles per hour, set the spreader to deliver 16.1 pounds of SP1304 granules per minute:

$$\frac{4 \text{ mph x 20 feet x 100 Lbs./A}}{495} = 16.1 \text{ lbs/min.}$$

Rates of application vary with susceptibility of weed species to SP1304, density of weed
mass at time of treatment, stage of growth, water depth, and rate of water exchange
through the treated area. Rates should be selected according to rates found in Table 4
(POUNDS OF SP1304 TO APPLY FOR TARGET CONCENTRATION OF 2 AND 4 PPM
2.4-D).

AQUATIC PLANT SUSCEPTIBILIT PLANT SPECIES	SP1304 PPM [†]	Mary Market Market
Susceptible Weeds	O1 100411 W	
watermilfoil (Myriophyllum spp.)	up to 1.0 to 4.0	
Slightly To Moderately Susceptible Weeds		
bladderwort (Utricularia spp.)		
coontail ^{††} (Ceratophyllym Demersum) water chestnut (Trapa natans)		
water stargrass ^{††} (Heteranthera dubia)		
watershield (Brasenia spp.)	up to 2.0 to 4.0	
white water lily <i>(Nymphaea</i> spp.)		
yellow water lily (Nuphar spp.)		(
or spatterdock ^{††}		

TLower concentrations may be effective under certain conditions, such as when treating larger/continuous areas or areas with less water exchange; etc.

^{††} Repeat applications may be necessary.

	T.		GET.
POUNDS OF SP1304 TO APPLY FOR A TARGET CONCENTRATION OF 2 AND 4 PPM 2,4-D [†]			
	Average Depth (feet)	For typical	For difficult
Surface Area		conditions – 2 ppm	conditions – 4 ppm
		(pounds)	(pounds)
1 acre	1	32.7	65.4
	2	65.4	130.9
	3	98.2	196.3
	4	130.9	261.8
	5	163.6	327.2

[†] Higher rates in the rate range are recommended: in areas of greater water exchange, when treating more mature plants, when targeting more difficult to control aquatic species, and when treating small areas in larger bodies of water (spot treatments). Lower concentrations are generally used when: conducting early season large-scale treatments; and treating larger areas, more immature plants, and in areas with less potential for rapid water exchange. Some areas may require a repeat application to control re-growth.

NOTE: The following formula can be used to calculate applications rates based on depths exceeding 5 feet deep or when using a concentration not in the table above:

Pounds of SP1304 per Acre = average depth x target ppm x 16.36

Example Calculation:

6 foot average depth x 2.25 ppm x 16.36 = 221 pounds of SP1304 per acre

NOTE: apply a maximum of 65.4 pounds of SP1304 (10.8 lbs a.e.)/acre-foot per application. Do not exceed 4.0 ppm during any single application.

STORAGE AND DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Do not contaminate water, food or feed by storage or disposal.

Pesticide Handling and Storage: Store product in original container in a safe place. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

Pesticide and Container Disposal: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (3) Refund of purchase price paid by buyer or user for product bought, or
- (4) Replacement of amount of product used.

To the extent consistent with applicable law SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. To the extent permitted by applicable law in no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitations of Remedies in any manner.

©Copyright ____ SePRO Corporation

[Base label for Household/Residential containers]



SP1304

AQUATIC HERBICIDE

READY-TO-USE

SePRO FAMILY OF PRODUCTS

Visit the SePRO website **www.lakelawnandpond.com** to learn more about the SePRO family of products for the best solution to your toughest pond and lakeshore aquatic weed and algae problems.

{Insert graphic – SePRO product family photo}

STORAGE AND DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Do not contaminate water, food or feed by storage or disposal.

Pesticide Handling and Storage: Store product in original container in a safe place. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste. Pesticide and Container Disposal: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Specialty Chemical: Do not ship or store with food, feeds, drugs or clothing.

Consumer Promise: 100% Satisfaction Guaranteed.

{graphic of phor	mments or Information ne} Call net/computer} www	· 		
(grapine or inter	nercompater, www		•	
Always read and Made in U.S.A.	d follow label directions.	•		
SePRO Corpor	ation 11550 N. Meridian Street,	Suite 600, Carmel, IN	46032 Ù.S	}_, , , . A.&
EPA Reg. No. 6 FPL100908	7690-XX	EPA Est. No SPC		· (((((((((((((((((((
© Copyright	SePRO Corporation.		•	' ((() ()

^{*}Trademark of SePRO Corporation