67690-50

4/25/2012



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

> > APR 252012

Laurent Mezin SePro Corporation 11550 North Meridian St Suite 600 Carmel, IN 46032-4565

Subject: Notification per PR Notice 98-10 SP1037 EPA Reg. No. 67690-50 Application dated January 25, 2012

Dear Ms. Mezin:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 of the subject product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

The alternate brand name "Renovate Max G" has been added to your file.

If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

Kathryn Montague

Product Manager 23 Herbicide Branch Registration Division (7505P)

Please read instructions on rev	erse before completing form.		I	orm Approv	ved. C	MB No. 2070	-0060	Print Form
<b>≎EPA</b> ⊧	United States Invironmental Protectio Washington, DC 204	-			F	Registratio Amendme Other	on	OPP Identifier Number
	Applicatio	n for F	Pesticide	e - Secti	on I			
1. Company/Product Number 67690-50				duct Manag Montague	ger			None Restricted
4. Company/Product (Name) SePRO / SP1037			<b>PM#</b> 23					None Restricted
5. Name and Address of Applic SePRO Corporation 11550 N. Meridian Street, S Carmel, IN 46032 Check if this is			(b)(i), my to: EPA Re		simila	ar or identica	l in cor NOT	FIFRA Section 3(c)(3) nposition and labeling TFICATION PR 25 2012
		Sect	tion - II					
X Notification - Explain be	se to Agency letter dated	1 and Se		inal printed i gency letter Me Too" Ap hther - Explai	r dated plicati	on.	,	
1. Material This Product Will B			ion - III					
Yes* XNo	Jnit Packaging Yes No f "Yes" No. per Jnit Packaging wgt. No. per container	If "Yes Packag	e wgt	No. per container		X	nteiner Aetel Mastic Glass Paper Other (Sj	oecify)
3. Location of Net Contents Inf	formation 4. Size(s) Ret. Itainer 1 - 250 Lbs	ail Contai		5	S. Loca	tion of Label On Label On Labeling		ns panying product
6. Manner in Which Label is Af	fixed to Product Lithogr Paper Stencil	raph glued led		Other				
		Sect	ion - IV					
1. Contact Point /Complete its	ems directly below for identification	n of indivi	dual to be d	contacted, if	neces	sary, to proce	ess this	application.)
Name Laurent Mezin		Title Mgr, Reg	gulatory Af	fairs			17-580-	No. (Include Ares Code) 8286
Certification       6. Dete Application         I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.       8. Dete Application         I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.       6. Dete Application					Received			
2. Signature	- te		Julatory Af	fairs		ci.		
4. Typed Name Laurent Mezin		<b>5. Date</b> 25 Janu	ary 2012			1.6		and a second

(

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

(

Yellow - Applicant Copy

2)29



SePRO Corporation • 11550 North Meridian Street • Suite 600 • Carmel, Indiana 46032-4565 *Phone:* (317) 580-8282 *Fax:* (317) 428-4577

25 January 2012

Kathryn Montague Product Manager 23 Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

#### RE: SP1037 (EPA Reg. No. 67690-50) Notification – Alternate Brand Name: <u>Renovate Max G</u>

Dear Ms. Montague:

SePRO Corporation is submitting a notification for the Alternate Brand Name "<u>Renovate Max G</u>' for the registered product SP1037 (EPA Registration No. 67690-50). Please note this alternate brand name was originally submitted in June 2009, but may not have been processed by the Agency as it was included in a submission of final printed labels.

Please find enclosed the following information:

- Transmittal document (this letter)
- Application for Pesticide Registration, EPA Form 8570-1
- One clean and one highlighted copy of the label.

This notification is consistent with the provisions of the PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

If you have any questions, please do not hesitate to contact me at (317) 580-8286 or at LaurentM@sepro.com.

Best Regards,

- Rézin auart

Laurent Mézin Regulatory Affairs Manager

Enclosures

## SP1037 (ABN: Renovate MAX G)

**Aquatic Herbicide** 

APR 2 5 2012

4/29

EPA Reg. No. 67690-50

### FPL20120125 Highlighted

Master Label (main brand name) – split label containing professional and consumer uses. ABN: Renovate Max G

**Registration Notes:** Based on EPA approved label with FPL20110927 (notification stamped 18 October 2011).

### **Table of Contents for Master Label**

I.	Main Label – Professional Uses a. Marketed under alternate brand name Renovate MAX G	Pages 2 - 14
II.	Main Label – Consumer/Homeowner Uses	Pages 15 - 26

### General label changes:

- 1. Includes the alternate brand name 'Renovate MAX G'
- 2. Updated FPL date

[Editor's note: This portion of the master label contains Professional Uses.]

[Front label for Non-dilutable, Non-Rigid containers, any size]



## SP1037 Renovate MAX G

AQUATIC HERBICIDE

FOR CONTROL OF AQUATIC WEEDS IN PONDS; LAKES; RESERVOIRS; MARSHES; BAYOUS; DRAINAGE DITCHES; NON-IRRIGATION CANALS; AND RIVERS AND STREAMS THAT ARE QUIESCENT OR SLOW-FLOWING.

### **Active Ingredients**

triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt	4.0%
2,4-dichlorophenoxyacetic acid, dimethylamine salt:	
Other Ingredients	
Total	
Acid equivalence (a.e.): 14.4%	

### KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes substantial, but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes.

AGRICULTURAL CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

Refer to inside of label booklet for additional precautionary information and directions for use including first aid and storage and disposal.

**NOTICE:** Read the entire label. Use only according to label directions. **Before using this** product, read *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies* at end of label booklet. If terms are unacceptable, return at once unopened.

EPA Reg. No. 67690-50 FPL201<u>20125</u>10927 EPA Est. No.

SePRO Corporation 11550 N. Meridian Street, Suite 600, Carmel, IN 46032 U S.A.

Aquatic Herbicide [Label booklet text]

Net Contents: \_\_\_\_\_

### PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

Causes substantial, but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes.

### KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

	FIRST AID
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
lf swallowed	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> </ul>
	Do not induce vomiting unless told to do so by a poison control center or doctor.
5 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	<ul> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If inhaled	Move person to fresh air.
	<ul> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> </ul>
	Call a poison control center or doctor for further treatment advice.
If on skin	Take off contaminated clothing.
or	Rinse skin immediately with plenty of water for 15 - 20 minutes.
clothing	Call a poison control center or doctor for treatment advice.
or going for t	duct container or label with you when calling a poison control center or doctor, reatment. In case of emergency endangering health or the environment product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

### All loaders, applicators, and other handlers must wear:

- · Long-sleeve shirt and long pants;
- Shoes and socks; and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENGINEERING CONTROLS**

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6].

### ENVIRONMENTAL HAZARDS

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, in quiescent waters, apply the product to areas separated by untreated sections that can be treated after vegetation in treated areas has disintegrated. During the growing season, weeds decompose in a 2 to 4 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

AGRICULTURAL CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

### **GENERAL INFORMATION**

SP1037Renovate MAX G herbicide may be applied directly to water for the control of aquatic weeds. SP1037Renovate MAX G enhances target weed control, and provides selective control of many broadleaf weeds in: ponds; lakes; reservoirs; marshes; bayous; drainage ditches; non-irrigation canals; and rivers and streams that are quiescent or slow-flowing.

SP1037Renovate MAX G is formulated on biodegradable granules that, when applied to water bodies, immediately delivers SP1037Renovate MAX G down to the critical area for controlling target weeds. SP1037Renovate MAX G is quickly absorbed from the water through plant stems and foliage and from the hydrosoil by roots. Herbicidal symptoms are initially expressed 2 to 14 days following application and usually involve bending and twisting of apical sections and shoots of susceptible plants. Initial symptoms are followed by necrosis of terminal buds and above ground tissue. Generally, target plants are controlled within 2 to 4 weeks after treatment, but depending on conditions and plant species can take up to 8 weaks for complete control.

When applying SP1037Renovate MAX G follow all applicable use directions precautions and limitations. All SP1037Renovate MAX G concentrations referred to in this label are based on acid equivalence (a.e.).

**Obtain Required Permits:** Consult with the State or local agency with primary responsibility for pesticide regulation before applying to public waters to determine if a permit or public notification is required.

**Recreational Use of Water in Treatment Area:** There are no restrictions on the use of treated water for recreational purposes, including swimming, fishing and domestic purposes.

**Livestock Use of Water from Treatment Area:** There are no restrictions on consumption of treated water for potable use by livestock, pets, or other animals.

### **GENERAL USE PRECAUTIONS AND RESTRICTIONS**

- <u>Do not</u> apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not enter or allow people (or pets) to enter the treated area until dusts have settled.
- For requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.
- Chemigation: <u>Do not</u> apply <u>SP1037Renovate MAX G</u> through any type of irrigation system.
- <u>Do not</u> apply to salt water bays or estuaries.
- Applications to target areas are limited to two (2) per season.
- Apply a maximum of 93.7 pounds of <u>SP1037Renovate MAX G</u> (13.5 lbs a.e.)/acre-foot per application. <u>Do not</u> exceed 5.0 ppm during any single application.
- Do not apply within 21 days of previous application except when conducting split treatments. Split treatments, over relatively short periods of time (e.g., 1 to 4 days), may be effective in some areas to maintain adequate exposure with target plants, such as small sites or sites with higher dilution potential.
- When treating moving bodies of water, applications must be made while traveling upstream to prevent concentration of herbicide downstream from the application.

### APPLICATION TO WATERS USED FOR IRRIGATION

### **Irrigation Restrictions**

- <u>Do not</u> use treated water for irrigating greenhouse or nursery plants unless triclopyr and 2,4-D residues are confirmed to be less than 1 ppb by laboratory analysis.
- Do not use water treated with SP1037Renovate MAX G for hydroponic farming.
- <u>Do not</u> apply <u>SP1037Renovate MAX G</u> directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers, or other desirable susceptible broadleaf plants, and <u>do not</u> permit dust to drift into these areas.
- This label describes both required and recommended uses of a chemical analyses for the active ingredients, triclopyr and 2,4-D. SePRO Corporation recommends the use of an Enzyme-Linked Immunoassay (ELISA) test for the determination of SP1037Renovate MAX <u>G</u> concentration in water. Contact SePRO Corporation for the incorporation of these analyses into your treatment program. Other proven chemical analysis for the active ingredients may also be used. The ELISA analysis is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredients in the water. Both triclopyr and 2,4-D can be analyzed from a single water sample.
- If <u>SP1037Renovate MAX G</u> treated water is intended to be used only for crope or non-crop areas that are labeled for direct treatment with triclopyr and 2,4-D such as pastures; turf, or established grasses, the treated water may be used to irrigate and/or mix sprays for these sites at any time during and after application.

- Due to potential phytotoxicity and/or residue considerations, the following restrictions are applicable to other uses of irrigation water:
  - If treated water is intended to be used to irrigate or mix sprays for crops not labeled for direct treatment with triclopyr and 2,4-D, the water must not be used unless <u>one</u> of the following restrictions has been observed:
    - A waiting period of 120 days from the time of application has elapsed; or
    - An approved assay indicates that the triclopyr concentration is 1.0 ppb or less and the 2,4-D concentration is 100 ppb or less at the water intake. See Table 2 (SAMPLING FOR DRINKING WATER ANALYSES) for the recommended waiting periods after application but before taking the initial water sample at water intake.
  - If treated water is intended to be used to irrigate non-crop areas not labeled for direct treatment with triclopyr and 2,4-D (e.g., landscape ornamentals) or for other irrigation uses not described, consult with SePRO Corporation prior to commencing irrigation if triclopyr concentrations exceed 1.0 ppb and 2,4-D concentrations are greater than 100 ppb.

### APPLICATIONS TO POTABLE WATER SOURCES

### **Potable Water Restrictions**

- The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of triclopyr in water is less than 400 ppb and the concentration of 2,4-D in water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that triclopyr and 2,4-D concentrations in potable water do not exceed 400 ppb and 70 ppb, respectively, at the time of consumption.
- The drinking water setback distances from functioning potable water intakes are provided in Table 1 (DRINKING WATER SETBACK DISTANCES).

	T	ABLE 1	
	DRINKING WATE	R SETBACK DISTAN	CES
Application conce		n setback distance (ft) ater intake	from functioning potable
<1 ppm <sup>†</sup>	1.1 to 2.0 ppm <sup>†</sup>	2.1 to 3.0 ppm <sup>†</sup>	3.1 to 5 ppm <sup>†</sup>
600	1,200	1,800	2,400

<sup>†</sup> ppm acid equivalent target water concentration

- Following each application of <u>SP1037Renovate MAX G</u>, treated water must not be used for potable water unless <u>one</u> of the following restrictions has been observed:
  - A setback distance described in Table 1 was used for the application;
  - o A waiting period of at least 21 days from the time of application has elapsed; or
  - An approved assay indicates that the triclopyr concentration is 400 ppb or less and the 2,4-D concentration is 70 ppb or less at the water intake. Sampling for drinking water analyses should occur no sooner than stated in Table 2. NOTE: Sampling for drinking water analysis should occur no sooner than 3 days after SP1037Renovate MAX G application. Analysis of 2,4-D in drinking water samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for 5valuating Solid Waste SW-846.

SAMPLING	FOR DRINKING WATER AN	NALYSES T
	after application before initial e functioning potable water in	
<1 ppm <sup>††</sup>	1.1 to 3.0 ppm <sup>††</sup>	3.1 to 5.0 ppm <sup>††</sup>
5	10	14

<sup>†</sup> These are general guidelines; the amount of time required for residues to reach concentrations acceptable for drinking or irrigation will depend on the total acres treated relative to water body size, application rates, water exchange rates, weed density, and various other factors. Consult a SePRO Aquatic Specialist for site specific recommendations.

<sup>††</sup> ppm acid equivalent target water concentration

- If no setback distance from Table 1 is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an advisory to shut off all potable water intakes inside the setback zone prior to SP1037Renovate MAX G application. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water. The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.
  - Example:
    - Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting should include the day and time of application. Posting may be removed if analyses of a sample collected at the intake, no sooner than stated in Table 2, shows that the triclopyr concentration in the water is 400 ppb or less and the 2,4-D concentration is 70 ppb or less, or after 21 days following application, whichever occurs first.
      - Text of notification: Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water unless water at functioning drinking water intakes is tested no sooner than [insert days from Table 2] and is demonstrated by assay to contain no more than 400 ppb triclopyr and 70 ppb 2,4-D. Application Date: \_\_\_\_\_, Time: \_\_\_\_\_.
- NOTE: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.
- Drinking water setback distances do not apply to terrestrial applications of triclopyr or 2,4-D adjacent to water bodies with potable water intakes.

### **GRAZING AND HAYING RESTRICTIONS**

Except for lactating dairy animals, there are no grazing restrictions following application of this product.

- Grazing Lactating Dairy Animals: <u>Do not</u> allow lactating dairy animals to graze treated areas until the next growing season following application of this product.
- <u>Do not</u> harvest hay for 14 days after application.
- Grazed areas of sites may be spot treated if they comprise no more than 10% of the total grazable area.

11/29

### SLAUGHTER RESTRICTIONS

During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

### DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Applying <u>SP1037Renovate MAX G</u> through an enclosed eductor or slurry injection system via a continuous stream of water and/or injected under the water surface further minimizes drift potential.

#### Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) near enough to be injured.

#### **Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below application height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**Susceptible Plants:** Do not apply under circumstances where drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of pesticide drift that might not be visible may injure susceptible broadleaf plants.

**Other State and Local Requirements:** Applicators must follow all state and local pesticide drift requirements regarding application of triclopyr or 2,4-D herbicides in aquatic sites. Where states have more stringent regulations, they must be observed.

### Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

### Aerial applications

- Apply <u>SP1037Renovate MAX G</u> at the lowest height consistent with efficacy and flight safety. <u>Do not</u> apply at a height greater than 10 feet above the water surface or plant canopy unless a greater height is required for aircraft safety.
- When applications are made with a crosswind, the swath will be displaced downwind. The
  applicator must compensate for this by adjusting the path of the aircraft upwind.

### AQUATIC WEEDS CONTROLLED BY SP1037RENOVATE MAX G

Efficacy and selectivity of SP1037<u>Renovate MAX G</u> is dependent upon dose, time of year, stage of growth, method of application, and water movement. The following categories –

12/29

highly susceptible, moderately susceptible, and less susceptible – are provided to define species that may be controlled using <u>SP1037Renovate MAX G</u>.

Efficacy and selectivity is dependent on many factors, and can be managed through selection of application rates, application techniques and timing, etc. Rate selection will be partially dependent on characteristics of the treatment area. Consult with SePRO Corporation to determine best treatment protocols to manage individual species and to meet specific aquatic plant management objectives. Plants listed as moderately susceptible and less susceptible can be controlled under most use conditions, but generally require higher application rates.

### HIGHLY SUSCEPTIBLE VASCULAR AQUATIC PLANTS

Eurasian watermilfoil (*Myriophyllum spicatum*) hybrid watermilfoil (*Myriophyllum spicatum x sibiricum*)

### MODERATELY SUSCEPTIBLE VASCULAR AQUATIC PLANTS

northern watermilfoil (*Myriophllyum sibiricum*) other milfoil species (*Myriophyllum* spp.) bladderwort (*Utricularia* spp.) white water lily (*Nymphaea* spp.) watershield (*Brasenia* spp.)

### LESS SUSCEPTIBLE VASCULAR AQUATIC PLANTS

variable-leaf milfoil (*Myriophyllum heterophyllum*) water stargrass (*Heteranthera dubia*) coontail (*Ceretophyllum demersum*) parrotsfeather (*Myriophyllum aquaticum*)<sup>††</sup> yellow water lily or spatterdock (*Nuphar* spp.) water chestnut (*Trapa natans*)<sup>†, ††</sup>

<sup>†</sup>Not for use in California

<sup>††</sup> Retreatment may be needed to achieve desired level of control.

### APPLICATION METHODS

### **Surface Application**

Use a mechanical spreader such as a fertilizer spreader, blower, mechanical seeder, an eductor system, or similar equipment capable of uniformly applying <u>SP1037Renovate MAX G</u>. Before spreading any product, carefully calibrate the application equipment. When using boats and power equipment, you must determine the proper combination of (1) boat speed, (2) rate of delivery from the spreader, and (3) width of swath covered by the granules.

Use the following formula to calibrate the spreader's delivery in pounds of SP1037Renovate MAX G per minute:

Pounds per Minute = miles per hour X swath width (feet) X pounds per acre 495

### Aerial Application (Helicopter Only)

Ensure uniform application. All equipment should be properly calibrated using blanks with a similar physical characteristics to <u>SP1037Renovate MAX G</u>. To avoid streaked, uneven or overlapped application, use an appropriate tracking device (e.g., GPS). Refer to the DRIFT

312

MANAGEMENT section of this label for additional precautions and instructions for aerial application.

### Floating-leaf and Emergent Weeds

For control of water lily's (*Nymphaea* spp. and *Nuphar* spp.), watershield (*Brasenia* spp.), and other susceptible emergent and floating-leaf herbaceous weeds, apply up to 1.0 to 5.0 ppm. Apply when plants are actively growing.

### **Submersed Weeds**

For control of Eurasian watermilfoil (*Myriophyllum spicatum*) and other submersed weeds, apply <u>SP1037Renovate MAX G</u> at concentrations up to 0.25 to 5.0 ppm in treated water. Rates should be selected according to the Table 3 (*CONCENTRATION OF* <u>SP1037RENOVATE MAX G</u> IN WATER). For optimal control, apply when Eurasian watermilfoil or other submersed weeds are actively growing.

When controlling Eurasian watermilfoil in plant communities containing other desirable susceptible species, selectivity may be enhanced generally by using a rate lower in the range, treatment timing, application technique, etc.; consult a SePRO Aquatic Specialist for site specific recommendations.

CONCENTRA	TION OF SP		LE 3 VATE MAX	G IN WA	TER (PPM	A.E.) †	
Average Water	0.25 ppm	0.5 ppm	1.0 ppm	2.0 ppm	4.0 ppm	5.0 ppm	
Depth (ft)							
1	4.7	9.4	18.8	37.5	75.0	93.7	
2	9.4	18.8	37.5	75.0	150.0	187.5	
3	14.1	28.1	56.3	112.5	225.0	281.2	
4	18.8	37.5	75.0	150.0	300.0	375.0	
5	23.4	46.9	93.8	187.5	375.0	468.7	

<sup>†</sup>Use of higher rates in the rate range is necessary to achieve desired control in areas of greater water exchange; when treating more mature plants; when targeting more difficult to control aquatic species; and when treating small areas in larger bodies of water (spot treatments). Lower concentrations are generally used when conducting early season large-scale treatments; and treating larger areas, more immature plants, and areas with less potential for rapid water exchange. Some areas may require a repeat application to control re-growth.

The following formula can be used to calculate applications rates based on depths exceeding 4 feet deep or when using a concentration not in the Table 3.

Pounds of SP1037Renovate MAX G per Acre = average depth x target ppm x 18.75

### Example Calculation:

6 foot average depth x 1.25 ppm x 18.75 = 140.6 pounds of SP1037Renovate MAX G per acre

**NOTE:** apply a maximum of 93.7 pounds of <u>SP1037Renovate MAX G</u> (13.5 it s a.e.)/acre-foot per application. Do not exceed 5.0 ppm during any single application.

### COMBINATIONS WITH OTHER HERBICIDES

SP1037<u>Renovate MAX G</u> may be combined or applied simultaneously with other herbicides to increase the weed control spectrum or enhance efficacy. Follow all applicable use directions, precautions, and restrictions on all labels used in the combination.

### IF ANY OF THE CONTENT OF THIS LABEL IS NOT UNDERSTOOD, OR YOU NEED FURTHER ASSISTANCE, CONTACT A SEPRO AQUATIC SPECIALIST WITH QUESTIONS SPECIFIC TO YOUR APPLICATION.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

**Nonrefillable Container Disposal (non-rigid, any size):** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### TERMS AND CONDITIONS OF USE

If terms of the following *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitations of Remedies*.

### WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation, as the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

### LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages are resulting from this product (including claims based on contract, negligence, strict iability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

(1) Refund of purchase price paid by buyer or user for product bought, or

(2) Replacement of amount of product used.

Lecci

5/29

To the extent consistent with applicable law SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. To the extent permitted by applicable law in no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* above and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer*" or *Limitations of Remedies* in any manner.

<sup>©</sup>Copyright \_\_\_\_\_ SePRO Corporation \*Trademark of SePRO Corporation [Base label for Non-dilutable, Non-Rigid containers, any size]



## SP1037 Renovate MAX G

AQUATIC HERBICIDE

### FOR CONTROL OF AQUATIC WEEDS IN PONDS; LAKES; RESERVOIRS; MARSHES; BAYOUS; DRAINAGE DITCHES; NON-IRRIGATION CANALS; AND RIVERS AND STREAMS THAT ARE QUIESCENT OR SLOW-FLOWING.

### **Active Ingredients**

triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt	4.0%
2,4-dichlorophenoxyacetic acid, dimethylamine salt:	
Other Ingredients	
Total	
Acid equivalence (a.e.): 14.4%	

### PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

Causes substantial, but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes.

### KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

	FIRST AID
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
lf swallowed	<ul> <li>Call a poison control center or doctor immediately for treatment advice</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If inhaled	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> </ul>

	Call a poison control center or doctor for further treatment advice.
If on skin	Take off contaminated clothing.
or clothing	<ul> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
	oduct container or label with you when calling a poison control center or doctor, or atment. In case of emergency endangering health or the environment involving

this product, call INFOTRAC at 1-800-535-5053.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

### ENVIRONMENTAL HAZARDS

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, in quiescent waters, apply the product to areas separated by untreated sections that can be treated after vegetation in treated areas has disintegrated. During the growing season, weeds decompose in a 2 to 4 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

AGRICULTURAL CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Nonrefillable Container Disposal (non-rigid, any size): Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refer to inside of label booklet for additional precautionary information and directions for use including first aid and storage and disposal.

**NOTICE:** Read the entire label. Use only according to label directions. **Before using this** product, read *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies* at end of label booklet. If terms are unacceptable, return at once unopened.

EPA Reg. No. 67690-50 FPL201<u>20125</u>10927

EPA Est. No.	cecce
SPC -	C

SePRO Corporation 11550 N. Meridian Street, Suite 600, Carmel, IN 46032 U.S.A.

**Aquatic Herbicide** 

Net Contents:

Lecce

[Editor's note: This portion of the master label contains Consumer/Homeowner Uses.]

[Front label for Household/Residential container]



## SP1037 Renovate MAX G

AQUATIC HERBICIDE

**READY-TO-USE** 

FOR CONTROL OF AQUATIC WEEDS IN PONDS AND WATER FRONT SHORELINES THAT ARE QUIESCENT OR SLOW-FLOWING.

### **Active Ingredients**

triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt	4.0%
2,4-dichlorophenoxyacetic acid, dimethylamine salt:	. 14.0%
Other Ingredients	. 82.0%
Total	100.0%
Acid equivalence (a.e.): 14.4%	

KEEP OUT OF REACH OF CHILDREN WARNING

Specialty Chemical: Do not ship or store with food, feeds, drugs or clothing.

See back panel booklet for additional precautionary statements, directions for use and warranty information.

### Net []

[Editorial Note: Bracketed [text] indicates marketing claims; text in curly brackets {text} show place holders for graphics.]

19/29

[Label booklet cover for Household/Residential containers]



## SP1037 Renovate MAX G

AQUATIC HERBICIDE

### **READY-TO-USE**

### **PRODUCT FACTS**


This product will provide selective herbicidal control of many unwanted aquatic plants species.

### AQUATIC WEEDS CONTROLLED

Controls Various Types of Nuisance Aquatic Weeds







(See complete list under the directions for use section of this label)

### How SP1037Renovate MAX G Works

SP1037Renovate MAX G is applied directly to water in a pond or water front shoreline. The active ingredients are quickly absorbed from the water through plant stems and foliage and from the hydrosoil by roots. SP1037Renovate MAX G is formulated on biodegradable granules that that when applied to water bodies immediately sink and deliver the product to the critical areas for controlling target weeds. Generally, target plants are controlled within 2 to 4 weeks after treatment, but depending on conditions and plant species can take up to 8 weeks or longer for complete control.

### Where to Apply

Apply SP1037Renovate MAX G herbicide to ponds and water front shorelines.

Questions, Comments or Information {graphic of phone} Call {graphic of internet/computer} www.

Open booklet for details. Made in U.S.A.

SePRO Corporation 11550 N. Meridian Street, Suite 600, Carmel, IN 46032 U.S.A.

EPA Reg. No. 67690-50 FPL201<u>20125</u>10927 SPC -

© Copyright \_\_\_\_\_ SePRO Corporation.

2029

[Label booklet text for Household/Residential containers]



## SP1037 Renovate MAX G

AQUATIC HERBICIDE

### PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Causes substantial, but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes.

# KEEP OUT OF REACH OF CHILDREN WARNING

	FIRST AID
lf in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
lf inhaled	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
lf on skin or clothing	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
or going for tre involving this p	uct container or label with you when calling a poison control center or doctor eatment. In case of emergency endangering health or the environment product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in

eccece

21/29

water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

### **GENERAL INFORMATION**

SP1037<u>Renovate MAX G</u> herbicide may be applied directly to water for the control of aquatic weeds. SP1037<u>Renovate MAX G</u> enhances target weed control, and provides selective control of many broadleaf weeds in ponds and on lake front shorelines.

SP1037<u>Renovate MAX G</u> is formulated on biodegradable granules that, when applied to water bodies, immediately delivers SP1037<u>Renovate MAX G</u> down to the critical areas for controlling target weeds. SP1037<u>Renovate MAX G</u> is quickly absorbed from the water through plant stems and foliage and from the hydrosoil by roots. Herbicidal symptoms are initially expressed 2 to 14 days following application and usually involve bending and twisting of apical sections and shoots of susceptible plants. Initial symptoms are followed by necrosis of terminal buds and above ground tissue. Generally, target plants are controlled within 2 to 4 weeks after treatment, but depending on conditions and plant species can take up to 8 weeks for complete control.

When applying <u>SP1037Renovate MAX G</u> follow all applicable use directions, precautions and limitations. All <u>SP1037Renovate MAX G</u> concentrations referred to in this label are based on acid equivalence (a.e.).

### IMPORTANT GENERAL USE INFORMATION AND PRECAUTIONS

- There are no restrictions on fishing, swimming, recreational use, or consumption of treated water for potable use by livestock, pet, or other animals following application of <u>SP1037Renovate MAX G</u>.
- <u>Do not</u> allow people or pets to enter the treatment area while <u>SP1037Renovate MAX G</u> is being applied.
- <u>Do not</u> allow people or pets to enter the treated area until dusts have settled.
- <u>Do not</u> apply to salt water bays or estuaries.
- Do not apply this product through any type of irrigation system.
- Applications to target areas are limited to two (2) per season.
- Apply a maximum of 93.7 pounds of SP1037Renovate MAX G (13.5 lbs a.e.)/acre-foot per application. <u>Do not</u> exceed 5.0 ppm during any single application.
- <u>Do not</u> apply within 21 days of previous application except when conducting split treatments. Split treatments, over relatively short periods of time (e.g., 1 to 4 days), may be effective in some areas to maintain adequate exposure with target plants, such as small sites or sites with higher dilution potential.
- Obtain Required Permits: Consult with appropriate state or local water authorities before applying <u>SP1037Renovate MAX G</u>. If the pond/lake is not privately owned by you be sure

to check with state or local water authorities who have regulatory authority over public water as permits may be required by state or local public agencies.

### APPLICATION TO WATERS USED FOR IRRIGATION Irrigation Restrictions

- <u>Do not</u> use treated water for irrigating greenhouse or nursery plants unless triclopyr and 2,4-D residues are confirmed to be less than 1 ppb by laboratory analysis.
- <u>Do not</u> use water treated with <u>SP1037Renovate MAX G</u> for hydroponic farming.
- <u>Do not</u> apply <u>SP1037Renovate MAX G</u> directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers, or other desirable susceptible broadleaf plants, and <u>do not</u> permit dust to drift into these areas.
- This label describes both required and recommended uses of a chemical analyses for the active ingredients, triclopyr and 2,4-D. SePRO Corporation recommends the use of an Enzyme-Linked Immunoassay (ELISA) test for the determination of SP1037Renovate MAX G concentration in water. Contact SePRO Corporation for the incorporation of these analyses into your treatment program. Other proven chemical analyses for the active ingredients may also be used. The ELISA analysis is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredients in the water. Both triclopyr and 2,4-D can be analyzed from a single water sample.
- If <u>SP1037Renovate MAX G</u> treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with triclopyr and 2,4-D such as pastures, turf or established grasses, the treated water may be used to irrigate and/or mix sprays for these sites at any time during and after the application.
- Due to potential phytotoxicity and/or residue considerations, the following restrictions are applicable to other uses of irrigation water.
  - If treated water is intended to be used to irrigate or mix sprays for crops not labeled for direct treatment with triclopyr and 2,4-D, the water must not be used unless <u>one</u> of the following restrictions has been observed:
    - A waiting period of 120 days from the time of application has elapsed; or
    - An approved assay indicates that the triclopyr concentration is 1.0 ppb or less and the 2,4-D concentration is 100 ppb or less at the water intake. See Table 2 (SAMPLING FOR DRINKING WATER ANALYSES) for the recommended waiting periods after application but before taking the initial water sample at water intake.
  - If treated water is intended to be used to irrigate non-crop areas not labeled for direct treatment with triclopyr and 2,4-D (e.g., landscape ornamentals) or for other irrigation uses not described, consult with SePRO Corporation prior to commencing irrigation if triclopyr concentrations exceed 1.0 ppb and 2,4-D concentrations are greater than 100 ppb.

### APPLICATIONS TO POTABLE WATER SOURCES Potable Water Restrictions

- The potable water use restrictions on this label are to ensure that consemption of water by the public is allowed only when the concentration of triclopyr in water is less than 400 pbb and the concentration of 2,4-D in water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that triclopyr and 2,4-D concentrations in potable water do not exceed 400 ppb and 70 ppb, respectively, at the time of consumption.
- The drinking water setback distances from functioning potable water intakes are provided in Table 1 (DRINKING WATER SETBACK DISTANCES).

23/29

		TABLE 1			
	DRINKING WATE	R SETBACK DISTAN	NCES		
Application conce		n setback distance (ft ater intake	) from functioning potable		
<u>≤</u> 1 ppm <sup>†</sup>	1.1 to 2.0 ppm <sup>†</sup>	2.1 to 3.0 ppm <sup>†</sup>	3.1 to 5.0 ppm <sup>†</sup>		
600 1,200 1,800 2,400					
nom acid equival	ent target water conc	entration			

ppm acid equivalent target water concentration

- Following each application of <u>SP1037Renovate MAX G</u>, treated water must not be used for potable water unless <u>one</u> of the following restrictions has been observed:
  - A setback distance described in Table 1 was used for the application;
  - A waiting period of at least 21 days from the time of application has elapsed; or
  - An approved assay indicates that the triclopyr concentration is 400 ppb or less and the 2,4-D concentration is 70 ppb or less at the water intake. Sampling for drinking water analyses should occur no sooner than stated in Table 2. NOTE: Sampling for drinking water analysis should occur no sooner than 3 days after SP1037Renovate MAX G application. Analysis of 2,4-D in drinking water samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

SAMPLIN	IG FOR DRINKING WATER A	NALYSES <sup>†</sup>
Minimum days after ap	oplication before initial water sa	ampling at the functioning
	potable water intake	
≤1 ppm <sup>††</sup>	1.1 to 3.0 ppm <sup>††</sup>	3.1 to 5.0 ppm <sup>††</sup>

These are general guidelines; the amount of time required for residues to reach concentrations acceptable for drinking or irrigation will depend on the total acres treated relative to water body size, application rates, water exchange rates, weed density, and various other factors. Consult a SePRO Aquatic Specialist for site specific recommendations.

<sup>††</sup> ppm acid equivalent target water concentration

If no setback distance from Table 1 is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an advisory to shut off all potable water intakes inside the setback zone prior to SP1037<u>Renovate MAX G</u> application. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water. The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

### • Example:

Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting should include the day and time of application. Posting may be removed if analyses of a sample collected at the intake, no sooner than stated in Table 2, shows that the triclopyr concentration in

24/29

the water is 400 ppb or less and the 2,4-D concentration is 70 ppb or less, or after 21 days following application, whichever occurs first.

- **Text of notification:** Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water unless water at functioning drinking water intakes is tested no sooner than [insert days from Table 2] and is demonstrated by assay to contain no more than 400 ppb triclopyr and 70 ppb 2,4-D. Application Date: \_\_\_\_\_ Time: \_\_\_\_.
- **NOTE**: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

### **GRAZING AND HAYING RESTRICTIONS**

Except for lactating dairy animals, there are no grazing restrictions following application of this product.

- Grazing Lactating Dairy Animals: <u>Do not</u> allow lactating dairy animals to graze treated areas until the next growing season following application of this product.
- Do not harvest hay for 14 days after application.
- Grazed areas of sites may be spot treated if they comprise no more than 10% of the total grazable area.

### SLAUGHTER RESTRICTIONS

During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

### AQUATIC WEEDS CONTROLLED BY SP1037RENOVATE MAX G

The following categories – highly susceptible, moderately susceptible, and less susceptible – are provided to define species that may be controlled using <u>SP1037Renovate MAX G</u>.

Efficacy and selectivity is dependent on many factors, and can be managed through selection of application rates, application techniques and timing, etc. Plants listed as moderately susceptible and less susceptible can be controlled under most use conditions, but generally require higher application rates. Consult with SePRO Corporation to determine best treatment protocols.

### HIGHLY SUSCEPTIBLE VASCULAR AQUATIC PLANTS

Eurasian watermilfoil (*Myriophyllum spicatum*) hybrid watermilfoil (*Myriophyllum spicatum x sibiricum*)

### MODERATELY SUSCEPTIBLE VASCULAR AQUATIC PLANTS

northern watermilfoil (*Myriophllyum sibiricum*) other milfoil species (*Myriophyllum* spp.) bladderwort (*Utricularia* spp.) white water lily (*Nymphaea* spp.) watershield (*Brasenia spp*.)

### LESS SUSCEPTIBLE VASCULAR AQUATIC PLANTS

variable-leaf milfoil (*Myriophyllum heterophyllum*) water stargrass (*Heteranthera dubia*) coontail (*Ceretophyllum demersum*) parrotsfeather (*Myriophyllum aquaticum*)<sup>††</sup>

Page 21 of 26

yellow water lily or spatterdock (*Nuphar* spp.) water chestnut (*Trapa natans*)<sup>†, ††</sup>

<sup>†</sup>Not for use in California

<sup>††</sup> Retreatment may be needed to achieve desired level of control.

### **APPLICATION METHODS**

Use a mechanical spreader such as a fertilizer spreader, blower, mechanical seeder, an educator system, or similar equipment capable of uniformly applying <u>SP1037Renovate MAX</u> <u>G</u>. Before spreading any product, carefully calibrate the application equipment according to equipment manufacturer's instructions.

Rates should be selected according to the Table 3 (CONCENTRATION OF <u>SP1037Renovate</u> <u>MAX G</u> IN WATER).

### Floating-leaf and Emergent Weeds

For control of susceptible emergent and floating-leaf herbaceous weeds, apply up to 1.0 to 5.0 ppm. Apply when plants are actively growing.

### Submersed Weeds

For control of submersed weeds, apply <u>SP1037Renovate MAX G</u> at concentrations up to 0.25 to 5.0 ppm in treated water.

			LE 3			
CONCENTRA	ATION OF SP	1037 <b>RENO</b>	VATE MAX	G IN WA	TER (PPM	A.E.) <sup>†</sup>
Avg. Water	0.25 ppm	0.5 ppm	1.0 ppm	2.0 ppm	4.0 ppm	5.0 ppm
Depth (ft)	Pounds SP1037Renovate MAX G / acre					
1	4.7	9.4	18.8	37.5	75.0	93.7
2	9.4	18.8	37.5	75.0	150.0	187.5
3	14.1	28.1	56.3	112.5	225.0	281.2
4	18.8	37.5	75.0	150.0	300.0	375.0
5	23.4	46.9	93.8	187.5	375.0	468.7

<sup>†</sup>Use of higher rates in the rate range is necessary to achieve desired control in areas of greater water exchange; when treating more mature plants; when targeting more difficult to control aquatic species; and when treating small areas in larger bodies of water (spot treatments). Lower concentrations are generally used when: conducting early season large-scale treatments; and treating larger areas, more immature plants, and areas with less potential for rapid water exchange. Some areas may require a repeat application to control re-growth.

The following formula can be used to calculate applications rates based on depths exceeding 5 feet deep or when using a concentration not in the Table 3.

Pounds of SP1037Renovate MAX G per Acre = average depth x target ppm x 18.75

### Example Calculation:

6 foot average depth x 1.25 ppm x 18.75 = 140.6 pounds of SP1037Renovate MAX G per acre

26/29

**NOTE:** apply a maximum of 93.7 pounds of <u>SP1037Renovate MAX G</u> (13.5 lbs a.e.)/acre-foot per application. Do not exceed 5.0 ppm during any single application.

### Small Site (Less than 1/2 Acre)/Spot Treatment Application

For small treatment sites of one-half (½) acre or less in larger bodies of water, use Table 4 (SP1037<u>Renovate MAX G</u> RECOMMENDED RATES FOR AREAS LESS THAN ½ ACRE), to determine the application rate depending on average water depth to target an application rate of 3.0 to 5.0 ppm. For best results, split the total application rate into three equal applications 8 to 12 hours apart and apply when water is relatively calm. Small treatment application of SP1037<u>Renovate MAX G</u> is recommended with waterproof gloves or a hand spreader to uniformly distribute on target weeds.

SP1037 <u>REN(</u>		TABLE 4 RECOMMENDED THAN ½ ACRE SP1037Renovat		REAS LESS	
Area (ft <sup>2</sup> )	3 foot ave	3 foot average depth		4 foot average depth	
	3.0 ppm	5.0 ppm	3.0 ppm	5.0 ppm	
500	1.9	3.2	2.6	4.3	
1,000	3.9	6.4	5.2	8.6	
4,000	15.5	25.8	20.7	34.4	
10,000	38.7	64.5	51.7	86.0	
20,000	77.5	129.1	103.3	172.1	

Example Scenario: A 100 ft. by 40 ft. lakeshore swimming area with a 4 ft. average depth, heavily infested with Eurasian watermilfoil

- Step 1: Determine the area to be treated in square feet (ft<sup>2</sup>) by multiplying the length of the area by the width.
   → 100 ft. x 40 ft. = 4,000 ft<sup>2</sup>
- Step 2: Determine the amount of SP1037<u>Renovate MAX G</u> to be used by consulting the Table 4.
   → Use 34.4 pounds based on 4 foot average depth and 5.0 ppm.
- Step 3: Apply SP1037Renovate MAX G uniformly over weeds in treatment site in three equal applications of 11.4 lbs. each, 8 12 hours apart.

**NOTE:** For applications with an area or depth not included in the above chart, the following formula can be used to calculate application rates.

Pounds of SP1037 Renovate MAX G = area (ft<sup>2</sup>)/43,560 x average depth x target ppm x 18.75

### Example Calculation:

8,250 ft<sup>2</sup>/43,560 x 2 foot average depth x 4.0 ppm x 18.75 = 28.4 pounds of <u>SP1037Renovate</u> MAX G

7/29

### IF ANY OF THE CONTENT OF THIS LABEL IS NOT UNDERSTOOD, OR YOU NEED FURTHER ASSISTANCE, CONTACT A SEPRO AQUATIC SPECIALIST WITH QUESTIONS SPECIFIC TO YOUR APPLICATION.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage**: Store product in original container in a safe place. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste.

**Pesticide and Container Disposal**: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. <u>If Empty</u>: Do not reuse this container. Place in trash or offer for recycling if available. <u>If Partly Filled</u>: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### TERMS AND CONDITIONS OF USE

If terms of the following *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitations of Remedies*.

### WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

### LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. To the extent permitted by applicable law in no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

28/29

The terms of the *Warranty Disclaimer* above and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer*" or *Limitations of Remedies* in any manner.

<sup>©</sup> Copyright _	SePRO Corporation
*Trademark	of SePRO Corporation

29/29

[Base label for Household/Residential containers]



## SP1037 Renovate MAX G

AQUATIC HERBICIDE

### **READY-TO-USE**

### SePRO FAMILY OF PRODUCTS

Visit the SePRO website **www.lakelawnandpond.com** to learn more about the SePRO family of products for the best solution to your toughest pond and lakeshore aquatic weed and algae problems.

{Insert graphic - SePRO product family photo}

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage**: Store product in original container in a safe place. Do not store near feed or foodstuffs. In case of spill, contain material and dispose as waste. **Pesticide and Container Disposal**: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. <u>If Empty</u>: Do not reuse this container. Place in trash or offer for recycling if available. <u>If Partly Filled</u>: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Specialty Chemical: Do not ship or store with food, feeds, drugs or clothing.

Consumer Promise: 100% Satisfaction Guaranteed.

### **Questions, Comments or Information**

{graphic of phone} Call \_\_\_\_\_\_ {graphic of internet/computer} www.

Always read and follow label directions. Made in U.S.A.

SePRO Corporation 11550 N. Meridian Street, Suite 600, Carmel, IN 46032 U.S.A.

EPA Reg. No. 67690-50 FPL201<u>20125</u>10927 EPA Est. No.

cecce

© Copyright \_\_\_\_\_ SePRO Corporation.