



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 17, 2018

Laurent C. Mezin  
Regulatory and Government Affairs Leader  
SePRO Corporation  
11550 N. Meridian Street, Suite 600  
Carmel, IN 46032-4562

Subject: Label Amendment – Amending CA specific use rates  
Product Name: SP5100 TURF GROWTH REGULATOR  
EPA Registration Number: 67690-51  
Application Date: 1/23/2018  
Decision Number: 537894

Dear Joshua Wood:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 537894  
Decision No. 67690-51

with FIFRA section 6. If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at [mellor.nathan@epa.gov](mailto:mellor.nathan@epa.gov).

A handwritten signature in cursive script that reads "Shaja B. Joyner".

Shaja B. Joyner, Product Manager 20  
Fungicide-Herbicide Branch  
Registration Division 7505P

Enclosure

*[Editor's note: Version A sub-label - turfgrass uses.]*  
*[Front label, ALL containers]*



# Cutless<sup>®</sup> MEC

## Turf Growth Regulator

**FOR GROWTH MANAGEMENT AND QUALITY IMPROVEMENT OF PERENNIAL COOL AND WARM SEASON TURFGRASSES ON GOLF COURSES.**

**Active Ingredient:**

flurprimidol:  $\alpha$ -(1-methylethyl)- $\alpha$ -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol ..... 16.0%

**Other Ingredients** ..... 84.0%

**TOTAL** ..... 100.0%

Contains 1.3 pounds active ingredient per one (1) U.S. gallon.

**Keep Out of Reach of Children**

**WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b>.</p>	

**Refer to inside of label booklet for additional Precautionary Statements and Directions for Use, including Storage and Disposal.**

**Notice:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements inside label booklet.**

®Cutless is a registered trademark of SePRO Corporation

EPA Reg. No. 67690-51  
FPL20180517

EPA Est. No. \_\_\_\_\_  
[P/N] - \_\_\_\_\_

**SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.**

**Net Contents \_\_\_\_\_ (Non-refillable)**

**[Editor's note: Version A sub-label - turfgrass uses.]**  
**[Label booklet text]**

---

## **PRECAUTIONARY STATEMENTS**

---

### **Hazards to Humans and Domestic Animals**

#### **Keep Out of Reach of Children**

#### **WARNING / AVISO**

**Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.**

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

#### **Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants;
- Shoes plus socks; and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### **USER SAFETY RECOMMENDATIONS**

#### **Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment water or rinsate.

---

## **DIRECTIONS FOR USE**

---

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying this product. Use only according to label directions.

**Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

### **Avoid Spray Drift**

Applications must be made only when there is no hazard for spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants.

**Do not spray when wind is greater than 10 mph.** Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

### **INFORMATION FOR GROWTH REGULATION OF PERENNIALTURFGRASSES**

Cutless MEC Turf Growth Regulator is a Type II Class B plant growth regulator (PGR) which reduces leaf blade length and stem internode elongation in turfgrasses resulting in a more compact growth form. Growth regulation results from suppression of gibberellic acid biosynthesis. Under normal growing conditions root growth and lateral expansion of turf are not affected. Follow an appropriate fertility program for the desired turf species in conjunction with this product's applications to provide best turfgrass enhancement and reduce potential for discoloration. Make broadcast treatments on medium to high quality turfgrass areas of uniform species composition. Turf containing significant amounts of coarse textured species including tall fescue, orchardgrass, timothy, dallisgrass, etc., may respond unevenly to treatment.

### **Benefits of Cutless MEC Applications to Perennial Turfgrasses may include:**

- Shoot growth suppression of warm and cool season turfgrasses resulting in decreased mowing frequency and turfgrass clippings;
- Increased turfgrass density, wear resistance, and improved color on warm and cool season turfgrass species resulting in improved turf quality;
- Suppressed growth of *Poa annua*, reducing populations and shifting competitive growth advantage towards perennial turfgrasses;
- Improved water use efficiency of warm and cool season turfgrass resulting in pre-drought stress conditioning.

**NOTICE TO USER:** The rates indicated may need to be adjusted within the approved rate ranges on this label to achieve the desired level of growth regulation on turfgrass species listed on this label. Turfgrass response to Cutless MEC may vary within turfgrass species due to the large number of cultivars and varieties available. The specified rate ranges permit the users to adjust the application rate to best address the growth conditions of the turfgrass being treated. Neither the manufacturer nor seller has determined if this product can be used safely or effectively on turfgrass species not mentioned on this label. Apply this product to a small test area to determine growth response and desired level of growth regulation prior to large scale applications for turfgrass species not listed on this label.

## Use Restrictions for Applications to Perennial Turfgrasses

- Do not apply to putting greens other than those where bentgrass is the desired turf species.
- Do not apply to bermudagrass putting greens, including those which are overseeded.
- Do not apply this product to sod farms, turfgrasses grown for seed, including plants or plant materials grown for sale or research purposes.
- Do not apply to shrubs, bedding plants, and/or food plants.
- Do not apply during prolonged periods of temperature (heat or cold) or moisture stress. Also avoid applications during periods of extreme disease and insect pressure.
- Do not apply until 6 to 8 weeks after turfgrass sprigging or laying sod. Turfgrass must be well established and actively growing prior to application.
- Do not apply to saturated soils or when a significant moisture event is anticipated. This product may accumulate in low lying areas and cause prolonged and excessive growth regulation in those areas.
- Do not apply to areas where *Poa annua* is the desired turfgrass species or areas that contain >80% *Poa annua*.
- Do not apply to turf used for livestock production.
- Do not apply more than 3.0 lbs AI/A/year or 295.4 fl. oz./A/year of Cutless MEC.
- **Chemigation:** Do not apply Cutless MEC through any type of irrigation system.

## Application Timing

Spring applications must be made after resumption of active seasonal growth of turfgrass. The final application of the season must be timed at least 4 weeks before the onset of inactive grass growth or winter dormancy. Applications to overseeded turfgrasses in dormant bermudagrass stands must be completed 4 weeks prior to expected bermudagrass green-up.

## Irrigation

Cutless MEC is primarily root absorbed. To maximize growth regulator activity, treated areas should receive 0.25 to 0.5 inch of rain or supplemental irrigation within 24 hours following application and prior to the first mowing after treatment.

## Turf Color and Post-Treatment Turf Management

Turfgrass treated with Cutless MEC may appear darker green in color. This color change, which appears 1 to 2 weeks after treatment, may persist an additional 3 to 6 weeks. Manage treated areas to encourage the growth of a healthy vigorous turf.

## *Poa annua* (Annual Bluegrass) Conversion to Perennial Turfgrasses

Applications of Cutless MEC followed by management practices designed to encourage vigorous growth of perennial turfgrasses can reduce the *Poa annua* (annual bluegrass) competition in cool season turf and increase the cover of more desirable perennial species. *Poa annua* is more sensitive to Cutless MEC treatments and is therefore more strongly suppressed than perennial turfgrass species. Discoloration of some *Poa annua*

biotypes can be expected from treatments that provide a desired level of growth regulator activity in perennial grass species. This effect becomes visible 7 to 10 days after treatment and lasts 3 to 6 weeks. The degree of discoloration will be proportional to the *Poa annua* composition of the turf. Application of this product in conjunction with soluble nitrogen fertilizers will also minimize discoloration. Application timing, rate ranges, and precautions for perennial grass conversion through selective reduction of *Poa annua* are provided in the *Poa annua (Annual Bluegrass) Conversion to Perennial Turfgrasses* section of this label.

## **APPLICATION DIRECTIONS**

Applications of this product to newly seeded turfgrasses should be delayed until turf is well established and actively growing. Additional turfgrass growth regulation may occur when Cutless MEC is tank mixed or used in conjunction with demethylation inhibitor (DMI) or sterol inhibiting fungicides.

**California:** For effective bermudagrass or seashore paspalum growth regulation in California, apply at a minimum 24.6 fl oz/A when not applied in combination with a Type II Class A PGR. When applied in combination with a Type II Class A PGR, apply Cutless MEC at a minimum 8.5 fl oz/A.

## **Mixing Directions**

Add Cutless MEC to a spray tank half filled with clean water. Begin agitation allowing sufficient mixing time to ensure complete dispersion and mixing of this product. Finish filling the spray tank. Continue agitation throughout the spraying operation to ensure uniform application. Apply using a boom-type sprayer with bypass and/or mechanical agitation calibrated to deliver 40 to 200 gallons/acre of spray solution (1 to 4.6 gallons/1000 ft<sup>2</sup>). In-line strainers and nozzle screens must be 50 mesh or larger. The use of a coloring agent to mark areas already sprayed is suggested for uniform application without skips and overlaps. Performance may be improved with the addition to the spray mix of a readily available nitrogen (N) source at 0.125 to 0.5 lbs N/1000 ft<sup>2</sup> or iron (Fe) at labeled rates.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## **Cutless MEC + Tank Mixtures**

Cutless MEC can be tank mixed and is compatible with most commonly-used pesticides and foliar nutrient products. However, test compatibility of this product with tank mix partners before use.



**NOTE:** Test the compatibility of this product in any tank-mix combination before use. To determine the physical compatibility with other products, use a jar test as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure sequence for adding required ingredients to the spray tank.

**Read and follow all label directions for each tank mix product.**

### **CUTLESS MEC RATES FOR GROWTH REDUCTION OF PERENNIAL TURFGRASS SPECIES – MULTIPLE APPLICATION PROGRAM**

The maximum number of annual applications is determined by the sum of the rates applied, not to exceed 3.0 lbs ai/A or 295.4 fl. oz. Cutless MEC/A (6-48 applications/year).

- **Cool Season Turfgrasses**

- Bentgrass (Golf Course Fairway Type Turf)

**Multiple application program:** Apply to fairway height bentgrass at a rate of 24.6 to 49.2 fl. oz./acre in early spring following resumption of active growth. Repeat applications of 12.3 to 49.2 fl. oz./acre of this product may be made at 2 to 6 week intervals until late summer or early fall.

- Bentgrass Putting Greens

**Multiple application program:** Apply to bentgrass as part of an overall greens management program. An initial application may be made after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Apply at a rate of 12.3 to 24.6 fl. oz. product/acre. Repeat applications of 6.1 to 24.6 fl. oz. product/acre may be made at 2 to 4 week intervals.

- Kentucky Bluegrass and Perennial Ryegrass

**Multiple application program:** Apply to Kentucky bluegrass and perennial ryegrass at a rate of 36.9 to 49.2 fl. oz. product/acre in early spring following resumption of active growth. Repeat applications of 24.6 to 49.2 fl. oz./acre may be made at 2 to 6 week intervals until late summer or early fall.

- **Warm Season Grasses**

- Seashore Paspalum

**Multiple application program:** Apply to Seashore paspalum at a rate of 12.3 to 49.2 fl. oz. product/acre when paspalum has completely recovered from winter

dormancy and is growing vigorously. Repeat applications of 12.3 to 49.2 fl. oz. product/acre may be made at 3 to 6 week intervals until late summer or early fall.

- Tifway (419), TifSport, and GN-1 Bermudagrass

**Multiple application program:** Apply to Tifway, TifSport, and GN-1 bermudagrass at a rate of 12.3 to 36.9 fl. oz. product/acre when bermudagrass has completely recovered from winter dormancy and is growing vigorously. Repeat applications of 12.3 to 36.9 fl. oz. product/acre may be made at 3 to 6 week intervals until late summer or early fall.

- Zoysiagrass

**Multiple application program:** Apply at a rate of 12.3 to 36.9 fl. oz. product/acre for growth regulation of zoysiagrass when the grass has completely recovered from winter dormancy and is growing vigorously. Repeat applications of 12.3 to 36.9 fl. oz. product/acre may be made at 3 to 6 week intervals. NOTE: late summer or early fall applications of this product are not to be used for growth regulation of zoysiagrass.

<b>TABLE 1</b>			
<b>Rate Ranges for Growth Regulation of Perennial Turfgrass Species with Cutless MEC Using a Multiple Application Program</b>			
<b>Turfgrass Species</b>	<b>Initial Spring Application<sup>†</sup> (Fl. Oz. Cutless MEC/A)</b>	<b>Repeat Applications<sup>†</sup></b>	
		<b>(Fl. Oz. Cutless MEC/A)</b>	<b>Treatment Interval</b>
<b>Cool Season Turfgrasses</b>			
Bentgrass (golf course fairway)	24.6 to 49.2	12.3 to 49.2	2 to 6 weeks
Bentgrass putting greens	12.3 to 24.6	6.1 to 24.6	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass	36.9 to 49.2	24.6 to 49.2	2 to 6 weeks
<b>Warm Season Turfgrasses</b>			
Seashore Paspalum <sup>††</sup>	12.3 to 49.2	12.3 to 49.2	3 to 6 weeks
Tifway, TifSport, and GN-1 Bermudagrass <sup>††</sup>	12.3 to 36.9	12.3 to 36.9	3 to 6 weeks
Zoysiagrass	12.3 to 36.9	12.3 to 36.9	3 to 6 weeks; not in late summer/fall

<sup>†</sup> Apply in early spring following resumption of active growth of the grass. Fall applications must be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

<sup>††</sup> California: Apply at 24.6 to 49.2 fl. oz./A

**APPLICATION INFORMATION - CUTLESS MEC TANK MIXES WITH PRIMO MAXX® (TRINEXAPAC-ETHYL)**

Rainfall or irrigation must be delayed at least 1 hour after application or until product has dried on the leaf surface but should occur within 24 hours after application.

PGR absorption via the foliage (Type II Class A; Primo MAXX®) and roots (Type II Class B; Cutless MEC) maximizes plant uptake of each material ensuring sufficient active ingredient is available for GA inhibition.

Tank mixing Cutless MEC with Primo MAXX® can provide enhanced growth suppression and improved turfgrass quality of perennial turfgrass species, versus using either product alone. Plant physiological advantages of tank mixing these two PGRs include:

1. Different mode of action within the gibberellic acid (GA) biosynthesis pathway; and
2. Difference in plant site of uptake.

Blocking GA biosynthesis early and late in the pathway regulates GA more efficiently than higher application rates of individual compounds. Turfgrass response from Cutless MEC tank mixes with Primo MAXX® at reduced rates may be greater than with normal use rates of either product applied alone. This combination of plant growth regulators and its use are protected by United States Patent No. 7,135,435. Additional patent rights pending.

Desirable growth regulation during early spring or late fall when turfgrass growth and vigor are reduced may be obtained with lower rates of both products.

**NOTICE TO USERS:** To the extent consistent with applicable law, this label makes no warranties concerning the performance of Type II Class A PGRs, including Primo MAXX®. Read and follow all label directions including Directions for Use, Precautionary Statements, and Restrictions and Limitations for Primo MAXX®.

<b>TABLE 2</b>			
<b>Rate Ranges for Cutless MEC Plus Type II Class A PGR Tank Mixes</b>			
<b>Turfgrass Species</b>	<b>Fl. Oz. Cutless MEC /A</b>	<b>Primo MAXX® (Trinexapac-ethyl)</b>	<b>Treatment Interval</b>
<b>Cool Season Turfgrasses</b>			
Bentgrass (golf course fairway)	6.1 to 24.6	½ labeled use rate	2 to 6 weeks
Bentgrass putting greens	6.1 to 12.3	½ labeled use rate	2 to 6 weeks
Kentucky Bluegrass/ Perennial Ryegrass	12.3 to 24.6	½ labeled use rate	2 to 6 weeks
<b>Warm Season Turfgrasses</b>			
Seashore Paspalum	12.3 to 24.6	½ labeled use rate	3 to 6 weeks
Tifway, TifSport, and GN-1 Bermudagrass†, ††	6.1 to 24.6	½ labeled use rate	3 to 6 weeks
Zoysiagrass	12.3 to 24.6	½ labeled use rate	3 to 6 weeks

† Do not use on bermudagrass putting greens.

†† California: Apply at 8.5 to 24.6 fl. oz./A

## **POA ANNUA (ANNUAL BLUEGRASS) CONVERSION TO PERENNIAL TURFGRASSES**

- **Multiple Application Program** - A multiple application perennial grass conversion program using Cutless MEC provides *Poa annua* suppression and cool season turf conversion. This program provides a gradual perennial grass conversion reducing *Poa annua* populations over one to several years. To maximize seedling establishment, do not apply this product two (2) weeks prior to and/or 2 weeks after interseeding or overseeding of perennial turfgrasses.

The maximum number of annual applications is determined by the sum of the rates applied, not to exceed 3.0 lbs ai/A or 295.4 fl. oz. product/A (6-48 applications/year).

- Bentgrass (Golf Course Fairway type turf)

Apply Cutless MEC to fairway height bentgrass at a rate of 24.6 to 49.2 fl. oz./acre in early spring following resumption of active growth of the grass. Repeat applications of 12.3 to 49.2 fl. oz. product/acre may be made at 2 to 6 week intervals until late summer or early fall. Normal management practices including fertilization, aeration and interseeding/overseeding will encourage growth of bentgrass.

- Bentgrass Putting Greens

Because annual turfgrass species, for example *Poa annua* are more strongly regulated by Cutless MEC, careful use of this product on putting greens with a high percentage of *Poa annua* can shift the competitive balance between bentgrass and *Poa annua* to favor bentgrass. Follow normal management practices including fertilization, aeration and interseeding/overseeding that encourages growth of bentgrass. Use of this product on bentgrass greens may increase putting speed without reducing the height of cut.

- Bentgrass Greens with less than 50% *Poa annua* (annual bluegrass): Apply Cutless MEC to bentgrass as part of an overall greens management program. An initial application may be made in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Apply at a rate of 12.3 to 24.6 fl. oz. product/acre. Repeat applications of 12.3 to 24.6 fl. oz. product/acre may be made at 2 to 4 week intervals through early fall.
- Bentgrass Greens with more than 50% *Poa annua* (annual bluegrass): Apply Cutless MEC to bentgrass greens at a rate of 12.3 fl. oz./acre in the spring months after bentgrass greens are growing vigorously and have been mowed 3 or 4 times. Repeat applications of 12.3 to 24.6 fl. oz. product/acre may be made at 2 to 4 week intervals through early fall.

- Kentucky Bluegrass, Perennial Ryegrass

Apply Cutless MEC to Kentucky bluegrass and perennial ryegrass at a rate of 36.9 to 49.2 fl. oz./acre in early spring following resumption of active growth of the grass. Repeat applications of 24.6 to 49.2 fl. oz. product/acre may be made at 3 to 6 week intervals until late summer or early fall. Normal management practices including fertilization, aeration and interseeding/overseeding will encourage growth of bluegrass and/or ryegrass.

Turfgrass Species	% <i>Poa annua</i>	Initial Spring Application <sup>†</sup>	Repeat Applications <sup>†</sup>	
		Fl. Oz. Cutless MEC /A	Fl. Oz. Cutless MEC /A	Treatment Interval
Bentgrass (golf course fairway)	0 - 80%	24.6 to 49.2	12.3 to 49.2	2 to 6 weeks
Bentgrass Putting Greens	Less than 50%	12.3 to 24.6	12.3 to 24.6	2 to 4 weeks
	More than 50%	12.3	12.3 to 24.6	2 to 4 weeks
Kentucky Bluegrass/ Perennial Ryegrass Fairways	0 - 80%	36.9 to 49.2	24.6 to 49.2	3 to 6 weeks

<sup>†</sup> Apply in early spring following resumption of active growth of the grass. Fall applications must be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

- **Alternative Spring/Fall Application Program** - The alternative spring/fall application program selectively suppresses *Poa annua* growth in perennial turfgrass species. Injury or discoloration of *Poa annua* is expected within 7 to 10 days of application. Perform cultural practices including fertilization, aeration and interseeding/overseeding in conjunction with Cutless MEC applications to encourage growth of the perennial turfgrass species.
  - **Bentgrass or Kentucky Bluegrass/Perennial Ryegrass fairways with less than 50% *Poa annua* (annual bluegrass):** An alternative spring/fall treatment program for management of *Poa annua* in fairways containing less than 50% *Poa annua* is to apply 73.8 fl. oz./acre of Cutless MEC following resumption of active bentgrass or Kentucky Bluegrass/Perennial Ryegrass growth in the spring. A second application of 49.2 fl. oz. product/acre of may be made in the early fall. Interseed/overseed with Bentgrass or Kentucky Bluegrass/Perennial Ryegrass 3 to 4 weeks following early fall application.
  - **Bentgrass or Kentucky Bluegrass/Perennial Ryegrass fairways with more than 50% *Poa annua* (annual bluegrass):** Apply Cutless MEC at a rate of 49.2 to 73.8 fl. oz./acre in late summer or early fall. Interseed/overseed with bentgrass or Kentucky Bluegrass/Perennial Ryegrass 3 to 4 weeks following late summer or early fall application. Apply an additional 49.2 fl. oz. product /acre of

the following spring after resumption of active growth of bentgrass or Kentucky Bluegrass/Perennial Ryegrass.

<b>TABLE 4</b> <b>Rate Ranges for <i>Poa annua</i> (Annual Bluegrass) Conversion to Perennial Turfgrasses with Cutless MEC Using an Alternative Spring/Fall Application Program</b>		
<b>Turfgrass Species</b>	<b>% <i>Poa annua</i></b>	<b>Spring/Fall application<sup>†</sup> (Fl. Oz. Cutless MEC /A)</b>
Bentgrass or Kentucky Bluegrass/Perennial Ryegrass Fairways	Less than 50%	73.8 in spring; 49.2 in fall
	More than 50%	49.2 to 73.8 in late summer or early fall; 49.2 the following spring

<sup>†</sup> Apply in early spring following resumption of active growth of the grass. Fall applications must be discontinued 4 weeks before the onset of inactive grass growth or winter dormancy.

**DOLLAR SPOT (*SCLEROTINIA HOMOEOCARPA*) SUPPRESSION BY CUTLESS MEC IN CREEPING BENTGRASS**

The active ingredient in Cutless MEC is from the pyrimidine class of chemistry which is structurally similar to pyrimidine fungicides that provide Dollar Spot control. Programmed applications of this product for turf growth suppression or *Poa annua* conversion have also been shown to suppress Dollar Spot incidence in creeping bentgrass fairways, greens and tees. Research results have shown Cutless MEC applications at labeled rates and application intervals can significantly reduce Dollar Spot incidence and populations when compared to untreated control plots. This product must not be used to replace labeled fungicides for the control of Dollar Spot; rather programmed use of this product may result in longer or improved control of Dollar Spot in conjunction with conventional fungicides, or delays in the appearance of Dollar Spot disease, thus leading to the potential for an overall reduction in annual fungicide use.

**EDGING AND BANDING APPLICATIONS FOR GROWTH REGULATION OF PERENNIAL TURFGRASS SPECIES**

Cutless MEC can be applied to turfgrass in edging and banding applications along the perimeter of lawns, landscape beds, sidewalks, curbs, parking lots, driveways, posts, mailboxes, building structures, gravestones, and fences. Apply this product in a 6 inch band width with a single nozzle sprayer. Repeat at 8 to 12 week intervals. The maximum number of annual applications is determined by the sum of the rates applied, not to exceed 3.0 lbs ai/A or 295.4 fl. oz. product/A (6-48 applications/year).

<b>TABLE 5</b> <b>Cutless MEC Edging/Banding Rates for Growth Regulation of Perennial Turfgrass</b>		
<b>Turfgrass Species</b>	<b>Fl. Oz. Cutless MEC /A</b>	<b>Fl. Oz. Cutless MEC per 1 Gallon of Water in Backpack Sprayers<sup>†</sup></b>

<b>Cool Season Turfgrasses</b>		
Bentgrass	49.2 to 98.4	1.2 to 2.5
Kentucky Bluegrass	73.8 to 147.6	1.9 to 3.7
Perennial Ryegrass	73.8 to 147.6	1.9 to 3.7
Tall Fescue	73.8 to 147.6	1.9 to 3.7
<b>Warm Season Turfgrasses</b>		
328 Hybrid Bermudagrass	36.9 to 49.2	0.9 to 1.2
419 Hybrid Bermudagrass	49.2 to 98.4	1.2 to 2.5
Common Bermudagrass	73.8 to 147.6	1.9 to 3.7
Seashore Paspalum	49.2 to 98.4	1.2 to 2.5
St. Augustinegrass	49.2 to 98.4	1.2 to 2.5
Zoysiagrass	49.2 to 98.4	1.2 to 2.5

† For backpack sprayers. Assuming a spray volume of 40 gallons per acre, one (1) gallon of spray solution will treat 2,180 linear feet with a six (6) inch band width.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used on site according to use directions or disposed of at an approved waste disposal facility.

**Nonrefillable Container Handling (rigid, 5 gallons or less):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container Handling (rigid, larger than 5 gal):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Warranty Disclaimer:** SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined

that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

**Misuse:** Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



© Copyright \_\_\_\_ SePRO Corporation

Primo MAXX® is a registered trademark of Syngenta Crop Protection.  
Cutless is a registered trademark of SePRO Corporation



[Editor's note: Version A sub-label - turfgrass uses.]  
[Base label for ALL containers]



# Cutless MEC

## Turf Growth Regulator

FOR GROWTH MANAGEMENT AND QUALITY IMPROVEMENT OF PERENNIAL COOL AND WARM SEASON TURFGRASSES ON GOLF COURSES.

### Active Ingredient:

flurprimidol:  $\alpha$ -(1-methylethyl)- $\alpha$ -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol ..... 16.0%

Other Ingredients ..... 84.0%

**TOTAL** ..... 100.0%

Contains 1.3 pounds active ingredient per one (1) U.S. gallon.

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING / AVISO

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

---

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

---

**Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used on site according to use directions or disposed of at an approved waste disposal facility.

**Nonrefillable Container Handling (rigid, 5 gallons or less):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container Handling (rigid, larger than 5 gal):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Refer to label booklet for additional Precautionary Statements and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse statements* inside label booklet.**

EPA Reg. No. 67690-51  
FPL20180517

EPA Est. No. \_\_\_\_\_  
[P/N] - \_\_\_\_\_

**SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.**

**Net Contents \_\_\_\_\_(Non-refillable)**

*[Editor's note: Version B sub-label - Ornamental Uses.]*  
*[Front label, ALL containers]*



# Cutless MEC

**FOR GROWTH MANAGEMENT AND TERMINAL GROWTH SUPPRESSION OF LANDSCAPE HEDGES, ORNAMENTAL SHRUBS, PERENNIAL GROUND COVERS AND HERBACEOUS PERENNIALS.**

**Active Ingredient:**

flurprimidol:  $\alpha$ -(1-methylethyl)- $\alpha$ -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol ..... 16.0%

**Other Ingredients** ..... 84.0%

**TOTAL** ..... 100.0%

Contains 1.3 pounds active ingredient per one (1) U.S. gallon.

**Keep Out of Reach of Children**

**WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

**Refer to label booklet for additional Precautionary Statements and Directions for Use, including Storage and Disposal.**

**Notice:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet.**

EPA Reg. No. 67690-51  
 FPL20180517

EPA Est. No. \_\_\_\_\_  
 [P/N] - \_\_\_\_\_

**SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.**

**Landscape/Ornamental Growth Regulator      Net Contents \_\_\_\_\_(Non-refillable)**

***[Editor's note: Version B sub-label - Ornamental Uses.]***  
***[Label booklet text]***

---

## **PRECAUTIONARY STATEMENTS**

---

### **Hazards to Humans and Domestic Animals**

**Keep Out of Reach of Children**  
**WARNING / AVISO**

**Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.**

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants;
- Shoes plus socks; and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### **USER SAFETY RECOMMENDATIONS**

**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment water or rinsate.

---

### **DIRECTIONS FOR USE**

---

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying this product. Use only according to label directions.

### **AVOID SPRAY DRIFT**

Applications must be made only when there is no hazard for spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. **Do not spray when wind is greater than 10 mph.** Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

## **INFORMATION FOR GROWTH MANAGEMENT AND TERMINAL GROWTH SUPPRESSION OF LANDSCAPE HEDGES, ORNAMENTAL SHRUBS, PERENNIAL GROUND COVERS AND HERBACEOUS PERENNIALS.**

Cutless MEC is a systemic plant growth regulator for use on landscape hedges, ornamental shrubs, perennial ground covers and herbaceous perennials. Treated plants require less trimming and exhibit a more compact growth form. Growth suppression is achieved by inhibition of gibberellin biosynthesis resulting in reduced internode elongation in terminal shoot growth. Smaller leaves and intensified greening occur in some species. Product absorbed by plant roots is translocated to above-ground growing points within water conducting tissues. Its movement in plants occurs in response to evaporative water loss (transpiration) from leaf surfaces.

Time required for onset of growth suppression may vary from 2 to 4 weeks for herbaceous species up to an entire year for certain woody plants. Time from application until onset of growth regulation depends on the rate of water loss from foliage and the rate of water movement through woody stems. Growth suppression will be slower to appear and of longer duration in plants with a larger mass of woody tissue. Duration of growth suppression may range from 3 to 12 months, depending upon plant species treated.

The amount of Cutless MEC required for optimum growth regulation is dependent upon plant characteristics (stem mass, height and foliage volume), environmental setting (exposure to sunlight, temperature conditions, rainfall/irrigation and soil texture), and species tolerance. Because of these interacting factors, plant response to a given application rate of this product may vary according to local conditions. Users should establish specific applications (within the ranges specified in Table 1) for different species in small scale treatments under actual use conditions prior to large scale applications.

Activation of Cutless MEC requires at least 1/2 inch of rainfall or sprinkler irrigation. Periodic rainfall or irrigation is needed to maintain uniform growth regulation activity. Avoid irrigation in excess of plant requirements.

### **Use Restrictions:**

- Use Cutless MEC only on vigorous, well-established plantings.

- Do not use this product on plants from which fruits, nuts or other commodities may be produced for human consumption or livestock feed.
- Do not apply to annual bedding plants or trees.
- Do not allow runoff to occur during initial irrigation of treated areas.
- Do not apply this product through any type of chemigation system.

**Use Precautions:**

- Avoid application when weather conditions favor movement of spray drift from treatment area.

**Note:** Nontarget plants with roots which extend into soil containing Cutless MEC may exhibit growth regulation.

**Antidote:** Gibberellin may be applied as an antidote to stimulate terminal shoot growth of over-regulated plants.

**Application Timing:**

Apply Cutless MEC as needed for shoot growth suppression of landscape hedges, ornamental shrubs, perennial ground covers and herbaceous perennials. Apply in conjunction with pruning or trimming of target species. For best results, shape or prune plants on the day of application or 1 to 5 days before or after application. Frequent trimming and loss of leaves encourages more rapid dissipation of growth regulator effects in treated plants.

**Reapplication:**

To avoid over-application of Cutless MEC, make repeat applications only after growth regulation from the prior treatment has expired (up to 1 year is required in most plants). Reapply only when growth regulation begins to decline from a desired level. Apply repeat treatments every 3 to 12 months.

**Application Rates – Ornamental Shrubs, Perennial Ground Covers, and Herbaceous Perennials**

The amount of Cutless MEC needed for an optimum growth regulation response is dependent upon the height and mass of woody stems, foliage volume, sunlight exposure, soil texture, rainfall and temperature conditions, and species tolerance. Cultural practices including frequency of irrigation, fertilization and timing of application may also affect the response to treatment. Because of these factors, plant response to a given application rate of this product may vary according to local conditions. Establish specific application rates for different species in small scale treatments under actual use conditions. Refer to Table 1, *Rate Directions for Established Ornamental Shrubs, Perennial Ground Covers and Herbaceous Perennials*, for rate directions.

**For spot treatment applications to landscape ornamentals contained in beds less than one acre (43,560 ft<sup>2</sup>), DO NOT exceed 3 lbs ai/A/year or 295 fl oz/A/year over the entire acre containing the spot treatments.**

<b>TABLE 1</b>	
<b>Rate Directions for Established Ornamental Shrubs and Hedges, Perennial Ground Covers and Herbaceous Perennials</b>	
<b>Landscape Ornamental</b>	<b>Cutless MEC (Fl. oz./1,000 ft<sup>2</sup>)</b>
Perennial Non-Woody Ground Covers and Herbaceous Perennials	1.3
	2.25
Shrubs and Hedges <sup>††</sup> , and Perennial Woody Ground Covers	2.25
	3.40
	4.50 <sup>†</sup>
	5.65 <sup>†</sup>
	6.78 <sup>†</sup>

<sup>†</sup> Rates designated for spot treatments only to dripline areas of landscape ornamentals less than 20,000 ft<sup>2</sup> over the entire acre containing the spot treatments. For spot treatments do not apply to areas greater than 20,000 ft<sup>2</sup>.

<sup>††</sup> Rates designated for spot treatment to individual plants, Do not apply more than 0.5 fl. oz. product/plant in water (35:1 water:Cutless MEC), to more than 590 plants per acre.

**Mixing and Application:**

Apply Cutless MEC in a spray volume of at least 5 gallons per 1000 ft<sup>2</sup>. Add the correct product volume and mix thoroughly. If backpack equipment is used, shake the spray tank to ensure thorough mixing and repeat every few minutes while applying to maintain a uniform spray mixture. If power equipment is used, agitate continuously.

Apply Cutless MEC spray uniformly over the treatment area. Use low pressure and coarse spray to prevent drift to nontarget plants. Sprays may be applied over the top of low growing woody plants and perennial ground covers or directed at the soil beneath shrubs or hedges.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used on site according to use directions or disposed of at an approved waste disposal facility.

**Nonrefillable Container Handling (rigid, 5 gallons or less):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



**Nonrefillable Container Handling (rigid, larger than 5 gal):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Warranty Disclaimer:** SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

**Misuse:** Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



© Copyright \_\_\_\_ SePRO Corporation

® Cutless is a registered trademark of SePRO Corporation

[Editor's note: Version B sub-label - Ornamental Uses.]  
[Base label for ALL containers]



## Cutless MEC

FOR GROWTH MANAGEMENT AND TERMINAL GROWTH SUPPRESSION OF  
LANDSCAPE HEDGES, ORNAMENTAL SHRUBS, PERENNIAL GROUND COVERS  
AND HERBACEOUS PERENNIALS.

**Active Ingredient:**

flurprimidol:  $\alpha$ -(1-methylethyl)- $\alpha$ -[4-(trifluoromethoxy)phenyl]-5-  
pyrimidinemethanol ..... 16.0%

**Other Ingredients** ..... 84.0%

**TOTAL** ..... 100.0%

Contains 1.3 pounds active ingredient per one (1) U.S. gallon.

---

### PRECAUTIONARY STATEMENTS

---

#### Hazards to Humans and Domestic Animals

**Keep Out of Reach of Children**  
**WARNING / AVISO**

**Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.**

FIRST AID	
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used on site according to use directions or disposed of at an approved waste disposal facility.

**Nonrefillable Container Handling (rigid, 5 gallons or less):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container Handling (rigid, larger than 5 gal):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, treatment area, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Refer to label booklet for additional Precautionary Statements and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse statements* inside label booklet.**

EPA Reg. No. 67690-51  
FPL20180517

EPA Est. No. \_\_\_\_\_  
[P/N] - \_\_\_\_\_

**SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.**

**Landscape/Ornamental Growth Regulator**

**Net Contents \_\_\_\_\_(Non-refillable)**

***[Text accessed through the weblink / QR code. This is NOT part of the printed label]***

---

## **TERMS AND CONDITIONS OF USE**

---

If terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label as well as the *Inherent Risks of Use* and *Limitation of Remedies* statements below are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Misuse*, *Inherent Risks of Use*, and *Limitation of Remedies*.

---

## **INHERENT RISKS OF USE**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including use under conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), the presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer and/or user of the product.

---

## **LIMITATION OF REMEDIES**

---

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use*, *Inherent Risks of Use* and *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use*, *Inherent Risks of Use* and *Limitation of Remedies* in any manner.