



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

January 7, 2015

Laurent C. Mézin, PhD  
Manager, Regulatory Affairs  
SePRO Corporation  
11550 North Meridian Street Suite 600  
Carmel, Indiana 46032-4565

Subject: Label Amendment – Addition of Alternate Brand Name and Homeowner Use  
Sublabel  
Product Name: Sonar H4C  
Alternate Brand Name: SePRO Total Pond - Prevent  
EPA Registration Number: 67690-61  
Application Date: October 8, 2014  
Decision Number: 496255

Dear Dr. Mézin:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 67690-61  
Decision No. 496255

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Sarah Meadows by phone at 703-347-0505, or via email at meadows.sarah@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending from the top of the "g" in "Montague".

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure


# Sonar® H4C

AQUATIC HERBICIDE

EPA Reg. No. 67690-61

FPL20141006 Clean

ABN: SePRO Total Pond – Prevent



Version A - Professional uses sub-label	pp. 1-15
Version B – Retail Consumer uses sub-label	pp. 16-25

**Registration Notes:** Based on EPA-Stamped label with FPL20140804.

## Label Notes:

### General Label notes:

Label changes:

1. Updated FPL date
2. Changed 'net content' to 'net weight'
3. Added a retail consumer sub-label; This will be a singular, fixed 3Lbs size (label text reflects 3lbs contents)
4. Adds an alternate brand name, included in the retail sub-label.

[Front of label booklet, ALL containers]



# Sonar® H4C

AQUATIC HERBICIDE

**AN HERBICIDE FOR MANAGEMENT OF AQUATIC VEGETATION IN FRESH WATER PONDS, LAKES, RESERVOIRS (INCLUDING INLETS AND TRIBUTARIES), POTABLE WATER SOURCES, DRAINAGE CANALS, IRRIGATION CANALS AND RIVERS.**

## Active Ingredient

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone ..... 2.7%

**Other Ingredients**..... 97.3%

**TOTAL**..... 100.0%

Contains 0.027 lb active ingredient per pound.

**Keep Out of Reach of Children**

## CAUTION / PRECAUCIÓN

**Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.**

**NOTICE:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use* and *Limitation of Remedies* inside label booklet.**

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

EPA Reg. No. 67690-61  
FPL20141006

EPA Est. No. \_\_\_\_\_  
[SPC -] \_\_\_\_\_

**SePRO Corporation • 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.**

**Net weight \_\_\_\_\_**  
**[Refillable/Nonrefillable container \_\_\_\_\_]**

[Label booklet text, ALL containers]

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing dust. Wear long sleeved shirt, long pants, shoes and socks.

## KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

### USER SAFETY RECOMMENDATIONS

- wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Follow use directions carefully so as to minimize adverse effects on non-target organisms. Trees and shrubs growing in water treated with Sonar H4C may occasionally develop chlorosis. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas.

---

## **DIRECTIONS FOR USE**

---

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all Directions Carefully Before Applying Sonar H4C.

### **PRODUCT INFORMATION**

Sonar H4C herbicide is a selective systemic aquatic herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs (including inlets and tributaries), drainage canals, irrigation canals, and rivers. Sonar H4C is a pelleted formulation containing 2.7% fluridone designed to provide enhanced numbers of pellets (greater coverage) in treated areas versus other Sonar pellet formulations at equivalent herbicide dosing. This higher density of pellets has the potential to improve herbicide contact with target vegetation in higher exchange treatment scenarios such as spot or small-partial application designs. Sonar H4C is absorbed from water by plant shoots and from hydrosol by the roots of aquatic vascular plants. It is important to maintain Sonar H4C in contact with the target plants for as long as possible. Rapid water movement or any condition which results in rapid dilution of Sonar H4C in treated water will reduce its effectiveness. In susceptible plants, Sonar H4C inhibits the formation of carotene. In the absence of carotene, chlorophyll is rapidly degraded by sunlight.

Herbicidal symptoms of Sonar H4C appear in seven to ten days and appear as white (chlorotic) or pink growing points. Under optimum conditions, 30 to 90 days are required before the desired level of aquatic weed management is achieved with Sonar H4C. Species susceptibility to Sonar H4C may vary depending on time of year, stage of growth and water movement. For best results, apply Sonar H4C prior to initiation of weed growth or when weeds begin active growth. Application to mature target plants may require an application rate at the higher end of the specified rate range and may take longer to control.

Sonar H4C is not corrosive to application equipment.

The label provides recommendations on the use of a chemical analysis for the active ingredient. SePRO Corporation recommends the use of a High-Performance Liquid Chromatography (HPLC) for the determination of the active ingredient concentration in the water. Contact SePRO Corporation to incorporate this test, known as a FastEST, into your treatment program. Other proven chemical analyses for the active ingredient may also be used. The FastEST is referenced in this label as the preferred method for the rapid determination of the concentration of the active ingredient in the water.

Application rates are provided in pounds of Sonar H4C to achieve a desired concentration of the active ingredient in parts per billion (ppb). **The maximum application rate or sum of all application rates is 90 ppb in ponds (< 10 Acres) and 150 ppb in lakes and reservoirs per annual growth cycle.** This maximum concentration is the amount of product calculated as the target application rate, NOT determined by testing the concentrations of the active ingredient in the treated water.

### **Use Restrictions**

- **Obtain Required Permits:** Consult with appropriate state or local water authorities before applying this product. Permits may be required by state or local public agencies.

- **New York State:** Application of Sonar H4C is not permitted in waters less than two (2) feet deep, except as permitted under FIFRA Section 24(c), Special Local Need registration.
- **Hydroponic Farming:** Do not use Sonar H4C treated water for hydroponic farming unless a FastEST has been run and confirmed that concentrations are less than 1 ppb.
- **Greenhouse and Nursery Plants:** Consult with SePRO Corporation for site-specific recommendations prior to any use of Sonar H4C treated water for irrigating greenhouse or nursery plants. Without site-specific guidance from SePRO, do not use Sonar H4C treated water for irrigating greenhouse or nursery plants unless a FastEST has been run and confirmed that concentrations are less than 1 ppb.

### Water Use Restrictions Following Applications With Sonar H4C (Days)

Application Rate	Drinking†	Fishing	Swimming	Livestock/Pet Consumption	Irrigation††
Maximum Rate (150 ppb) or less	0	0	0	0	See irrigation instructions below

† Note below, under *Potable Water Intakes*, the information for application of Sonar H4C within ¼ miles (1,320) feet of a functioning potable water intake.

†† Note below, under *Irrigation*, specific time frames or fluridone concentrations that provide the widest safety margin for irrigating with fluridone treated water.

- **Potable Water Intakes:** Concentrations of the active ingredient fluridone up to 150 ppb are allowed in potable water sources; however, in lakes and reservoirs or other sources of potable water, do not apply Sonar H4C at application rates greater than 20 ppb within one-fourth (1/4) mile (1,320 feet) of any functioning potable water intake. At application rates of 8-20 ppb, Sonar H4C may be applied where functioning potable water intakes are present. **NOTE: Existing potable water intakes which are no longer in use, such as those replaced by connections to potable water wells or a municipal water system, are not considered to be functioning potable water intakes.**
- **Irrigation:** For tobacco, tomatoes, peppers or other plants within the Solanaceae Family and newly seeded crops or newly seeded grasses such as overseeded golf course greens, do not use Sonar H4C treated water if concentrations are greater than 5 ppb; furthermore, when rotating crops, do not plant members of the Solanaceae family in land that has been previously irrigated with fluridone concentrations in excess of 5 ppb without consultation with a SePRO Aquatic Specialist. It is recommended that a SePRO Aquatic Specialist be consulted prior to commencing irrigation of these sites.

### Use Precautions

- **Irrigation:** Irrigation with Sonar H4C treated water may result in injury to the irrigated vegetation. Follow these precautions and inform those who irrigate from areas treated with Sonar H4C of the irrigation time frames or FastEST requirements presented in the table below. These time frames and FastEST recommendations are suggestions which should be followed to reduce the potential for injury to vegetation irrigated with water treated with Sonar H4C. Greater potential for crop injury occurs where Sonar H4C treated water is applied to crops grown on low organic and sandy soils.



Application Site	Days After Application		
	Established Tree Crops	Established Row Crops/ Turf/Plants	Newly Seeded Crops/Seedbeds or Areas to be Planted Including Overseeded Golf Course Greens
Ponds and Static Canals †	7	30	FasTEST required
Canals	7	7	FasTEST required
Rivers	7	7	FasTEST required
Lakes and Reservoirs ††	7	7	FasTEST required

† For purposes of Sonar H4C labeling, a pond is defined as a body of water 10 acres or less in size. A lake or reservoir is greater than 10 acres.

†† In lakes and reservoirs where one-half or greater of the body of water is treated, use the pond and static canal irrigation precautions.

Where the use of Sonar H4C treated water is desired for irrigating crops prior to the time frames established above, the use of a FasTEST is recommended to measure the concentration in the treated water. Where a FasTEST has determined that concentrations are less than 10 parts per billion, there are no irrigation precautions for irrigating established tree crops, established row crops or turf.

### **PLANT CONTROL INFORMATION**

Sonar H4C selectivity is dependent upon dosage, time of year, stage of growth, method of application, and water movement. The following categories, controlled, partially controlled, and not controlled are provided to describe expected efficacy under ideal treatment conditions using higher to maximum label rates. Use of lower rates will increase selectivity of some species listed as controlled or partially controlled. Additional aquatic plants may be controlled, partially controlled, or tolerant to Sonar H4C. It is recommended to consult a SePRO Aquatic Specialist prior to application of Sonar H4C to determine a plant's susceptibility to Sonar H4C.

#### **Vascular Aquatic Plants Controlled by Sonar H4C: <sup>1</sup>**

##### **Submersed Plants:**

bladderwort (*Utricularia* spp.)  
 common coontail (*Ceratophyllum demersum*) †  
 common Elodea (*Elodea canadensis*) †  
 egeria, Brazilian Elodea (*Egeria densa*)  
 fanwort, Cabomba (*Cabomba caroliniana*)  
 hydrilla (*Hydrilla verticillata*)  
 naiad (*Najas* spp.) †  
 pondweed (*Potamogeton* spp., except Illinois pondweed) †  
 watermilfoil (*Myriophyllum* spp. except variable-leaf milfoil)

##### **Floating Plants:**

azolla (*Azolla* spp.)  
 duckweed (*Lemna*, *Landoltia*, and *Spirodela* spp.)

**Shoreline Grasses:**

paragrass (*Urochloa mutica*)

<sup>1</sup> Species denoted by a dagger (†) are native plants that are often tolerant to fluridone at lower use rates. Please consult a SePRO Aquatic Specialist for recommended Sonar H4C use rates (not to exceed maximum labeled rates) when selective control of exotic species is desired.

**Vascular Aquatic Plants Partially Controlled by Sonar H4C:**

**Submersed Plants:**

Illinois pondweed (*Potamogeton illinoensis*)

limnophila (*Limnophila sessiliflora*)

tapegrass, American eelgrass (*Vallisneria americana*)

watermilfoil--variable-leaf (*Myriophyllum heterophyllum*)

**Emerald Plants:**

alligatorweed (*Alternanthera philoxeroides*)

American lotus (*Nelumbo lutea*)

cattail (*Typha* spp.)

creeping waterprimrose (*Ludwigia peploides*)

parrotfeather (*Myriophyllum aquaticum*)

smartweed (*Polygonum* spp.)

spatterdock (*Nuphar luteum*)

spikerush (*Eleocharis* spp.)

waterlily (*Nymphaea* spp.)

waterpurslane (*Ludwigia palustris*)

watershield (*Brasenia schreberi*)

**Floating Plants:**

Salvinia (*Salvinia* spp.)

**Shoreline Grasses:**

barnyardgrass (*Echinochloa crusgalli*)

giant cutgrass (*Zizaniopsis miliacea*)

reed canarygrass (*Phalaris arundinaceae*)

southern watergrass (*Hydrochloa caroliniensis*)

torpedograss (*Panicum repens*)

**Vascular Aquatic Plants Not Controlled by Sonar H4C:**

**Emerald Plants:**

American frogbit (*Limnobium spongia*)

arrowhead (*Sagittaria* spp.)

bacopa (*Bacopa* spp.)

big floatingheart, banana lily (*Nymphoides aquatica*)

bulrush (*Scirpus* spp.)

pickerelweed, lanceleaf (*Pontederia* spp.)

rush (*Juncus* spp.)

water pennywort (*Hydrocotyle* spp.)

**Floating Plants:**

floating waterhyacinth (*Eichhornia crassipes*)  
waterlettuce (*Pistia stratiotes*)

**Shoreline Grasses:**

maidencane (*Panicum hemitomon*)

**NOTE:** algae (chara, nitella, and filamentous species) are not controlled by Sonar H4C

**APPLICATION DIRECTIONS**

The aquatic plants present in the treatment site should be identified prior to application to determine their susceptibility to Sonar H4C. It is important to determine the area (acres) to be treated and the average depth in order to select the proper application rate. Do not exceed the maximum labeled rate for a given treatment site per annual growth cycle.

**Application to Ponds**

Sonar H4C may be applied to the entire surface area of a pond. For single applications, rates may be selected to provide 45 to 90 ppb to the treated water, although actual concentrations in treated water may be substantially lower at any point in time due to the slow-release formulation of this product. When treating for optimum selective control, lower rates may be applied for sensitive target species. Use the higher rate within the rate range where there is a dense weed mass, when treating more difficult to control species, and for ponds less than 5 acres in size with an average depth less than 4 feet. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the *Application Rate Calculation—Ponds, Lakes and Reservoirs* section of this label. Split or multiple applications may be used where dilution of treated water is anticipated; however, the sum of all applications should total 45 to 90 ppb and must not exceed a total of 90 ppb per annual growth cycle.

Average Water Depth of Treatment Site (feet)	Pounds of Sonar H4C per Treated Surface Acre	
	45 ppb	90 ppb
1	4.5	9
2	9	18
3	13.5	27
4	18	36
5	22.5	45
6	27	54
7	31.5	63
8	36	72
9	40.5	81
10	45	90

**Application to Lakes and Reservoirs**

The following treatments may be used for treating both whole lakes or reservoirs and partial areas of lakes or reservoirs (bays, etc.). For best results in treating partial lakes and

reservoirs, Sonar H4C treatment areas should be a minimum of 5 acres in size. Treatment of areas smaller than 5 acres or treatment of narrow strips such as boat lanes or shorelines may not produce satisfactory results due to dilution by untreated water. Rate ranges are provided as a guide to include a wide range of environmental factors, such as target species, plant susceptibility, selectivity and other aquatic plant management objectives. Application rates and methods should be selected to meet the specific lake/reservoir aquatic plant management goals.

## A. Whole Lake or Reservoir Treatments (Limited or No Water Discharge)

### Single Application to Whole Lakes or Reservoirs

Where single applications to whole lakes or reservoirs are desired, apply Sonar H4C at an application rate not to exceed 90 ppb, and in a suggested range of 16 to 90 ppb. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the *Application Rate Calculation—Ponds, Lakes and Reservoirs* section of this label. Choose an application rate not to exceed 90 ppb to meet the aquatic plant management objective. **Where greater plant selectivity is desired such as when controlling Eurasian watermilfoil and curlyleaf pondweed, choose an application rate lower in the rate range.** For other plant species, SePRO recommends contacting a SePRO Aquatic Specialist in determining when to choose application rates lower in the rate range to meet specific plant management goals. Use the higher rate within the rate range where there is a dense weed mass or when treating more difficult to control plant species or in the event of a heavy rainfall event where dilution has occurred. In these cases, a second application or more may be required; however, the sum of all applications cannot exceed 150 ppb per annual growth cycle. Refer to the section of this label entitled, *Split or Multiple Applications to Whole Lakes or Reservoirs*, for guidelines and maximum rate allowed.

Average Water Depth of Treatment Site (feet)	Pounds of Sonar H4C Per Treated Surface Acre	
	16 ppb	90 ppb
1	1.6	9
2	3.2	18
3	4.8	27
4	6.4	36
5	8	45
6	9.6	54
7	11.2	63
8	12.8	72
9	14.4	81
10	16	90
11	17.6	99
12	19.2	108
13	20.8	117
14	22.4	126
15	24	135
16	25.6	144
17	27.2	153

18	28.8	162
19	30.4	171
20	32	180

### Split or Multiple Applications to Whole Lakes or Reservoirs

To meet certain plant management objectives, split or multiple applications may be desired in making whole lake treatments. Split or multiple application programs are desirable when the objective is to use the minimum effective dose and to maintain this lower dose for the sufficient time to ensure efficacy and enhance selectivity. Under these situations, use the lower rates within the rate range. **In controlling Eurasian watermilfoil and curlyleaf pondweed and where greater plant selectivity is desired, choose an application rate lower in the rate range.** For other plant species, SePRO recommends contacting a SePRO Aquatic Specialist in determining when to choose application rates lower in the rate range to meet specific plant management goals. For split or repeated applications, the sum of all applications must not exceed 150 ppb per annual growth cycle.

**NOTE:** In treating lakes or reservoirs that contain potable water intakes and when the application requires treating within ¼ mile of a potable water intake, no single application can exceed 20 ppb. Additionally, the sum of all applications cannot exceed 150 ppb per annual growth cycle.

### **B. Partial Lake or Reservoir Treatments**

Where dilution of Sonar H4C with untreated water is anticipated, such as in partial lake or reservoir treatments, split or multiple applications may be used to extend the contact time to the target plants. The application rate and use frequency of Sonar H4C in a partial lake is highly dependent upon the treatment area. An application rate at the higher end of the specified rate range may be required and frequency of applications will vary depending upon the potential of untreated water diluting the Sonar H4C concentration in the treatment area. Use a rate at the higher end of the rate range where greater dilution with untreated water is anticipated.

#### Application Sites Greater Than ¼ Mile from a Functioning Potable Water Intake

For single applications, apply Sonar H4C at application rates not to exceed 150 ppb, and in a suggested range of 45 to 150 ppb. Split or multiple applications may be made; however, the sum of all applications cannot exceed 150 ppb per annual growth cycle. Split applications should be conducted to maintain a sufficient concentration in the target area for a period of 45 days or longer. The use of a FastEST is recommended to maintain the desired concentration in the target area over time.

#### Application Sites within ¼ Mile of a Functioning Potable Water Intake

In treatment areas that are within ¼ mile of a potable water intake, no single application can exceed 20 ppb. When utilizing split or repeated applications of Sonar H4C for sites which contain a potable water intake, a FastEST is required to determine the actual concentration in the water. Additionally, the sum of all applications cannot exceed 150 ppb per annual growth cycle.

**Application Rate Calculation — Ponds, Lakes and Reservoirs**

The amount of Sonar H4C to be applied to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

$$\text{Pounds of Sonar H4C required per treated acre} = \text{Average water depth of treatment site} \\ \times \text{Desired ppb concentration of active ingredient} \times 0.1$$

For example, the pounds per acre of Sonar H4C required to provide a concentration of 25 ppb of active ingredient in water with an average depth of 5 feet is calculated as follows:

$$5 \times 25 \times 0.1 = 12.5 \text{ pounds per treated surface acre.}$$

**NOTE:** Calculated rates may not exceed the maximum allowable rate in pounds per treated surface acre for the water depth listed in the application rate table for the site to be treated.

**Application to Drainage Canals, Irrigation Canals and Rivers****Static Canals:**

In static drainage and irrigation canals, apply Sonar H4C at typical use rates of 37 to 74 pounds per surface acre.

**Moving Water Canals and Rivers:**

The performance of Sonar H4C will be enhanced by restricting or reducing water flow. In slow moving bodies of water use an application technique that maintains a concentration of 10 to 40 ppb in the applied area for typically a minimum of 45 days. Sonar H4C can be applied by split or multiple broadcast applications or by metering in the product to provide a uniform concentration of the herbicide based upon the flow pattern. The use of a FasTEST is recommended to maintain the desired concentration in the target area over time.

**Static or Moving Water Canals or Rivers Containing a Functioning Potable Water Intake**

In treating a static or moving water canal or river which contains a functioning potable water intake, applications of Sonar H4C greater than 20 ppb must be made more than ¼ mile from a functioning potable water intake. Applications less than 20 ppb may be applied within ¼ mile from a functioning potable water intake; however, if applications of Sonar H4C are made within ¼ mile from a functioning water intake, a FasTEST must be utilized to demonstrate that concentrations do not exceed 150 ppb at the potable water intake.

**Application Rate Calculation — Drainage Canals, Irrigation Canals and Rivers**

The amount of Sonar H4C to be applied through a metering system to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

1. Average flow rate (feet per second) x average width (ft.) x average depth (ft.) x 0.9 = CFS (cubic feet per second)
2. CFS x 1.98 = acre feet per day (water movement)
3. Acre feet per day x desired ppb x 0.1 = pounds Sonar H4C required per day.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

### Container Handling

**Nonrefillable Container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity ≤ 50 pounds) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Triple rinse containers too large to shake (capacity >50 pounds) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

---

## TERMS AND CONDITIONS OF USE

---

If terms of the following *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the

buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies*.

---

## **WARRANTY DISCLAIMER**

---

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

## **INHERENT RISKS OF USE**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

---

## **LIMITATION OF REMEDIES**

---

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer*, *Inherent Risks of Use* and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or this *Limitation of Remedies* in any manner.

© Copyright \_\_\_\_ SePRO Corporation



[Base label for all containers]



# Sonar® H4C

AQUATIC HERBICIDE

## Active Ingredient

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone ..... 2.7%Other Ingredients ..... 97.3%

TOTAL ..... 100.0%

Contains 0.027 lbs active ingredient per pound.

## Keep Out of Reach of Children CAUTION / PRECAUCIÓN

---

### PRECAUTIONARY STATEMENTS

---

#### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing dust. Wear long sleeved shirt, long pants, shoes and socks.

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b> .	

**ENVIRONMENTAL HAZARDS**

Follow use directions carefully so as to minimize adverse effects on non-target organisms. Trees and shrubs growing in water treated with Sonar H4C may occasionally develop chlorosis. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

**Container Handling**

**Nonrefillable Container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Refillable Container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**See attached booklet for complete container disposal directions including triple rinsing and pressure rinsing instructions.**

**Refer to the inside of the label booklet for additional precautionary information and directions for use.**

**NOTICE:** Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* inside label booklet.**

EPA Reg. No. 67690-61  
FPL20141006

EPA Est. No. \_\_\_\_\_  
[SPC -] \_\_\_\_\_

**SePRO Corporation • 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.**

**Net weight \_\_\_\_\_**  
**[Refillable/Nonrefillable container \_\_\_\_\_]**

[Front Panel, STP ONLY containers]



# SePRO Total Pond – Prevent®

Contains Sonar® for Early-Season Floating and Submersed Weed Removal and Prevention

### Active Ingredient

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone..... 2.7%

**Other Ingredients**..... 97.3%

**TOTAL** ..... 100.0%

Contains 0.027 lb active ingredient per pound.



**Keep Out of Reach of Children**

## CAUTION

### Controls many submersed and floating aquatic weeds

- Apply early-season to prevent weeds from ever becoming a problem.
- Apply in-season to remove nuisance aquatic weeds
- Systemic, complete kill including the roots
- Easy-to-apply granular formulation

Duckweed  
 Pondweed  
 Milfoil  
 Hydrilla  
 PLUS many more

See back panel booklet for additional precautionary statements.

**Net weight 3Lbs (Non-refillable)**

[Label booklet Cover, STP ONLY containers]



## SePRO Total Pond – Prevent®



Contains Sonar® for Early-Season Floating and Submersed Weed Removal and Prevention

SePRO Total Pond - Prevent is a selective systemic aquatic herbicide for management of fresh water aquatic vegetation in ponds such as swimming ponds, fishing ponds, livestock tanks, lakes, reservoirs and potable waterbodies.

### Quick facts:

SePRO Total Pond – Prevent is powered by Sonar®, a leading aquatic herbicide for over 20 years. This easy to use formulation provides broad spectrum weed control.

- Stops photosynthesis in treated weeds
- Safe for watering livestock
- No fishing or swimming restrictions following application.

### When and where to use:

Apply early-season to prevent weeds from ever becoming a problem. Can also be used in-season to control established weeds.

- Controls most submersed weeds
- Controls most floating leaf weeds
- Results within 30 – 45 days with early application
- Season long control

### How to use:

- Apply the product directly to the water evenly around the perimeter of the waterbody or directly over areas where weeds typically grow. See inside label booklet for detailed use instructions.
- Treats up to 8,000 square feet, 4 feet deep

**Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.**

Copyright ©\_\_\_\_ SePRO Corporation

Manufactured by

SePRO Corporation; 11550 N. Meridian Street, Suite 600; Carmel, IN 46032

Customer service: 1-844-766-2727

EPA Reg. No. 67690-61

EPA Est. No. \_\_\_\_\_

[SPC -] \_\_\_\_\_

FPL20141006

Formulated and Packaged in U.S.A.

SePRO Total Pond – Prevent and Sonar are registered trademarks of SePRO Corporation

[Label booklet text, STP ONLY containers]

---

**STOP! PLEASE READ ENTIRE LABEL BEFORE USE.**

---

### DIRECTIONS FOR USE

It is a violation for Federal Law to use this product in a manner inconsistent with its labeling.

### HOW SePRO Total Pond – Prevent WORKS


SePRO Total Pond – Prevent controls unwanted aquatic vegetation by restricting the plant's ability to make food, causing them to die. This involves inhibiting development of yellow pigments that protect a plant's chlorophyll from decomposition by sunlight. As the plant's chlorophyll decreases, so does its capacity to produce food. The visual symptom of SePRO Total Pond – Prevent's action is bleaching (turning white) on the terminal bud or growing points of the plant.

### What to Expect

SePRO Total Pond – Prevent is selective and controls target weeds slowly. It lets desirable vegetation re-establish itself, and does not result in oxygen depletion or fish kill. Aquatic weeds remain under control for a full season and often longer. It takes 30 to 60 days for this product to control established weeds. You should expect full results within 90 days. You'll be able to observe the visual symptoms of SePRO Total Pond – Prevent's action as chlorosis, or bleaching, gradually appearing on the terminal buds, or growing points, of the plant. Some plants are less susceptible and may require treatment at the maximum label rate.

### WHEN TO APPLY

SePRO Total Pond – Prevent can be applied in an **Early-season Program** before many nuisance aquatic plants begin to actively grow – *or* – in an **In-season Program** to control many established nuisance aquatic plants. Regardless of your situation, for best results apply in the spring or early summer.



**Early-season Program:** Before weeds are present, or are still very small, split the treatment into 2 applications to provide optimal results and extended herbicide exposure. Find your region on the map. Apply  $\frac{1}{2}$  the appropriate rate from Table 1 or 2 based on target weed species during the range of dates indicated (or earlier if submersed plant growth is evident). Wait one month and make another application of  $\frac{1}{2}$  the indicated rate to complete the program.

**In-season Program** - If your expected treatment start date is later than the range of dates indicated for your region on the map, apply the recommended dosage indicated in Table 1 or 2 based on your target weed species in a single application.

*For Example:* A ¼-acre pond with an average depth of 4 feet, targeting control of duckweed and pondweed (Table 1). A total of 3.8 lbs. of SePRO Total Pond – Prevent is needed for the season. For the Early Season Program, apply this amount of product over two applications of 1.9 lbs. each, one month apart. For the In-season Program, apply 3.8 lbs. of product in one application.

**Susceptible Aquatic Weeds for Use Rates in Table 1.**

- Bladderwort
- Common Elodea
- Egeria, Brazilian Elodea
- Hydrilla
- Naiad
- Pondweed (except Illinois pondweed)
- Watermilfoil
- Azolla
- Duckweed
- Tapegrass, American eelgrass

**Table 1.  
SePRO Total Pond - Prevent rates (lbs.)**

Average Pond Depth (ft.)	Pond Size (ac)		
	1/8	1/4	1/2
1	0.5	0.9	1.9
2	0.9	1.9	3.8
3	1.4	2.8	5.6
4	1.9	3.8	7.5
5	2.3	4.7	9.4
6	2.8	5.6	11.3
7	3.3	6.6	13.1
8	3.8	7.5	15.0

**Susceptible Aquatic Weeds for Use Rates in Table 2.**

- Common Coontail
- Fanwort, Cabomba
- Illinois Pondweed
- American Lotus
- Parrotfeather
- Spatterdock
- Spikerush
- Waterlily
- Waterpurslane
- Watershield

**Table 2.  
SePRO Total Pond - Prevent rates (lbs.)**

Average Pond Depth (ft.)	Pond Size (ac)		
	1/8	1/4	1/2
1	0.9	1.9	3.8
2	1.9	3.8	7.5
3	2.8	5.6	11.3
4	3.8	7.5	15.0
5	4.7	9.4	18.8
6	5.6	11.3	22.5
7	6.6	13.1	26.3
8	7.5	15.0	30.0

The maximum application rate or sum of all application rates per annual growing cycle is 36 lbs./acre, 4 feet deep in ponds less than 10 acres in size.

**USE RESTRICTIONS**

**Obtain Required Permits:** Consult with appropriate state or local water authorities before applying this product. Permits may be required by state or local public agencies.

For purposes of SePRO Total Pond – Prevent labeling, calculate parts per billion (ppb) with the following conversion:

$$ppb = \text{SePRO Total Pond – Prevent per surface acre (lbs.)} \div \text{average depth (ft)} \div 0.1$$

**Water Use Restrictions Following Applications With SePRO Total Pond - Prevent**

**Drinking, Fishing and Swimming:** Drinking, fishing and swimming are allowed following application. Note below application restrictions around potable water intakes.

**Livestock/Pet Consumption:** Drinking by livestock and pets is allowed following application.

**Hydroponic Farming, Greenhouse and Nursery Plants:** Do not use SePRO Total Pond – Prevent treated water for hydroponic farming, or greenhouse or nursery plants irrigation, unless concentrations are less than 1 ppb as determined by an approved laboratory analysis. (Please contact a SePRO Total Pond Advisor for details about this service.)

**Potable Water Intakes:** Concentrations of the active ingredient up to 150 ppb are allowed in potable water sources; however, in lakes and reservoirs or other sources of potable water, do not apply Total Pond – Prevent at application rates greater than 20 ppb within one-fourth (1/4) mile (1,320 feet) of any functioning potable water intake. At application rates of 8-20 ppb, Total Pond – Prevent may be applied where functioning potable water intakes are present.

**Irrigation:** For tobacco, tomatoes, peppers or other plants within the Solanaceae Family and newly seeded crops or newly seeded grasses such as overseeded golf course greens, do not use SePRO Total Pond - Prevent treated water if concentrations are greater than 5 ppb; furthermore, when rotating crops, do not plant members of the Solanaceae family in land that has been previously irrigated with concentrations in excess of 5 ppb without consultation with a SePRO Total Pond Advisor. It is recommended that a SePRO Total Pond Advisor be consulted prior to commencing irrigation of these sites.

**USE PRECAUTIONS**

**Irrigation:** Irrigation with SePRO Total Pond - Prevent treated water may result in injury to the irrigated vegetation. Follow these precautions and inform those who irrigate from areas treated with this product of the irrigation time frames and approved laboratory analysis requirements presented in the table below. These time frames and analysis recommendations are suggestions which should be followed to reduce the potential for injury to vegetation irrigated with water treated with this product. Greater potential for crop injury occurs where SePRO Total Pond - Prevent treated water is applied to crops grown on low organic and sandy soils.

Application Site	Days After Application Before Irrigation		
	Established Tree Crops	Established Row Crops/ Turf/Plants	Newly Seeded Crops/Seedbeds or Areas to be Planted Including Overseeded Golf Course Greens
Ponds and Static Canals <sup>†</sup>	7	30	approved laboratory analysis required <sup>†††</sup>
Lakes and Reservoirs <sup>††</sup>	7	7	approved laboratory analysis required <sup>†††</sup>

<sup>†</sup> For purposes of SePRO Total Pond - Prevent labeling, a pond is defined as a body of water 10 acres or less in size. A lake or reservoir is greater than 10 acres.

<sup>††</sup> In lakes and reservoirs where one-half or greater of the body of water is treated, use the pond and static canal irrigation precautions.

<sup>†††</sup> Please contact a SePRO Total Pond Advisor for details about this service.

Where the use of SePRO Total Pond - Prevent treated water is desired for irrigating crops prior to the time frames established above, the use of an approved laboratory analysis is recommended to measure the concentration in the treated water. (Please contact a SePRO Total Pond Advisor for details about this service.) Where such an analysis has determined that concentrations are less than 10 parts per billion, there are no irrigation precautions for irrigating established tree crops, established row crops or turf.

For more information on how to obtain a FastEST sample kit, contact a SePRO Total Pond Advisor TOLL FREE at 1-844-766-2727 or email [service@seprototal.com](mailto:service@seprototal.com)



## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing dust. Wear long sleeved shirt, long pants, shoes and socks.

## KEEP OUT OF REACH OF CHILDREN CAUTION

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC at 1-800-535-5053</b> .	

### **ENVIRONMENTAL HAZARDS**

Follow use directions carefully so as to minimize adverse effects on non-target organisms. Trees and shrubs growing in water treated with SePRO Total Pond - PREVENT may occasionally develop chlorosis.

<b>STORAGE AND DISPOSAL</b>
<p><b>Pesticide Storage:</b> Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, contain material and dispose as waste.</p> <p><b>Container Handling</b>  <b>Nonrefillable Container. DO NOT reuse or refill this container.</b>  <b>If empty:</b> Place in trash or offer for recycling if available. <b>If partly filled:</b> Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>

**Notice:** Research and testing have determined that the “directions for use” are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the buyer assumes responsibility for lack of performance or safety if not used according to the directions. To the extent consistent with applicable law, Manufacturer makes no other express or implied warranty of fitness or merchantability nor any other express or

**implied warranty. In no case shall Manufacturer be liable for consequential, special or indirect damages resulting from the use, handling or application of this product.**

---



**Need help? A SePRO Pond Advisor can remotely and accurately measure the size of your pond and discuss a personalized strategy to achieve your management objectives. Call us TOLL FREE at 1-844-766-2727 or email [service@seprototal.com](mailto:service@seprototal.com)**

SePRO Total Pond - Prevent is a registered trademark of SePRO Corporation.  
Sonar is a registered trademark of SePRO Corporation.

Copyright ©\_\_\_\_\_ SePRO Corporation

[Base label for STP ONLY containers]



## SePRO Total Pond – Prevent®

Contains Sonar® for Early-Season Floating and Submersed Weed Control

### Active Ingredient

fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone ..... 2.7%

**Other Ingredients** ..... 97.3%

**TOTAL** ..... 100.0%

Contains 0.027 lbs active ingredient per pound.

## Keep Out of Reach of Children

## CAUTION

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing dust. Wear long sleeved shirt, long pants, shoes and socks.

**Refer to the inside of the label booklet for additional precautionary information and directions for use.**

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC</b> at <b>1-800-535-5053</b>.</p>	

**STORAGE AND DISPOSAL**

**Pesticide Storage:** Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, contain material and dispose as waste.

**Container Handling**

**Nonrefillable Container. DO NOT reuse or refill this container.**

**If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

SePRO Total Pond - Prevent is a registered trademark of SePRO Corporation.  
Sonar is a registered trademark of SePRO Corporation.

Copyright ©\_\_\_\_ SePRO Corporation

**SePRO Corporation** 11550 North Meridian Street, Suite 600, Carmel, IN 46032

1-844-766-2727

EPA Reg. No. 67690-61  
FPL20141006

EPA Est. No. \_\_\_\_\_  
[SPC -] \_\_\_\_\_

**SePRO Corporation • 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.**

**Net weight 3Lbs (Non-refillable)**