67702-4

1-6-1999

1/2

# NEU1160 Vegetable Oil Insecticide

For Use in Homes, Home Gardens and Home Greenhouses

Active Ingredient	By Wt.
Canola Oil	96%
Inert Ingredients	<u>4%</u>
Total	100%

This product contains 7.368 lbs. of active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN

ACCEPTED

JAN 6 1999

Under the Federal Insecticides. Fungicide. and Rodenticide Art. as amended, for the posticide registered under 67702-4

# CAUTION

FIRST AID

IF SWALLOWED; Ingestion may irritate the digestive tract, each medical attention if symptoms persist or if material is ingested in large quantities. Call a physician for advice.

IF ON SKIN: Wash with plenty of soap and water, Get medical attention if trritation persists

PRECAUTIONARY STATEMENTS - HOUSEHOLD

Hazards to Humans and Domestic Animals: Harmful If swallowed or absorbed through skin, Avoid contact with skin, eyes or clothing. Wash thoroughly with some and water efter handling.

Environmental Hazards; Do not apply directly to water, Do not contaminate water when disposing of equipment washwaters of rineste

DIRECTIONS FOR USE - HOUSEHOLD

It is a violation of Federal law to use this product in a menner inconsistant with its labeling.

## STORAGE AND DISPOSAL - HOUSEHOLD

STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animats.

DISPOSAL: Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

Please read entire label. Use only as directed. Shake well before each use. Thoroughly wet all parts of the pasts and plants. Repeat treatment in 1 - 2 weeks if needed.

#### DIRECTIONS FOR MIXING - HOUSEHOLD

For 2% formulation; mix 4 teaspoons of concentrate with 1 quart water (or 2 fluid oz, concentrate with 3 quarts water). Shake well before each use and stir or shake diluted spray frequently during use. Apply one quart of formulation for 3 square yards.

#### PESTS/SITES AND APPLICATION INSTRUCTIONS-HOUSEHOLD

Apply at a rate of 2% formulation. Using hand-held trigger sprayers, hand sprayers or backpack sprayers, thoroughly spray insect posts with diluted product until plants are wet. Repeat treatment in 1-2 weeks if needed, Apply one quart of formulation for 3 square yards.

Controls:		
Adelgids	Aphids	Mites
Leathoppers	Meslybugs	Scale
Phylioxena	Psyttida .	Immature plant bugs
Sawfly larvae	Whitefly	

72

F	OΓ	UBB	on

VEGETABLES	FRUITS & NUTS	ORNAMENTAL & SHADE TREES
Beens	Apples	Birch
Cabbage	Cherrise	Everproens
Corn	Citrus	Holly
Curcurbits	Figs	Oek
Melone	Grapes	
Peas	Peers	
Potatoes	Nüts	•
Soybeans	Raspberries	
Tomatoes	Strawberries	

#### DIRECTIONS FOR DORMANT USE - HOUSEHOLD

Controls egg stages of pasts present in dormant season and scale insects, mites and mealybugs on dormant shrubs (such as roses), evergreens, woody plants, fruit trees and shade trees. For dormant use apply a 2% formulation, mix 4 teaspoons of concentrate with 1 quart water (or 2 fluid ounces concentrate with 3 quarts water). Thoroughly wat the dormant plants or trees.

#### **USE PRECAUTIONS - HOUSEHOLD**

Oil removes the glaucous (blue) bloom from such evergreene as Colorado blue spruce and Koster spruce. Use caution on blooms and buds and on oil sensitive plants such as: Chamsecypenis, Cryptomeria, Japanese Holly, Red Cedar and Smoke Tree. If concerned about sensitivity of plants, test spray a portion of the plant and well several days to determine if burn occurs before spraying larger areas.

Do not combine with sulfur and do not follow a previous application of sulfur or apply sulfur following an application of this material until sufficient time has elapsed for safety (at least 30 days).

When used in combination with or immediately before or after spraying, this product may increase the phytotodoxy of dinitro compounds, or fungicides such as Captan, Foipet, Karathene and Morestan, or copper compounds. If in doubt, treat a small area with a combination spray and well several days to determine if burn occurs.

Do not use NEU1160 spray when temperatures are above 90°F. When temperatures drop below 32°F during the night, do not spray tree and vine crops until daytime temperatures have warmed to a point above 35°F and crops are free from moisture due to metting frost or heavy daws.

Use of the NEU1160 on citrus after October 1 may increase the tree's susceptibility to cold damage.

#### NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on this label and is researcably fit for purposes stated on this label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not researcably foreseeable to seller. Buyer assumes all risk of any such use. Seller makes no other warranties, either expressed or implied.

- Dorment and growing season insect spray.
- Kills all stages of insects, including eggs.
- Based on plant oits.
- Residual protection against mites.
- Dose not persist in the environment.
- · Contact insecticide, ovicide.
- Insecticide from plants for plants.
- Acts as a repellent (ovipositional and feeding deterrent).
- Use to control insects, mites, scale.
- · For flowers, fruits and vegetables, houseplants, trees.
- For use indoors, autdoors and in greenhouses.

Registrant:

W. Neudorff GmbH KG, Postfach 1209, an der Mühle 3,

D-31660 Emmerthal, Germany

EPA Registration #: 67702-4

EPA Establishment # 67702-WG-1

Gallons

**NET CONTENTS:** 

DEC-22-1998 16:11

7037595548

99%