

67702-24

7/28/2011

10 of 21



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr. Walter G. Talarek
W. Neudorff GmbH KG
1008 Riva Ridge Dr
Great Falls, VA 22066-1620

JUL 28 2011

JUL 28 2011

Subject: Bug-N-Sluggo Insect, Slug, and Snail Bait, EPA Reg. No. 67702-24
Date of Registrant Submission: April 29, 2011
Decision: 448819

Dear Mr. Talarek:

The labeling referred to above, submitted in connection with registration under the Federal insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

Should you wish to retain a reference to the company's website on your label, please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance. As an alternative, you may refer consumers to the company's phone number or email address.

Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records. If you have any questions regarding this letter, please contact Samantha Hulkower at (703) 603-0683.

Sincerely,


Mark Suarez
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure: Copy of Label Stamped "Accepted"

MASTER LABEL

**BUG-N-SLUGGO®
INSECT, SLUG AND SNAIL BAIT**

Active Ingredient:	By weight
Iron phosphate.....	0.97%
Spinosad (a mixture of spinosyn A and spinosyn D)	0.07%
Other Ingredients:	<u>98.96%</u>
Total	100.0%



SUBLABEL A: Home and Garden Use
SUBLABEL B: Commercial Agriculture Use

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Registration #67702-24

EPA Establishment #

Registrant: W. Neudorff GmbH KG
An der Muhle 3, Postfach 1209
D-31860 Emmerthal, Germany
www.neudorff.com

[in the alternative, the distributor's telephone number and website address may be used]

Home and Garden Use: NET WEIGHT 1oz, 13oz, 20oz, 26oz, 30oz, 0.5lb, 1 lb, 1.25lbs,
1.5 lbs, 2lbs, 2.5lbs, 3lbs, 3.2lbs, 4 lbs, 5lbs, 5.25lbs, 5.5lbs, 6lbs, 7lbs, 8lbs, 10lbs, 20lbs,
25lbs, 40 lbs, 1lb. 4oz., 1lb. 14oz., 2lb. 8oz., 2116 lbs (960kg)

Commercial Agriculture Use: NET WEIGHT 5lbs, 10lbs, 20lbs, 25lbs, 40lbs, 45lbs,
50lbs, 55lbs (25 kg), 1984lbs (900 kg), 2011lbs (912 kg), 2116 lbs (960kg)

SUBLABEL A: Home and Garden Use**BUG-N-SLUGGO®
INSECT, SLUG AND SNAIL BAIT****For Outdoor Use Around Residential Sites**

Active Ingredient:	By weight
Iron phosphate.....	0.97%
Spinosad (a mixture of spinosyn A and spinosyn D)	0.07%
Other Ingredients:	<u>98.96%</u>
Total	100.0%

NET WEIGHT 1oz, 13oz, 20oz, 26oz, 30oz, 0.5lb, 1 lb, 1.25lbs, 1.5 lbs, 2lbs, 2.5lbs,
3lbs, 3.2lbs, 4 lbs, 5lbs, 5.25lbs, 5.5lbs, 6lbs, 7lbs, 8lbs, 10lbs, 20lbs, 25lbs, 40 lbs, 1lb.
4oz., 1lb. 14oz., 2lb. 8oz., 2116 lbs (960kg)

KEEP OUT OF REACH OF CHILDREN**CAUTION****FIRST AID**

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes.
Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or
doctor, or going for treatment. You may also contact the National Pesticide Information
Center hotline at 1-800-858-7378 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Caution. Causes moderate eye irritation.
Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling
and before eating, drinking, chewing gum, or using tobacco.

Environmental Hazards: To protect the environment, do not allow pesticide to enter or
run off into storm drains, drainage ditches, gutters or surface waters. Applying this
product in calm weather when rain is not predicted for the next 24 hours will help to
ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping

any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PESTS CONTROLLED: earwigs, cutworms, sowbugs, pillbugs, crickets, ants (excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants), slugs, snails

APPLICATION DIRECTIONS:

Scatter the bait granules on the soil around or near the plants to be protected with a suitable hand or power operated spreader (gravity or rotary) to ensure uniform coverage over the treatment area. Apply bait evenly at approximately 0.5 to 1 lb per 1000 square feet, or for smaller areas, 0.075 to 0.15 oz, or 0.5 to 1 teaspoon, per square yard. Do not place in piles. For slugs and snails also scatter the bait around the perimeter of the area to provide a protective "barrier" for slugs and snails entering the area. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. Do not use food utensils such as tablespoons, cups, etc. for food purposes after pesticide use.

Do not apply more than 3 times in any 30 day period. Apply at the higher rate if the infestation is severe. Re-apply after heavy rain or watering. Do not apply within 3 days of harvest.

For seedlings spread the bait around the base of the stem. For trees or shrubs, spread the bait around the base of the tree. For small plots, treat around the base of the plants to be protected. In greenhouses, scatter the bait in the pots of plants being damaged or around pots on greenhouse benches.

Evening is the best time to apply the bait as pests travel and feed mostly by night or early morning. Treat all likely areas of infestation, especially around the perimeter of garden plots, for slugs and snails because these pests travel into plant areas from daytime refuges.

USE SITES:

Vegetables: artichokes (globe), asparagus, beans, beets, broccoli, Brussels sprouts, cabbage, cantaloupe, carrots, cauliflower, corn, cucumbers; eggplants, garlic, lettuce, onions, peas, peppers, potatoes, radishes, rutabagas, spinach, soybeans, squash, Swiss chard, tomatoes and turnips; **Fruits:** apples, avocados, apricots, cherries, grapes, melons, peaches, plums, citrus, pears; **Berries:** strawberries, blueberries, loganberries; **Outdoor Ornamentals; Greenhouses (Non-Commercial); Lawns; Turf.**

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

CONTAINER HANDLING:

Nonrefillable. Do not reuse or refill this container.:

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

BATCH CODE:

The registrant may use one or both of these two optional statements, either:

"WARRANTY

To the extent consistent with applicable law, seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law, seller makes no other warranties, either expressed or implied."

or

OUR COMMITMENT TO YOUR SATISFACTION: If you are not happy with the way this product works in your garden, please let us know. We will be happy to replace the product or refund your money. Simply send your name, address and proof of purchase to [insert name and address of supplemental registrant]."

[The following claims and product information, may or may not be presented on the product's label and labeling:

-Quick Facts

-Product Facts

-What it does: Kills insects, snails and slugs

-Where to Use: For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens and in greenhouses

-Read the Label First!

-**NOTE:** This package is sold by weight. Contents may have settled during shipment.



-For Organic Gardening

-US Patent number x,xxx,xxx.

- Patent Pending
- This container is made from XX% recycled materials.
- The highly compressed granules (pellets) are easy to use, clean to handle and economical.
- Patented technology.
- Patented snail & (and) slug killer.
- Patented snail, slug and listed insect killer
- Patented insect, snail and slug killer
- Patented formula
- Easy-to-use (ready-to-use) (RTU) granular (pellet) formulation.
- Easy to apply granules (pellets).
- Kills snails & (and) slugs.
- Pellets
- Kills slugs, snails and listed insects.
- SLUG, SNAIL AND INSECT BAIT
- SLUGGO plus insect killer
- Slug and bug control
- Slug and bug killer
- INSECT, SLUG AND SNAIL (BAIT)(PELLETS)
- bug and slug control
- bug and slug killer
- Kills (earwigs,) (cutworm,) (sowbugs,) (pillbugs,) (crickets,) (ants, excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants)) (slugs,) (snails,)
- Spinosad is derived from a naturally occurring soil dwelling bacterium.
- Spinosad is biologically derived from *Saccharopolyspora spinosa*, a naturally occurring soil organism.
- Spinosad is derived from a soil dwelling bacterium.
- Treats (will treat) x,xxx sq. ft.
- 1 lb treats 2000 ft²
- Effective for up to 4 weeks.
- Lasts up to 4 weeks.
- Remains effective for up to 4 weeks.
- Proven snail & (and) slug & (and) insect killer (kill, control).
- Convenient. Easy-to-use. Requires no mixing, spraying, or special applicators.
- SATISFACTION GUARANTEED.
- Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.
- Can be used in vegetable gardens.
- Ideal for vegetable gardens.
- For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens, and in greenhouses (non-commercial).
- Use around flowers, gardens, and fruit trees.
- Lawn insecticide
- Can be used on lawns
- Home and Garden

- Read Entire Container Label Before Using This Product.
- Unconditionally guaranteed by [insert name of supplemental registrant].
- Baits and Kills-For household home garden use.
- The slugs and snails controlled by this product: *Deroceras reticulatum* (Field slug), *Deroceras laeve* (Smooth slug), *Arion subfuscus* (Dusky slug), *Arion circumscriptus* (Gray garden slug), *Arion hortensis* (Black field slug), *Arion rufus* (Large red slug), *Arion ater* (Large black slug), *Limax flavus* (Spotted garden slug), *Limax tenellus* (Slender slug), *Ariolimax columbianus* (Banana slug), *Helix* spp., *Helicella* spp., and *Cepaea* spp.
- Kills a wide variety of slugs and snails
- Effective against a wide variety of slugs, snails and listed insects.
- Effective against a wide variety of insects, slugs and snails.
- Lures slugs, snails and insects from hiding places and plants.
- Lures insects, slugs and snails from hiding places and plants.
- No slime trails
- No messy cleanup
- Dead snails, no slime trails
- Attracts and kills slugs, snails and insects
- Attracts and kills insects, slugs and snails
- Attracts and kills-For control (or suppression) of (earwigs), (cutworm), (sowbugs), (pillbugs), (crickets), (ants, excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants)), (slugs), (snails)
- Contains Spinosad
- Contains Ferramol™ Slug and Snail Bait, a trademark of W. Neudorff GmbH KG
- Made with Ferramol™ Slug and Snail Bait, a trademark of W. Neudorff GmbH KG
- Ferramol™ is a trademark of W. Neudorff GmbH KG, Germany.
- Sluggo® is a registered trademark of W. Neudorff GmbH KG, Germany
- www.neudorff.com
- Manufactured under a license of W. Neudorff GmbH KG, Germany.



-For more information on Bug-N-Sluggo and other [sub registrant] products call 1-xxx-[sub registrant]-x or visit our web site at www. [sub registrant].com

-[Slugs and snails] [They] favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra, lawns, ivy patches, and other ground cover where they obtain shelter by day.

Registrant: W. Neudorff GmbH KG, Postfach 1209, An der Mühle 3,
 D-31860 Emmerthal, Germany
 011-49-5155-62421

EPA Registration #67702-24

EPA Establishment #67702-DEU-2

SUBLABEL B: Commercial Agriculture Use

**BUG-N-SLUGGO®
INSECT, SLUG AND SNAIL BAIT**

Active Ingredient:	By weight
Iron phosphate.....	0.97%
Spinosad (a mixture of spinosyn A and spinosyn D)	0.07%
Other Ingredients:	<u>98.96%</u>
Total	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Registration #67702-24

EPA Establishment # 67702-DEU-

NET WEIGHT 5lbs, 10lbs, 20lbs, 25lbs, 40 lbs, 45lbs, 50lbs, 55lbs (25 kg), 1984lbs
(900 kg), 2011lbs (912 kg), 2116lbs (960kg)

FIRST AID
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Hotline Number
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticide Information Center at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: www.npic.orst.edu) [alternately insert subregistrants toll free number and hours of operation]

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE) Requirements: Applicators and other handlers must wear: long-sleeved shirts and long pants; and shoes plus socks. Follow manufacturer's instructions for maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This product is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a manner that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency representative responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and the restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

There is a restricted-entry interval of 4 hours for this product.

COMMERCIAL AGRICULTURE

HOW TO APPLY: Scatter the bait granules on the soil around or near the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply the higher rates if the infestation is severe or if the area is heavily watered or after long periods of heavy rain. Reapply as the bait is consumed or at least every 2-4 weeks. Do not apply more than 3 times in any 30 day period.

Do not place in piles. The product works best when soil is moist but with little or no standing water. See specific directions for different use sites.

PESTS CONTROLLED: earwigs, cutworms, sowbugs, pillbugs, crickets, ants (excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants), slugs, snails

Bug-N-Sluggo contains 0.0007lbs of spinosad per lb. Use the following table to assist in calculating applied lbs spinosad per acre.

Application Rate Reference Table:

Application Rate of Bug-N-Sluggo	Spinosad Equivalent (lb ai/acre)
20 lbs/acre	0.014
25 lbs/acre	0.0175
30 lbs/acre	0.021
35lbs/acre	0.0245
40lbs/acre	0.028
44lbs/acre	0.0308

USE SITES:

VEGETABLES

Scatter the bait around the perimeter of the field or greenhouse vegetable plantings at the rate of 20-44 lbs per acre (0.5-1 lb per 1,000 square feet) to provide a protective "barrier" for pests entering the vegetable plantings. If pests are inside the rows, then scatter the bait on the soil around the plants and between the rows.

Artichoke	Do not apply within 2 days of harvest. Do not make applications less than 7 days apart. Do not make more than 4 applications per crop. Do not apply more than a total of 0.33 lb ai spinosad (including foliar uses) per crop.
Asparagus	For post harvest protection of ferns only. Do not apply within 60 day of spear harvest. Do not make applications less than 4 days apart. Do not make more than 3 applications per crop. Do not apply more than a total of 0.28 lb a.i. spinosad per acre per season. Do not feed treated ferns to meat or dairy animals.

Root and Tuber Vegetables:

Potatoes, ginger, sweet potatoes, yams	Do not apply within 7 days of harvest. Do not make applications less than 7 days apart. Do not make more than 4 applications per crop. Do not apply more than a total of 0.33 lb ai spinosad (including foliar uses) per crop.
Carrots and Parsnips	Do not apply within 3 days of harvest. Do not make applications less than 5 days apart. Do not make more than 4 applications per calendar year. Do not apply more than a total of 0.3 lb ai spinosad (including foliar uses) per acre per crop.
Beets and Sugar Beets	Do not apply within 3 days of harvest. Do not make applications less than 7 days apart. Do not make more than 4 applications per crop. Do not apply more than a total of 0.33 lb ai spinosad (including foliar uses) per crop.

Bulb Vegetables

dry bulb onion, garlic, great-headed elephant garlic, green onion, leek, shallot, Welch onion	Do not apply within one day of harvest. Do not make applications less than 4 days apart. Do not apply more than a total of 0.45 lb ai spinosad (including foliar and seed uses) per
---	---

	acre per crop. Do not make more than 5 applications per year.
--	---

Leafy Vegetables (except Brassicas) and leaves of root and tuber and legume vegetables

arugula, beets, celery, peas, lettuce, rhubarb, spinach, Swiss chard, turnip greens and watercress	Do not apply within one day of harvest for leafy vegetables and watercress; 3 days for leaves of root, tuber and legume vegetables. Do not make applications less than 4 days apart. Do not make for than 6 applications per year. Do not apply more than a total of 0.45lb ai spinosad (including foliar and seed uses) per acre of crop. Do not apply to seedling leafy crops grown for transplant within a greenhouse or shade house.
--	--

Cole Crops

broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi,	Do not apply within one day of harvest. Do not make applications less than 4 days apart. Do not make for than 6 applications per year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre of crop. Do not apply to seedling cole crops grown for transplant.
--	---

Legume Vegetables

Succulent and dried beans and peas	Do not make applications less than 5 days apart. Do not make more than 6 applications per crop. Succulent beans and peas: do not apply within 3 days of harvest. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per season. Dried beans and peas: do not apply within 28 days of harvest. Do not apply more than a total of 0.188lb ai spinosad per acre per season.
Soybean	Do not apply within 28 days of harvest. Do not make applications less than 4 days apart. Do not make more than 4 applications per calendar year. Do not apply more than a total of 0.186lb ai spinosad (including foliar uses) per acre per year. Do not feed treated forage or hay to meat or dairy animals.

Cucurbit Vegetables

cucumber, edible gourds, melons, muskmelons, cantaloupe, honeydew, pumpkin, summer squash, watermelon, and winter squash	Do not make more than 6 applications per crop. Do not apply within one day of harvest for cucumber; 3 days for other cucurbits. Do not make applications less than 5 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar and seed uses) per acre per season.
--	---

Fruiting Vegetables

eggplant, groundcherry, pepino, pepper (except black), tomatillo, tomato and okra	Do not apply within one day of harvest. Do not make applications less than 4 days apart. Do not make more than 6 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop. Do not apply to seedling fruiting vegetables and okra grown for transplant within a greenhouse, shade house, or field plot.
---	--

Corn

Corn – field corn	Do not apply within 28 days of harvest for grain or fodder; 7 days for forage. Do not make more than 3 applications per calendar year. Do not apply more than a total of 0.188lb ai spinosad (including foliar uses) per acre per year.
Corn – sweet corn	Do not apply within one day of harvest for grains; 7 days for forage, 28 days for fodder. Do not make more than 6 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year.

FRUITS INCLUDING CITRUS, AND SMALL FRUITS

Apply at 0.15 oz., or 1 level teaspoon, per square yard, in a 6 inch circular band around the base of the plants to be protected in the field or greenhouse. For older trees, spread the bait around the base of the trees to intercept pests traveling to the trunk. Apply the bait at 20-44 lbs. per acre (0.5-1 lb per 1,000 square feet) in orchards, using standard

fertilizer granular spreaders. For seedlings spread the bait around the base of the stem. Use the higher rates for heavy infestations.

Banana and plantain	Do not apply within 8 weeks of harvest. Do not make applications less than 7 days apart. Do not make more than 6 applications per calendar year or 4 applications per crop. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.
Fig	Do not apply within 7 days of harvest. Do not make applications less than 7 days apart. Do not make more than 4 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad per acre per crop.
Grape	Do not apply within 7 days of harvest. Do not make applications less than 5 days apart. Do not make more than 5 applications per calendar year. Do not apply more than a total of 0.45 lb ai spinosad per acre per season west of the Rocky Mountains, and no more than 0.36 lb ai spinosad per acre per season east of the Rocky Mountains.
Tropical tree fruits (avocado, guava, mango, papaya, passion fruit)	Do not apply within one day of harvest. Do not make more than 2 applications per calendar year. Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.

Citrus Fruits

grapefruit, lemons, limes, oranges, tangerines	Do not apply within one day of harvest. Do not make applications less than 6 days apart. Do not make more than 3 applications per calendar year. Do not apply more than a total of 0.45 ai spinosad (including foliar uses) per acre per crop.
--	--

Pome Fruits

apples, crabapples, loquat, mayhaw, pears, quince	Do not apply within 7 days of harvest. Do not make applications less than 10 days apart. Do not make more than 4 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.
---	--

Stone Fruits

cherries, plums, prunes	Do not apply within 7 days of harvest. Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year.
peaches, nectarines, apricots	Do not apply within 14 days of harvest for apricots or within 1 day of harvest for nectarines and peaches. Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year.

BERRIES

Spread the bait around the perimeter of the field or greenhouse to intercept pests migrating toward the berries. Use a rate of 20-44 lbs. per acre (0.5-1 lb per 1,000 square feet) and scatter by hand or with granular spreaders. If pests are already in the plots, then carefully spread bait between the furrows near the base of the plants.

Crop	Restrictions
Bushberries (blueberry, currant, elderberry, gooseberry, huckleberry, juneberry, ligonberry, salal)	Do not apply within 3 days of harvest. Do not make applications less than 6 days apart. Do not make more than 6 applications per calendar year or more than 3 applications per crop. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.
Caneberries (blackberry, loganberry, red and black raspberry)	Do not apply within one day of harvest. Do not make applications less than 5 days apart. Do not make more than 6 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.
Cranberry	Do not apply within 21 days of harvest. Do not make applications less than 7 days apart. Do not make more than 6 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.
Strawberry	Do not apply within one day of harvest. Do not make applications less than 5 days apart. Do not make more than 5 applications per calendar year. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.

TREE NUTS, PEANUTS AND PISTACHIOS

Use a rate of 20-44 lbs. per acre (0.5-1 lb per 1,000 square feet) and scatter by hand or with granular spreaders. For seedlings spread the bait around the base of the stem. For older trees, spread around the base of the trees to intercept pests traveling to the trunk. Use the higher rates for heavy infestations.

Crop	Restrictions
Peanut (not for use in California)	Do not apply within 3 days of nut harvest or 14 days of forage. Do not make applications less than 7 days apart. Do not make more than 3 applications per calendar year. Do not apply more than a total of 0.28lb ai spinosad (including foliar uses) per acre per crop. Do not allow grazing of crop residue or harvest of crop residue for hay until 14 days after the last application.
Tree nuts (almond, cashew, chestnut, filbert, macadamia nut, pecan, walnut); Pistachios	Do not apply within 1 day of harvest. Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.

HERBS

At the seedling and later stages, apply the bait between the rows and around the perimeter of the field or greenhouse. Use the higher dosage for heavy infestations.

Application Rate: - Use 20-44 lbs/acre (0.5-1 lb/1000 sq ft).

Crop	Restrictions
angelica, balm, basil, borage, burnet, chamomile, catnip, chervil (dried), chive, clary, costmary, cilantro, curry, dillweed, horehound, hyssop, lavender, lemongrass, lovage, marigold, marjoram, nasturtium, parsley, pennyroyal, rosemary, rue, sage, savory, sweet bay, tansy, tarragon, thyme, wintergreen, woodruff and wormwood	Do not apply within one day of harvest. Do not make applications less than 5 days apart. Do not make more than 5 applications per calendar year or more than 3 applications per crop. Do not apply more than a total of 0.47lb ai spinosad (including foliar uses) per acre per crop.
Peppermint and spearmint	Do not apply within 7 days of harvest. Do not make applications less than 4 days apart. Do not make more than 4 applications per calendar year or more than 3 applications per crop. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per crop.

GRASS FORAGE, FODDER, HAY AND CEREAL GRAINS

When pests are detected, scatter the bait by hand or with a granular spreader at a rate of approximately 0.23 to 1 lb. per 1000 square feet (10-44 lbs. per acre). Use the higher rate for heavy infestations.

Small Cereal Grains

Crop	Restrictions
barley, buckwheat, grain amaranth, milo, oats, pearl millet, proso millet, rye, sorghum, triticale, wheat	Do not use within 3 days of harvest for forage, fodder or hay. Do not apply within 21 days of harvest for grain or straw. Do not make applications less than 4 days apart. Do not make more than 3 applications per calendar year. Do not apply more than a total of 0.28lb ai spinosad (including foliar uses) per acre per year.

Grass Forage, Fodder, Hay

Grass crops, grass grown for pastures and rangeland	Do not use within 3 days of harvest for hay or fodder. No preharvest interval for forage. Do not make more than 6 applications per season. Do not apply more than a total of 0.186lb ai spinosad per acre per season.
---	---

GRASS GROWN FOR SEED PRODUCTION

When pests are detected, scatter the bait by hand or with a granular spreader at a rate of approximately 0.23 to 1 lb. per 1000 square feet (10-44 lbs. per acre). Use the higher rate for heavy infestations. Do not make more than 6 applications per season. Do not apply more than a total of 0.186lb ai spinosad per acre per season.

OUTDOOR ORNAMENTALS

Scatter bait in a 6 inch circular band around the base of the plants to be protected at 0.15 oz., or 1 level teaspoon, per square yard. If plants are next to a grassy area, spread the bait between the ornamentals and the grass. Pests traveling to the plants will encounter the bait before reaching the plant. In these situations, scatter the bait around the perimeter of the plot at 1 lb. per 1000 square feet to intercept pests traveling to the plot. Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year.

GREENHOUSE ORNAMENTALS

Where insects, slugs or snails are a problem in greenhouses, scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply about ½ teaspoon per 9 inch pot. Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year. Do not apply more than 10 times in 12 months.

TREE FARMS AND PLANTATIONS

Use the bait on tree farms and plantations. Spread the bait at a rate of 0.5 to 1 lb. per 1000 square feet (20-44 lbs. per acre). Use the higher rate for heavy infestations. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year.

OUTDOOR CONTAINER-GROWN NURSERY PLANTS

Where pests are a problem in outdoor nurseries, scatter the bait in the plant containers at the rate of 1 tablespoon per container of plants being damaged, or scatter around the soil near the containers at the rate of 1 teaspoon per square yard (24-44 lbs. per acre). Do not make applications less than 7 days apart. Do not apply more than a total of 0.45lb ai spinosad (including foliar uses) per acre per year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty metal drum, plastic bag, box or plastic tote into application equipment. Then dispose of empty metal drum, plastic bag, box or plastic tote in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BATCH CODE

The registrant may use any or none of these optional statements:

"WARRANTY

To the extent consistent with applicable law, the seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with the directions for use. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law, seller makes no other warranties, either expressed or implied.

OR

OUR COMMITMENT TO YOUR SATISFACTION: If you are not happy with the way this product works in your garden, please let us know. We will be happy to replace the product or refund your money. Simply send your name, address and proof of purchase to [insert name and address of supplemental registrant].

OR

SATISFACTION GUARANTEED.

OR

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

OR

Unconditionally guaranteed by [insert name of supplemental registrant]."

[The following claims and product information, may or may not be presented on the product's label and labeling:

- Quick Facts
- Product Facts

- What it does: Kills listed insects, snails and slugs
- Where to Use: For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, trees, and in greenhouses
- Read the Label First!
- Read Entire Container Label Before Using This Product.
- NOTE: This package is sold by weight. Contents may have settled during shipment.
- US Patent number .x,xxx,xxx.
- This container is made from XX% recycled materials.
- The highly compressed granules (pellets) are easy to use and clean to handle.
- Patented technology. Patented insect, snail & (and) slug killer.
- Patented snail, slug & insect (and) killer.
- Unique, patented formula.
- Patented formula
- SLUG, SNAIL AND INSECT BAIT
- SLUGGO plus insect killer
- Slug and bug control
- Slug and bug killer
- INSECT, SLUG AND SNAIL (BAIT)(PELLETS)
- bug and slug control
- bug and slug killer
- Kills (earwigs,) (cutworm,) (sowbugs,) (pillbugs,) (crickets,) (ants, excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants,)(slugs,) (snails,)
- Kills (earwigs) (cutworm)
- Spinosad is derived from a naturally occurring soil dwelling bacterium.
- Spinosad is derived from *Saccharopolyspora spinosa*, a naturally occurring soil organism.
- Spinosad is derived from a soil dwelling bacterium.
- 1 lb treats 2000 ft²
- Easy-to-use (ready-to-use) (RTU) granular (pellet) formulation.
- Easy to apply granules.
- Easy to use small (short) pellets
- Small (short) pellets for greater coverage.
- Shorts
- Pellets
- Kills insects, snails & (and) slugs.
- Treats (will treat) x,xxx sq. ft.
- Remains effective after rain or sprinkling.
- Proven insect, snail & (and) slug killer (kill, control).
- Convenient. Easy-to-use. Requires no mixing, spraying, or special applicators.
- For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, and in greenhouses.
- Read Entire Container Label Before Using This Product.
- For broadcast application standard broadcast applicators may be used, such as (but not limited to): Cyclone.

- For row application standard granular spreaders may be used, such as (but not limited to): Gandy and Clampco.
- Baits and Kills
- The slugs and snails controlled by this product: *Deroceras reticulatum* (Field slug), *Deroceras laeve* (Smooth slug), *Arion subfuscus* (Dusky slug), *Arion circumscriptus* (Gray garden slug), *Arion hortensis* (Black field slug), *Arion rufus* (Large red slug), *Arion ater* (Large black slug), *Limax flavus* (Spotted garden slug), *Limax tenellus* (Slender slug), *Ariolimax columbianus* (Banana slug), *Helix* spp., *Helicella* spp., and *Cepaea* spp.
- Kills a wide variety of insects, slugs and snails
- Effective against a wide variety of listed insects, slugs and snails.
- Effective against a wide variety of, listed slugs, snails and insects.
- No slime trails
- No messy cleanup
- Dead snails, no slime trails
- For control (or suppression) of (earwigs), (cutworm), (sowbugs), (pillbugs), (crickets), (ants, excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants)), (slugs), (snails)
- Contains Spinosad
- Sluggo® is a registered trademark of W. Neudorff GmbH KG, Germany
- May be used as part of an Integrated Pest Management (IPM) program or for organic production.
- Patented insect killer
- Contains Ferramol™ Slug and Snail Bait, a trademark of W. Neudorff GmbH KG
- Made with Ferramol™ Slug and Snail Bait, a trademark of W. Neudorff GmbH KG
- Ferramol™ is a trademark of W. Neudorff GmbH KG, Germany.
- www.neudorff.com
- Manufactured under a license of W. Neudorff GmbH KG, Germany.
- Sold under a license of W. Neudorff GmbH KG, Germany



-For Organic Production

Registrant: W. Neudorff GmbH KG, Postfach 1209, An der Mühle 3,
D-31860 Emmerthal, Germany
Phone: 250-652-5888
www.neudorff.com
[in the alternative, the distributor's telephone number and website address
may be used]