

67702-25

08/30/2012

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON D C 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Mr Walter G Talarek  
W Neudorff GmbH KG  
1008 Riva Ridge Dr  
Great Falls VA 22066 1620

Subject Neudorff Bug Bait EPA Reg No 67702 25  
Date of Registrant Submission July 5 2012  
Decision 467324

AUG 30 2012

Dear Mr Talarek

The labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act as amended is acceptable

Should you wish to retain a reference to the company's website on your label please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition regardless of whether a website is referenced on your product's label claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration the website will be referred to the EPA's Office of Enforcement and Compliance Assurance. As an alternative you may refer consumers to the company's phone number or email address.

Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records. If you have any questions regarding this letter please contact Samantha Hulkower at (703) 603 0683.

Sincerely,


A handwritten signature in black ink, appearing to read "Mark Suarez", is written over the word "Sincerely,".

Mark Suarez  
Product Manager 13  
Insecticide Branch  
Registration Division (7505P)

Enclosure Copy of Label Stamped Accepted

# NEUDORFF BUG BAIT

Active Ingredient	By weight
Spinosad (a mixture of spinosyn A and spinosyn D)	0.07%
Other Ingredients	99.93%
<b>Total</b>	<b>100.00%</b>



**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

FIRST AID
<b>If in eyes</b> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [For emergency information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: <a href="http://www.npic.orst.edu">www.npic.orst.edu</a> ) (alternately insert subregistrants toll free number and hours of operation). During other times, call the poison control center 1-800-222-1222.]

NET WEIGHT 10 lbs 20 lbs 25 lbs 40 lbs 45 lbs 50 lbs 55 lbs (25 kg) 1984 lbs (900 kg) 2000 lbs 2011 lbs (912 kg) and 2116 lbs (960 kg)

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**PERSONAL PROTECTIVE EQUIPMENT (PPE) REQUIREMENTS** Applicators and other handlers must wear long-sleeved shirts and long pants and shoes plus socks. Follow manufacturer's instructions for maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS** This product is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a manner that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency representative responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and the restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

**APPLICATION DIRECTIONS**

**HOW TO APPLY:** Spread the bait on the soil around or near the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply at the higher rate if the infestation is severe. Reapply after heavy rain or watering. Reapply as the bait is consumed or at least every 2 to 4 weeks. Do not apply more than 3 times in any 30 day period.

Do not place in piles. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. See specific directions for different use sites.

Neudorff Bug Bait contains 0.0007 lbs of spinosad per lb. Use the following table to assist in calculating applied a.i. per acre.

**APPLICATION RATE REFERENCE TABLE**

Application Rate of Neudorff Bug Bait	Active Ingredient Equivalent (lb ai/acre)
20 lbs/acre	0.014
25 lbs/acre	0.0175
30 lbs/acre	0.021
35 lbs/acre	0.0245
40 lbs/acre	0.028
44 lbs/acre	0.0308

**PESTS CONTROLLED:** Earwigs, cutworms

**BERRIES:** Spread the bait around the perimeter of the plot to intercept pests migrating towards the berries. If pests are already in the plots, then carefully spread bait between the furrows near the base of the plants. For small plots, treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively. Use the higher rate for heavy infestations.

Application Rate: 20-44 lbs/acre (0.5-1 lb/1000 sq ft)

<b>CROP</b>	<b>Maximum number of applications</b>	<b>Application Interval</b>	<b>Pre Harvest Interval (PHI)</b>	<b>Additional Restrictions</b>
Bushberries (blueberry currant elderberry gooseberry huckleberry juneberry ligonberry salal)	Do not make more than 6 applications per calendar year or more than 3 applications per crop	Do not make applications less than 6 days apart	Do not apply within 3 days of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop
Caneberries (blackberry loganberry red and black raspberry)	Do not make more than 6 applications per calendar year	Do not make applications less than 5 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop
Cranberry	Do not make more than 6 applications per calendar year	Do not make applications less than 7 days apart	Do not apply within 21 days of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop
Strawberry	Do not make more than 5 applications per calendar year	Do not make applications less than 5 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop

**FRUITS INCLUDING CITRUS** Apply in orchards using standard fertilizer granular spreaders. For seedlings, spread the bait around the base of the stem. Apply at 0.15 oz or 1 level teaspoon per square yard in a 6 inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the trees to intercept pests traveling to the trunk. Use the higher rates for heavy infestations. Application Rate: 20-44 lbs/acre (0.5-1 lb/1000 sq ft)

<b>CROP</b>	<b>Maximum number of applications</b>	<b>Application Interval</b>	<b>Pre Harvest Interval (PHI)</b>	<b>Additional Restrictions</b>
Banana and plantain	Do not make more than 6 applications per calendar year or more than 4 applications per crop	Do not make applications less than 7 days apart	Do not apply within 8 weeks of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop
Citrus (grapefruit lemons limes oranges tangerines)	Do not make more than 3 applications per calendar year	Do not make applications less than 6 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop

Fig	Do not make more than 4 applications per calendar year	Do not make applications less than 7 days apart	Do not apply within 7 days of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop
Grape	Do not make more than 5 applications per calendar year	Do not make applications less than 5 days apart	Do not apply within 7 days of harvest	Do not apply more than a total of 0.45 lb ai spinosad per acre per season west of the Rocky Mountains and no more than 0.36 lb ai spinosad per acre per season east of the Rocky Mountains
Pome fruits (apples crabapple loquat mayhaw pears quince)	Do not make more than 4 applications per calendar year	Do not make applications less than 10 days apart	Do not apply within 7 days of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop
Stone fruits – cherries plums prunes		Do not make applications less than 7 days apart	Do not apply within 7 days of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per year
Stone fruits – peaches nectarines apricots		Do not make applications less than 7 days apart	Do not apply within 14 days of harvest for apricots or within 1 days of harvest for nectarines and peaches	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per year
Tropical Tree fruits (avocado guava mango papaya passionfruit)	Do not make more than 2 applications per calendar year	Do not make applications less than 7 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop

VEGETABLES Scatter the bait around the perimeter of the vegetable plantings to provide a protective barrier for pests entering the vegetable plantings. If pests are inside the rows then scatter the bait on the soil around the plants and between the rows. Application Rate 20-44 lbs/acre (0.5-1 lb/1000 sq ft)

<b>CROP</b>	<b>Maximum number of applications</b>	<b>Application Interval</b>	<b>Pre Harvest Interval (PHI)</b>	<b>Additional Restrictions</b>
Artichoke	Do not make more than 4 applications per crop	Do not make applications less than 7 days apart	Do not apply within 2 days of harvest	Do not apply more than a total of 0.33 lb ai spinosad (including foliar uses) per crop
Asparagus (For post harvest protection of ferns only)	Do not make more than 3 applications per crop	Do not make applications less than 4 days apart	Do not apply within 60 days of spear harvest	Do not apply more than a total of 0.28 lb ai spinosad (including foliar uses) per crop. Do not feed treated ferns to meat or dairy animals.
Legume vegetables (succulent and dried beans and peas)	Do not make more than 6 applications per crop	Do not make applications less than 5 days apart	Succulent beans and peas: Do not apply within 3 days of harvest. Dried beans and peas: Do not apply within 28 days of harvest.	Succulent beans and peas: Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per season. Dried beans and peas: Do not apply more than a total of 0.188 lb ai spinosad per acre per season.
Bulb vegetables (dry bulb onion, garlic, great headed (elephant) garlic, green onion, leek, shallot, welsch onion)	Do not make more than 5 applications per calendar year	Do not make applications less than 4 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar and seed uses) per acre per crop.
Cole crops (broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, izuna, mustard greens, mustard spinach, rape greens)	Do not make more than 6 applications per calendar year	Do not make applications less than 4 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop. Do not apply to seedling cole crops grown for transplant.

Corn – field corn	Do not make more than 3 applications per calendar year		Do not apply within 28 days of harvest for grain or fodder 7 days for forage	Do not apply more than a total of 0.188 lb ai spinosad (including foliar uses) per acre per year
Corn – sweet corn	Do not make more than 6 applications per calendar year		Do not apply within one day of harvest for grains 7 days for forage 28 days for fodder	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per year
Cucurbits (cucumber edible gourds melons muskmelons cantaloupe honeydew pumpkin summer squash watermelon winter squash)	Do not make more than 6 applications per crop	Do not make applications less than 5 days apart	Do not apply within one day of harvest for cucumber 3 days for other cucurbits	Do not apply more than a total of 0.45 lb ai spinosad (including foliar and seed uses) per acre per season
Fruiting vegetables (eggplant groundcherry pepino pepper (except black) tomatillo tomato) and okra	Do not make more than 6 applications per calendar year	Do not make applications less than 4 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop Do not apply to seedling fruiting vegetables and okra grown for transplant within a greenhouse shade house or field plot
Leafy vegetables (except Brassica) and leaves of root and tuber and legume vegetables (arugula beets celery peas lettuce rhubarb spinach swiss chard) turnip greens and watercress	Do not make more than 6 applications per calendar year	Do not make applications less than 4 days apart	Do not apply within one day of harvest for leafy vegetables and watercress 3 days for leaves of root tuber and legume vegetables	Do not apply more than a total of 0.45 lb ai spinosad (including foliar and seed uses) per acre per crop Do not apply to seedling leafy crops grown for transplant within a greenhouse or shade house
Potatoes ginger sweet potatoes yams	Do not make more than 4 applications per crop	Do not make applications less than 7 days apart	Do not apply within 7 days of harvest	Do not apply more than a total of 0.33 lb ai spinosad (including foliar uses) per crop

Root vegetables – Carrot parsnip	Do not make more than 4 applications per calendar year	Do not make applications less than 5 days apart	Do not apply within 3 days of harvest	Do not apply more than a total of 0.3 lb ai spinosad (including foliar uses) per acre per crop
Root vegetables – Radish rutabaga turnip	Do not make more than 3 applications per calendar year	Do not make applications less than 5 days apart	Do not apply within 3 days of harvest	Do not apply more than a total of 0.28 lb ai spinosad (including foliar uses) per acre per crop
Root and Tuber Vegetables (beets sugarbeets)	Do not make more than 4 applications per crop	Do not make applications less than 7 days apart	Do not apply within 3 days of harvest	Do not apply more than a total of 0.33 lb ai spinosad (including foliar uses) per acre per crop
Soybean	Do not make more than 4 applications per calendar year	Do not make applications less than 4 days apart	Do not apply within 28 days of harvest	Do not apply more than a total of 0.186 lb ai spinosad (including foliar uses) per acre per year. Do not feed treated forage or hay to meat or dairy animals.

**TREE NUTS PEANUTS AND PISTACHIOS** For seedlings spread the bait around the base of the stem. For older trees spread around the base of the trees to intercept pests traveling to the trunk. Use the higher rates for heavy infestations. Application Rate 20-44 lbs/acre (0.5-1 lb/1000 sq ft)

<b>CROP</b>	<b>Maximum number of applications</b>	<b>Application Interval</b>	<b>Pre Harvest Interval (PHI)</b>	<b>Additional Restrictions</b>
Peanut (not for use in California)	Do not make more than 3 applications per calendar year	Do not make applications less than 7 days apart	Do not apply within 3 days of nut harvest or 14 days of forage	Do not apply more than a total of 0.28 lb ai spinosad (including foliar uses) per acre per crop. Do not allow grazing of crop residue or harvest of crop residue for hay until 14 days after the last application.
Tree nuts (almond cashew chestnut filbert (hazelnut) macadamia nut pecan walnut) Pistachios		Do not make applications less than 7 days apart	Do not apply within 1 day of harvest	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop



HERBS At the seedling and later stages apply the bait between the rows and around the perimeter of the field Use the higher dosage for heavy infestations Application Rate 20 44 lbs/acre (0 5 1 lb/1000 sq ft)

CROP	Maximum number of applications	Application Interval	Pre Harvest Interval (PHI)	Additional Restrictions
Herbs (angelica balm basil borage burnet chamomile catnip chervil (dried) chive chive (Chinese) cilantro (leaf) clary coriander (leaf) costmary cilantro curry (leaf) dillweed horehound hyssop lavender lemongrass lovage (leaf) marigold marjoram nasturtium parsley (dried) pennyroyal rosemary rue sage savory (summer and winter) sweet bay tansy tarragon thyme wintergreen woodruff and wormwood)	Do not make more than 5 applications per calendar year or more than 3 applications per crop	Do not make applications less than 5 days apart	Do not apply within one day of harvest	Do not apply more than a total of 0 47 lb ai spinosad (including foliar uses) per acre per crop
Peppermint and spearmint	Do not make more than 4 applications per calendar year or more than 3 applications per crop	Do not make applications less than 4 days apart	Do not apply within 7 days of harvest	Do not apply more than a total of 0 45 lb ai spinosad (including foliar uses) per acre per crop

**GRASS GROWN FOR SEED PRODUCTION AND CEREAL GRAINS** When pests are detected scatter the bait by hand or with a granular spreader Use the higher rate for heavy infestations Application Rate 20 44 lbs/acre (0 5 1 lb/1000 sq ft)

<b>CROP</b>	<b>Maximum number of applications</b>	<b>Application Interval</b>	<b>Pre Harvest Interval (PHI)</b>	<b>Additional Restrictions</b>
Small cereal grains (barley buckwheat grain amaranth milo oats pearl millet proso millet rye sorghum triticale wheat)	Do not make more than 3 applications per calendar year	Do not make applications less than 4 days apart	Do not apply within 3 days of harvest for forage fodder or hay Do not apply within 21 days of harvest for grain or straw	Do not apply more than a total of 0 28 lb ai spinosad (including foliar uses) per acre per year
Grass crops grass grown for seed pastures and rangeland	Do not make more than 6 applications per season	No preharvest interval for forage	Do not use within 3 days of harvest for hay or fodder	Do not apply more than a total of 0 186 lb ai spinosad (including foliar uses) per acre per season

#### **TURF AND ORNAMENTALS**

**PESTS CONTROLLED** Ants (excluding fire ants harvester ants Pharaoh s ants and carpenter ants earwigs cutworms

When pests are detected broadcast the bait by hand or with a granular spreader Alternatively in or around plant pots on greenhouse benches or in or around container grown plants apply at 0 15 oz or 1 level teaspoon per square yard in a 6 inch circular band around the base of the plants to be protected Use the higher rate for heavy infestations Application Rate 20 44 lbs/acre (0 5 1 lb/1000 sq ft)

<b>CROP</b>	<b>Maximum number of applications</b>	<b>Application Interval</b>	<b>Additional Restrictions</b>
Turf (lawns athletic field and recreational areas commercial turf sod golf courses and grass buffers around gardens crop areas and ornamentals)		Do not make applications less than 7 days apart	Do not apply more than a total of 0 45 lb ai spinosad (including foliar uses) per acre per year
Outdoor Ornamentals (herbaceous and woody plants)		Do not make applications less than 7 days apart	Do not apply more than a total of 0 45 lb ai spinosad (including foliar uses) per acre per year
Greenhouse Ornamentals	Do not make more than 10 applications per calendar year	Do not make applications less than 7 days apart	Do not apply more than a total of 0 45 lb ai spinosad (including foliar uses) per acre per year
Outdoor container grown nursery plants		Do not make applications less than 7 days apart	Do not apply more than a total of 0 45 lb ai spinosad (including

			foliar uses) per acre per year
Tree Farms and Plantations (Conifers including Christmas trees and deciduous trees)	Do not make more than 6 applications per calendar year	Do not make applications less than 7 days apart	Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per year
Non crop areas (pastures rangeland forests parks playgrounds cemeteries picnic areas roadsides campgrounds industrial sites fallow land barrier strips and buffer zones)			

#### STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

**PESTICIDE STORAGE** Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals

**PESTICIDE DISPOSAL** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

**CONTAINER HANDLING** Nonrefillable container Do not reuse or refill this container Completely empty drum bag box or tote into application equipment by shaking and trapping sides and bottom to loosen clinging particles Then offer for recycling if available or dispose of empty drum bag box or tote in a sanitary landfill or by incineration or by other procedures allowed by State and local authorities

#### BATCH CODE

[The registrant may use one or both of these two optional statements either

#### WARRANTY

To the extent consistent with applicable law seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated on this label only when used in accordance with directions under normal use conditions To the extent consistent with applicable law this warranty does not extend to use of this product contrary to label directions or under abnormal use conditions or under conditions not reasonably foreseeable to seller To the extent consistent with applicable law seller makes no other warranties either expressed or implied

or



#### OUR COMMITMENT TO YOUR SATISFACTION

If you are not happy with the way this product works in your garden please let us know We will be happy to replace the product or refund your money Simply send your name address and proof of purchase to [insert name and address of supplemental registrant]

]

[The following claims and product information may or may not be presented on the product's label and labeling

- Quick Facts
- Product Facts
  - What it does Kills insects

- Where to Use For use around vegetables fruit trees citrus berries ornamentals shrubs flowers trees gardens and in greenhouses
- Read the Label First!
- **NOTE** This package is sold by weight Contents may have settled during shipment
-  For Organic Gardening
-  For Organic Production
- Neudorff Bug Bait may be used as part of an Integrated Pest Management (IPM) program or for organic production
- Patent Pending
- This container is made from XX% recycled materials
- The highly compressed granules (pellets) are easy to use clean to handle and economical
- Patented technology
- Patented insect killer
- Patented formula
- Easy to use (ready to use) (RTU) granular (pellet) formulation
- Easy to apply granules (pellets)
- Pellets
- Kills insects
- INSECT BAIT
- INSECT (BAIT) (PELLETS)
- Kills (ants excluding fire ants harvester ants Pharaoh s ants and carpenter ants ) (earwigs ) (cutworm)
- Spinosad is derived from the fermentation of a naturally occurring soil dwelling organism
- Treats (will treat) x xxx sq ft
- 1 lb treats 2000 ft<sup>2</sup>
- Effective for up to 4 weeks
- Lasts up to 4 weeks
- Remains effective for up to 4 weeks
- The highly compressed easy to apply granules can be effective up to four (4) weeks
- Neudorff Bug Bait contains spinosad derived from naturally occurring soil dwelling bacterium in a patented granular formulation that requires no mixing spraying or special applicators
- Proven insect (killer) (kill) (control)
- Convenient Easy to use Requires no mixing spraying or special applicators
- SATISFACTION GUARANTEED
- Money Back Guarantee If you are not satisfied with this product we will gladly refund your original purchase price
- Can be used in vegetable gardens
- Ideal for vegetable gardens
- For use around listed vegetables fruit trees citrus berries ornamentals shrubs flowers trees gardens and in greenhouses
- For use on lawns commercial turf recreational areas and golf courses
- Use around flowers gardens and fruit trees
- Use around turf and ornamentals
- Can be used around turf ornamentals and non crop areas
- Read Entire Container Label Before Using This Product
- Unconditionally guaranteed by [insert name of supplemental registrant]
- Baits and Kills
- Effective against a wide variety of insects
- Lures insects from hiding places and plants
- Attracts and kills insects
- Attracts and kills

- For control (or suppression) of (ants excluding fire ants harvester ants Pharaoh s ants and carpenter ants) (earwigs) (cutworm)
- Contains Spinosad
- [www.neudorff.de](http://www.neudorff.de)
- Manufactured under a license of W Neudorff GmbH KG Germany



- **NEUDORFF**
- Evening is the best time to apply the bait as pests travel and feed mostly by night or early morning
- Ant Bait (excluding fire ants harvester ants Pharaoh s ants and carpenter ants)
- Ant Killer (excluding fire ants harvester ants Pharaoh s ants and carpenter ants)
- Controls foraging ants (excluding fire ants harvester ants Pharaoh s ants and carpenter ants)
- Controls common household ants (excluding fire ants harvester ants Pharaoh s ants and carpenter ants)
- Can be used on listed household ants around the home
- Cutworm Bait
- Cutworm Killer
- Earwig Bait
- Earwig Killer
- Earwig and Cutworm Killer
- Earwig and Cutworm Bait
- Ant Earwig and Cutworm Killer
- Ant Earwig and Cutworm Bait
- Neudorff Bug Bait is a soil applied granular insecticidal bait containing the active ingredient spinosad that attracts and kills insects including ants (excluding fire ants harvester ants Pharaoh s ants and carpenter ants) earwigs and cutworms in commercial agricultural crops ornamentals herbs seed crops and lawns
- Listed by the Organic Materials Review Institute (OMRI) for use in organic production
- OMRI Listed
- **OMRI**  
LISTED
- ]

EPA Registration #67702 25

EPA Establishment #67702 DEU 2

Registrant W Neudorff GmbH KG Postfach 1209 An der Muhle 3  
D 31860 Emmerthal Germany  
011 49 5155 624 100  
[www.neudorff\[pro\].com](http://www.neudorff[pro].com)  
[in the alternative the distributor s telephone number and website address may be used]