



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 19, 2023

Mr. Walter G. Talarek
Authorized Agent
W. Neudorff GmbH KG
c/o Walter G. Talarek PC
5153 Allison Marshall Dr.
Warrenton, Va. 20187

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Adding to/revising sublabel(s): container size; optional statements; notes to reviewer; and miscellaneous editorial revisions
Product Name: NEU1572H
EPA Registration Number: 67702-63
EPA Receipt Date: 02/09/2023
Action Case Number: 00431253

Dear Mr. Talarek:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Page 2 of 2


EPA Reg. No. 67702-63

Action Case No. 00431253

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Nina Naimy via email at naimy.nina@epa.gov.


Sincerely,

A handwritten signature in blue ink, appearing to read "James Parker", is written over a thin blue horizontal line.

James Parker, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

Enclosure

Text in braces { } is for information only and will not appear on the label
Text in brackets [] is optional or a 'place holder' for graphics and may or may not appear on the label
Text in parentheses () will appear on the label



MASTER LABEL

NEU1572H
ABN: Fiesta Granular Weed & Feed
ABN: Fiesta Granular Turf Weed Killer
ABN: Fiesta Weed and Feed Granule
ABN: Fiesta Granular Selective Herbicide

Active Ingredient:	
Sodium Ferric EDTA.....	14.00%
Other Ingredients:.....	86.00%
Total:.....	100.00%

SUBLABEL A: Household Use
SUBLABEL B: Commercial Use

EPA REG. NO. 67702-63

EPA EST.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Registrant: W. Neudorff GmbH KG
An der Muhle 3,
31860 Emmerthal, Germany
www.neudorff.com

Household Use: NET WEIGHT: 1 oz, 13 oz, 26 oz, 0.5 lb, 1 lb, 1.25 lbs (20 oz), 1.5 lbs, 30 oz, 2 lbs, 2.25 lbs, 2.5 lbs (40 oz), 3 lbs, 3.2 lbs, 4 lbs, 5 lbs, 5.25 lbs, 5.5 lbs, 6 lbs, 7 lbs, 7.5 lbs, 8 lbs, 10 lbs, 12lbs, 15lbs, 20 lbs, 25 lbs, 40 lbs, 50 lbs, 2116 lbs

Commercial Use: NET WEIGHT: 5 lbs, 10 lbs, 20 lbs, 25 lbs, 40 lbs, 45 lbs, 50 lbs, 55 lbs (25 kg), 1984 lbs (900 kg), 2011 lbs (912 kg), 2116 lbs (960 kg)

SUBLABEL A – HOUSEHOLD USE

NEU1572H

Active Ingredient:

Sodium Ferric EDTA.....	14.00%
Other Ingredients:.....	86.00%
Total:.....	100.00%

EPA REG. NO. 67702-

EPA EST.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID [⊕]

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. {Optionally, supplemental distributors may use the following wording and own toll-free hotline number;} [For medical emergency information, [you may call toll free 1-XXX-XXX-XXXX] {insert operating hours and days of service} [During other times] call the poison control center at 1-800-222-1222.]

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not expected for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

{Use one of the following two formats; Latin names are optional throughout directions for use}

{Format 1:}

Use Sites:

Use to selectively control/suppress broadleaf weeds, moss, algae*, lichens*, liverworts* and diseases* growing in lawns.

* Not for use in California

Pests:

Controls or suppresses weeds such as but not limited to:

- Black medic (*Medicago lupulina*)[§]
- Broadleaf plantain (*Plantago major*)*

- Bull thistle (*Cirsium vulgare*)
- Canada thistle (*Cirsium arvense*)*
- Common chickweed (*Stellaria media*)§
- Crabgrass (*Digitaria spp.*)- pre-emergent control only
- Creeping buttercup (*Ranunculus repens*)
- Creeping Charlie/Ground ivy (*Glechoma hederacea*)*
- Dandelion (*Taraxicum officinale*)
- Dovefoot geranium (*Geranium molle*)*
- English daisy (*Bellis perennis*)§
- False dandelion (*Hypochoeris radicata*)*
- Heal-all (*Prunella vulgaris*)*
- Lawn burweed (*Soliva ptero-sperma*)*
- Narrowleaf plantain (*Plantago lanceolata*)*
- Oxalis (*Oxalis spp.*)*
- Prostrate knotweed (*Polygonum aviculare*)*
- Shepherd's-purse (*Capsella bursa-pastoris*)*
- Silverweed cinquefoil (*Potentilla anserina*)*
- *Speedwell spp.*(including Persian Speedwell (*Veronica persica*) and Slender speedwell (*Veronica filiformis*)*)
- White clover (*Trifolium repens*)
- Wild chamomile (*Matricaria chamomilla*)*
- Wild geranium (*Geranium robertianum*)*
- moss, algae*, lichens*, liverworts*

Controls or suppresses diseases such as but not limited to:

- Dollar Spot*
- Rust*
- Snow Mold*

* Not for use in California

§ only approved for suppression in California

[Application Directions:

Apply at a rate of 3-8 lbs /1000 ft² (0.5 -1.15 oz /yard²), or 5-8 lbs/1000 ft² (0.72 - 1.15 oz /yard²) in California. {Optional:} [The standard rate for this product is 5lbs /1000 ft² (0.72 oz /yard²); use the higher rate of 8 lbs /1000 ft² (1.15 oz /yard²) on tough perennial weeds, such as common chickweed[§], buttercup and broadleaf plantain*. Use at 3 lbs /1000 ft² † (0.5 oz /yard²) on moss, and weed seedlings of false dandelion*, English daisy*, dandelion* or white clover*.] Repeat treatment in 3 to 4 weeks for best results.

For Spot Treatment, apply 5 lbs /1000 ft² (0.72 oz /yard²). Repeat treatment in 3 to 4 weeks for best results.

For Pre-emergent control of broadleaf weeds and crabgrass, begin application when soil temperatures reach 50 to 55 °F. Apply 5-8 lbs/1000 ft² (0.72 -1.15 oz /yard²), depending on weed pressure. Repeat treatment in 3 to 6 weeks for best results.

{Optional table}

Rate	Weed Species	Diseases
3 lbs/ 1000 ft ² (0.43 oz/yd ²) [†]	Weed seedlings of English daisy* (<i>Bellis perennis</i>), dandelion* (<i>Taraxacum officinale</i>) false dandelion* (<i>Hypochaeris radicata</i>), young moss	Dollar spot *, snow mold*, and rust*. Use the higher rate when disease pressure is high.
4 lbs/ 1000 ft ² (0.58 oz/yd ²) [†]	Moss, speedwell species, including Persian speedwell, (<i>Veronica persica</i>) and slender speedwell* (<i>Veronica filiformis</i>), wild geranium* (<i>Geranium robertianum</i>)	
5 lbs/ 1000 ft ² (0.72 oz/yd ²)	Black medic (<i>Medicago lupulina</i>) [§] , bull thistle (<i>Cirsium vulgare</i>), Canada thistle* (<i>Cirsium arvense</i>), creeping charlie/ground ivy* (<i>Glechoma hederacea</i>), dandelion (<i>Taraxacum officinale</i>), dove's foot geranium* (<i>Geranium molle</i>), English daisy (<i>Bellis perennis</i>), false dandelion* (<i>Hypochaeris radicata</i>), heal-all* (<i>Prunella vulgaris</i>), lawn burweed* (<i>Soliva ptero-sperma</i>), narrowleaf plantain* (<i>Plantago lanceolata</i>), oxalis* (<i>Oxalis</i> spp.), prostrate knotweed* (<i>Polygonum aviculare</i>), Shepherd's-purse* (<i>Capsella bursa-pastoris</i>), silverweed cinquefoil* (<i>Potentilla anserina</i>), algae*, lichens*, liverworts*. Preemergent control of crabgrass (<i>Digitaria</i> spp.).	
8 lbs/ 1000 ft ² (1.15 oz/yd ²)	Common chickweed [§] (<i>Stellaria media</i>), creeping buttercup (<i>Ranunculus repens</i>), broadleaf plantain* (<i>Plantago major</i>), white clover* (<i>Trifolium repens</i>).	

]

[†] Lower application rate not for use in California. Minimum application rate in California is 5 lbs/1000 ft²

* Not for use in California

§ only approved for suppression in California

How to Apply:

Contents may have settled during shipment; invert package several times to ensure product is mixed. For best results, apply a few days after mowing when temperature is between 50°F and 85°F. If temperature is below 80°F ensure lawn is wet from dew, or water lawn lightly before application. If temperature is above 80°F do not pre-wet grass and apply product in the late afternoon or evening. Do not exceed the 5 lbs/1000 ft² rate when temperatures are above 85°F. Do not apply when daytime temperature exceeds 90°F.

Uniform coverage is important to prevent lawn damage. Apply with a broadcast/rotary, drop, or handheld spreader set to the correct application rate. See table below for spreader settings. Wait at least 24 hours after application to water lawn. Lawn areas can be re-sown one day after treatment.

Apply product only on lawn. Application on hard surfaces can cause staining. If product gets on surface, sweep off if the surface is dry, rinse off immediately if the surface is wet. If product gets on clothing or footwear, wash immediately.

[Spreader settings:

Weed Species	Rate	{Optional column} Scotts® Broadcast and Rotary Spreader Setting (All Models)	{Optional column} Scotts® Drop Spreader Setting (All Models)	{Optional column} Scotts® Hand-Held Spreader Setting (All models)
Weed seedlings of English daisy* (<i>Bellis perennis</i>), dandelion* (<i>Taraxacum officinale</i>) false dandelion* (<i>Hypochaeris radicata</i>), young moss	3 lbs per 1000 square feet	2.5	5	3
Black medic (<i>Medicago lupulina</i>) [§] , bull thistle (<i>Cirsium vulgare</i>), Canada thistle* (<i>Cirsium arvense</i>), creeping charlie/ground ivy* (<i>Glechoma hederacea</i>), dandelion (<i>Taraxacum officinale</i>), dove's foot geranium* (<i>Geranium molle</i>), English daisy (<i>Bellis perennis</i>), false dandelion* (<i>Hypochaeris radicata</i>), heal-all* (<i>Prunella vulgaris</i>), lawn burweed* (<i>Soliva pterosperma</i>), narrowleaf plantain* (<i>Plantago lanceolata</i>), oxalis* (<i>Oxalis</i> spp.), prostrate knotweed* (<i>Polygonum aviculare</i>), Shepherd's-purse* (<i>Capsella bursa-pastoris</i>), silverweed cinquefoil* (<i>Potentilla anserina</i>), speedwell species, including Persian speedwell, (<i>Veronica persica</i>) and slender speedwell* (<i>Veronica filiformis</i>), wild geranium* (<i>Geranium robertianum</i>), moss, algae*, lichens*, liverworts*. Preemergent control of crabgrass (<i>Digitaria</i> spp.).	5 lbs/1000 square feet	3	6	3.5
Common chickweed [§] (<i>Stellaria media</i>), creeping buttercup (<i>Ranunculus repens</i>), broadleaf plantain* (<i>Plantago major</i>), white clover* (<i>Trifolium repens</i>).	8 lbs/1000 square feet	4.5	7.5	5

† Lower application rate not for use in California. Minimum application rate in California is 5 lbs/1000 ft²

* Not for use in California

§ only approved for suppression in California

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.] [Spreaders other than those listed in this table may be used, as long as they are calibrated prior to use to deliver the product within the use rates on this label.]

What will I see?

Works quickly; dying weeds will be seen within 24 hours of application. Treated weeds, moss, or algae* will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however, the grass will recover within a few days to weeks. May stain thatch (dead grass under lawn); this does not affect the health of the grass. If lawn is damaged, irrigation 24 hours after application, followed by mowing to remove the lawn tip burn can help lawn recover faster.

* Not for use in California

Use Precautions:

Do not apply to drought stressed grass; ensure lawn is well watered prior to application. Do not apply when the daytime temperature will exceed 90°F. Do not apply to bentgrass. If unsure of grass sensitivity, make a test application to a small area of your lawn. Avoid applying to desirable plants. Do not apply if rainfall is expected within 24 hours of application. Do not irrigate within 24 hours of application.

]

{OR}

{Format 2:}

[

DIRECTIONS FOR USE	
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.	
BEFORE YOU USE	Contents may have settled during shipment; invert bag several times to ensure product is mixed. For best results, apply a few days after mowing when temperature is between 50°F and 85°F. If temperature is below 80°F ensure lawn is wet from dew, or water lawn lightly before application. If temperature is above 80°F do not pre-wet grass and apply product in the late afternoon or evening. Do not exceed the 5 lbs/1000 ft ² rate when temperatures are above 85°F. Do not apply when daytime temperature exceeds 90°F.
WHERE TO USE	Use to selectively control/suppress [broadleaf weeds], [moss], [algae]*, [lichens]*, [liverworts]*, and [diseases]* growing in lawns
HOW TO USE	Apply at a rate of 3-8 lbs /1000 ft ² (0.5 -1.15 oz /yard ²), or 5-8 lbs/1000 ft ² (0.72 -1.15 oz /yard ²) in California. {Optional:} [The

<p>[Spot treatment]</p> <p>[Pre-emergent]</p>	<p>standard rate for this product is 5 lbs /1000 ft² (0.72 oz /yard²); use the higher rate of 8 lbs /1000 ft² (1.15 oz /yard²) on tough perennial weeds, such as common chickweed[§], buttercup and broadleaf plantain*. Use at 3 lbs /1000 ft² † (0.5 oz /yard²) on moss, and weed seedlings of false dandelion*, English daisy*, dandelion* or white clover*.] Repeat treatment in 3 to 4 weeks for best results.</p> <p>For Spot Treatment, apply 5 lbs /1000 ft² (0.72 oz /yard²). Repeat treatment in 3 to 4 weeks for best results.</p> <p>For Pre-emergent control of broadleaf weeds and crabgrass, begin application when soil temperatures reach 50 to 55 °F. Apply 5-8 lbs/1000 ft² (0.72 -1.15 oz /yard²), depending on weed pressure. Repeat treatment in 3 to 6 weeks for best results.</p> <p>Uniform coverage is important to prevent lawn damage. Apply with a broadcast/rotary, drop, or handheld spreader set to the correct application rate. See table below for spreader settings. Wait at least 24 hours after application to water lawn. Lawn areas can be re-sown one day after treatment.</p> <p>Apply product only on lawn. Application on hard surfaces can cause staining. If product gets on surface, sweep off if the surface is dry, rinse off immediately if the surface is wet. If product gets on clothing or footwear, wash immediately.</p>
<p>WHAT WILL I SEE?</p>	<p>Works quickly; dying weeds will be seen within 24 hours of application. Treated weeds, moss, or algae* will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however, the grass will recover within a few days to weeks. May stain thatch (dead grass under lawn); this does not affect the health of the grass. If lawn is damaged, irrigation 24 hours after application, followed by mowing to remove the lawn tip burn can help lawn recover faster.</p>
<p>PESTS (WEEDS, DISEASES*, MOSS, LICHENS* & ALGAE*)</p>	<ul style="list-style-type: none"> • Black medic (<i>Medicago lupulina</i>)[§] • Broadleaf plantain (<i>Plantago major</i>)* • Bull thistle (<i>Cirsium vulgare</i>) • Canada thistle (<i>Cirsium arvense</i>)* • Common chickweed (<i>Stellaria media</i>)[§] • Crabgrass (<i>Digitaria spp.</i>)- pre-emergent control only • Creeping buttercup (<i>Ranunculus repens</i>) • Creeping Charlie/Ground ivy (<i>Glechoma hederacea</i>)* • Dandelion (<i>Taraxicum officinale</i>) • Dovefoot geranium (<i>Geranium molle</i>)* • English daisy (<i>Bellis perennis</i>)[§] • False dandelion (<i>Hypochoeris radicata</i>)* • Heal-all (<i>Prunella vulgaris</i>)* • Lawn burweed (<i>Soliva ptero-sperma</i>)*

	<ul style="list-style-type: none"> • Narrow-leaf plantain (<i>Plantago lanceolata</i>)* • Oxalis (<i>Oxalis</i> spp.)* • Prostrate knotweed (<i>Polygonum aviculare</i>)* • Shepherd's-purse (<i>Capsella bursa-pastoris</i>)* • Silverweed cinquefoil (<i>Potentilla anserina</i>)* • <i>Speedwell</i> spp.(including Persian Speedwell (<i>Veronica persica</i>) and Slender speedwell (<i>Veronica filiformis</i>)*) • White clover (<i>Trifolium repens</i>) • Wild chamomile (<i>Matricaria chamomilla</i>)* • Wild geranium (<i>Geranium robertianum</i>)* • moss, algae*, lichens*, liverworts* <p>Controls or suppresses diseases such as but not limited to:</p> <ul style="list-style-type: none"> • Dollar Spot* • Rust* • Snow Mold*
<p>USE PRECAUTIONS</p>	<ul style="list-style-type: none"> • Do not apply to drought stressed grass. Ensure that lawn is well watered prior to application. • Do not apply when the daytime temperature will exceed 90°F. • Do not apply to bentgrass. • If unsure of grass sensitivity, make a test application to a small area of your lawn. • Avoid applying to desirable plants. • Do not apply if rainfall is expected within 24 hours of application. Do not irrigate within 24 hours of application.

† Lower application rate not for use in California. Minimum application rate in California is 5 lbs/1000 ft²

* Not for use in California

§ only approved for suppression in California

Spreader settings:

{Spreader setting table to be inserted here including graphics, if applicable}

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.

]

<p>STORAGE AND DISPOSAL </p> <p>PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. Store unused material in a cool, dry place tightly closed, in original container only.</p> <p>PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available.</p>

If partly filled: Call your local solid waste agency [or call [insert supplemental distributor's toll free number]] for disposal instructions. Never place unused product down any indoor or outdoor drain.

BATCH CODE:

{Optional fertilizer guarantee information. The following table may be used in whole or in part. Fertilizer information denoted by "x" will be added to FPL. NPK will not exceed the endpoints tested for acute toxicology studies.}

[

Guaranteed Fertilizer Analysis XX-XX-XX {insert N-P-K ratio}	
Total nitrogen (N) X% Ammoniacal nitrogen X% Urea nitrogen X% Other water soluble nitrogen X% Water insoluble nitrogen	X%
Available phosphate (P ₂ O ₅)	X%
Iron (Fe)	X%
Sulfur (S)	X%
Magnesium (Mg) (Total) X% water soluble magnesium X% chelated magnesium	X%
Sulfur X% combined sulfur X% free sulfur	X%
Soluble potash (K ₂ O)	X%
Calcium (Ca)	X%
Derived From: {Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer.}	

]

{Note to reviewer: Spanish label text may also be included in addition to English label text.}

{The following claims and product information, may or may not be presented on the product's label and labeling:}

[

{For products being sold into states requiring a metals internet statement on fertilizer products, the following statement will be used:}

-Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

{For products being sold into Florida, the following statements will be used:}

-FL License No.: FXXXXXX

-Florida Applications:

We recommend that you follow the Florida Green Industries BMP's at:

<http://www.flrules.org/Gateway/reference.asp?No=Ref-04706>

We recommend that you follow the Florida Golf Course BMP's at:

<http://www.flrules.org/Gateway/reference.asp?No=Ref-04702>


{For states with nitrogen application restrictions, including but not limited to, MD, NH, NJ, PA and VA, the following statement will be used:}

-Nitrogen Notice: Do not apply more than X.XX lbs. of this product per 1,000 sq.ft.

-Phosphate Notice: This lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.

-www.neudorff.com

-www.lawnbox.com



-[Moss], [weed], [algae*], [lichen*], [liverwort*] and [disease*] killer [for lawns]

-Lawn [Moss], [weed], [algae*], [lichen*], [liverwort*] and [disease*] [Killer]
[Control]

-[Selective] [Broadleaf] [Weed], [algae*], [lichen*], [liverwort*] and [disease*]
[Killer][Control]

-[Selective Control of][Controls] [Broadleaf] Weeds in Lawns

-Kills [moss], [weed], [algae*], [lichen*], [liverwort*] and [diseases*], not
[lawns][grass]

-Kills [moss], [weed], [algae*], [lichen*], [liverwort*] and [diseases*] without
harming lawn

-Kills [weeds], [algae*], [moss], [liverworts*] and [lichens*]









-Easy-to-use [moss], [weed], [algae*], [lichen*], [liverwort*] and [disease*] killer
for the lawn

-Cure your [weed], [moss] and [disease*] problems fast

- Ideal [weed], [moss] & [disease*] control for lawn
- Iron-based [disease], [moss], [disease*] & [weed] control for lawn(s)
- Controls dollar spot*, snow mold*, and rust* [on lawns]
- Lawn [Fungicide*], [Herbicide] and [Fertilizer]
- Kills [dandelions], [clover] and [most] other [major][broadleaf] lawn weeds* *as listed
- Prevents and controls [Moss], [weed], [algae*], [lichen*], [liverwort*], and [lawn diseases*]
- [Controls] [kills] [dandelions] [buttercup] [English daisy] [daisy] [clover]
- Controls disease* in the lawn, kills the weeds.
- [Disease*], [moss] and [Weed] Killer in one!
- Tough on [moss] [weeds] and [disease*], easy on lawns
- Prevent disease* and kill the weeds!
- Kills most common lawn diseases* fast!
- Kills [weeds][dandelions], down to the root
- NEU1572H is derived from iron that is used to selectively control/ suppress [broadleaf weeds], [moss], and [diseases*] growing in lawns
- Herbicide and Fungicide* in 1 for the Lawn!
- NEU1572H is formulated to feed your lawn and control [broadleaf weeds], [moss], [algae*], [diseases*] commonly found in lawns.
- Pre- and Post-emergent activity of broadleaf weeds with one application
- Prevents crabgrass and broadleaf weeds from germinating
- Pre and post emergent [N-P-K] herbicide
- Pre and post emergent weed control for lawns
- Controls weeds before they sprout
- Provides pre-emergent control of weeds in lawns
- Pre-emergent control of [broadleaf weeds] and [crabgrass] [in lawns]
- Controls weeds before you see them (!)
- Apply pre-emergence for extended weed control
- Controls Weeds and Feeds your Lawn.
- Builds thick, green lawn [from the roots up]
- Builds thick, green lawn while killing listed weeds
- Strengthens and protects your lawn
- Greens up lawn in 7 days
- Thickens lawn to crowd out weeds
- [Fast] [quick] green-up
- Greens your lawn fast
- Two [2] in One [1] fertilizer and [weed][moss] killer
- Kills [weeds] and [moss], feeds your lawn
- Provides fast green-up of lawn
- Contains Iron, a plant nutrient
- [Guaranteed Analysis]: xx-xx-xx {insert N-P-K ratio} [Fe 1.82%]
- [X-X-X {insert N-P-K ratio}] [1.82% Fe] Nitrogen, Phosphorus, Potash / Potassium, [Iron]
- [N] Nitrogen [and [Fe] Iron] for greening [P] Phosphorus helps develop a stronger, hardier root system [K] Potassium / Potash improves plant hardiness and drought resistance
- Healthier, stronger and greener lawn

-WEED AND FEED

- All you need for a thicker, greener lawn without dandelions
- [6 in 1!][5 in 1!][4 in 1!][3 in 1!][2 in 1!][for use with the corresponding number of the following pests]: Controls [Disease*], [Weeds], [Moss], [Algae*] and [Lichens] in your lawn [while also fertilizing!]
- 6 in 1! Fertilizes your lawn while controlling weeds, moss, disease*, algae and lichens!
- 3 in 1- Weeds, feeds, and controls diseases
- Treats up to XXXX ft² [sq. ft] [of lawn]
- Value [package]. Treats xxx square feet
- xx lb container covers up to xxx sq. ft.
- Covers X,XXX square feet/sq. ft.
- Treats [and]/[Feeds][Fertilizes] XXXX sq. ft.
- Tip: 1,000 sq. ft. = 20 ft. x 50 ft
- Works in temperatures [between] [from] 50°F and 90°F [10°C and 32°C]
- Works in cool weather [down to 50°F]
- Low temperature weed control [down to 50°F]
- [Even][also] works in cool weather
- Works fast
- See results in 24 hours
- Weeds start dying within 24 hours
- Fast and effective
- Eliminates unwanted moss, algae* and lichens* quickly
- Kills [weeds], [moss], [disease*], [algae*], and [lichens*] [quickly and effectively] [fast]
- Fast Acting!
- visible overnight results
- For a healthier greener lawn
- Fertilizes the lawn with Iron, a plant nutrient
- For a greener lawn in one week
- Formulated with Iron to help build a thick, green lawn from the roots up
- Greening power in just one week
- Promotes a green dandelion free lawn
- Effective for lawn spot treatments
- Kills [weeds][dandelions][&] [clover], down to the root
- 3x1 [action] [Herbicide, Fungicide & Fertilizer]
- Triple Action Formula [Herbicide, Fungicide & Fertilizer]
- Can be used on new lawns prior to [grass emergence] [germination]
- Can be used on newly [reseeded][overseeded][seeded][planted] lawns [prior to germination]
- Can apply [1] [one] day after [planting][seeding][reseeding][overseeding] lawn
- Can apply [1] [one] day before [planting][seeding][reseeding][overseeding] lawn
- Lawn areas can be [re-sown] [overseeded] one day after treatment
- Can [re-seed] [seed][overseed][plant] lawn 1 day after applying
- Apply, and [reseed][overseed][plant][seed] lawn the next day
- Weed killer for newly planted lawn
- Spring and fall weed control
- For a healthy lawn

- Patent Pending
- Rainfast in 24 hours
- No unpleasant odor
- Derived from iron [Fe]
- For non-selective [moss] weed and grass control, use [H01 or NEU1171H – or Finalsan Weed & Grass Killer or Fireworxx 80]
- Can be used to spot treat problem areas [in lawns]
- [Helpful Hints:] Best results are obtained when applied in the Spring or Fall when environmental conditions are cool and there is ample moisture
- [Helpful Hints:] The best times to treat weeds are in the Spring when they are small and growing and in the Fall to eliminate many of next year’s weeds
- People and pets can enter treated area after granules are no longer visible (approximately 6 hours)
- Clears out [clover], [dandelion] and other broadleaf weeds and helps prepare your lawn for winter
- Promotes healthy, weed- and disease*-free lawns
- Broadcast application recommended
- Apply in early spring for crabgrass prevention
- Prevents crabgrass [for up to 6 weeks]
- Stop crabgrass before it [starts] [invades] [overtakes] [your lawn]
- [In warm winter climates,] a fall application can prevent fall and winter weeds from germinating
- Builds strong, deep roots
- *not for use in California {for use with any marketing claim containing an asterisk}
- Before/After [application]
- Before/After [application] -
-**READ THE LABEL FIRST!**
- [Open box] [Box contains 2 [containers][bags][bottles] of product]
- Refer to leaflet for complete [First Aid], [Directions for Use], [Precautionary Statements], and [Storage and Disposal Instructions]
- [See][refer to] [inside][side][back][panel][attached booklet][container label] for [additional] [complete] [precautionary statements], [directions for use], [application instructions], [first aid statements], [storage and disposal], [use precautions] {Note to reviewer - When the first aid statement will appear on the side or back panel, either the claim “See side panel for First Aid” or “See back panel for First Aid” will appear on the front panel}
- For questions or comments, call [toll free] 1-XXX-XXX-XXXX [][][] [or visit us online at www.neudorff.com []
- For emergency medical information, you may call [toll free] 1-XXX-XXX-XXXX [][][] [or visit us online at www.neudorff.com []
- Satisfaction guaranteed


- Easy-to-use (ready-to-use) (RTU) dry formulation
- Easy to apply (dry product) (granules).
- The product is easy to use, clean to handle and economical.

-NOTE: This package is sold by weight. Contents may have settled during shipment.

-This container is made from XX% recycled materials.

-Cut Here To Open

-Convenient. Easy-to-use. Requires no mixing or spraying.

QUICK FACTS	
{Insert list from optional statement section}	
CONTROLS	<ul style="list-style-type: none">• [Dandelion], [Common Chickweed[§]], [Clover] and other Listed Broadleaf Weeds• [Moss], [Algae]* and [Lichens*][Liverworts*]• [Rust]*, [Dollar Spot]* and [Snow Mold]*
WHERE TO USE	<ul style="list-style-type: none">• On lawns [do not apply to bentgrass]
	For questions or comments, call toll-free [insert subregistrants toll-free number].

* Not for use in California

§ only approved for suppression in California

{Optional warranty statements. Registrant or distributor has choice of using any or none.}

-NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

{OR}

-Buyer assumes all responsibility for safety and use not in accordance with directions.

{OR}

-Unconditionally Guaranteed by (company name). **OUR COMMITMENT TO YOUR SATISFACTION:** If you are not happy with the way this product works in your garden, please let us know. We will be happy to replace the product, refund your money, or substitute another (brand name) product. Simply send your name, address and proof of purchase to (Company name & address).

{OR}

(company name) guarantees if for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

{OR}

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

{OR}

Satisfaction guaranteed (or your money back)

]

Registrant: W. Neudorff GmbH KG
An der Muhle 3,
31860 Emmerthal, Germany
www.neudorff.com

SUBLABEL B – COMMERCIAL USE

NEU1572H

Active Ingredient:

Sodium Ferric EDTA.....	14.00%
Other Ingredients:.....	86.00%
Total:.....	100.00%

EPA REG. NO. 67702-63

EPA EST. 67702-DEU-

1

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. {Optionally, supplemental distributors may use the following wording and own toll-free hotline number:} [For medical emergency information, [you may call toll free 1-XXX-XXX-XXXX] {insert operating hours and days of service}[During other times,] call the poison control center at 1-800-222-1222.]

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

{Latin names are optional throughout directions for use}

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable directions and precautions on this label before using.

SITES: Lawns, turf (use on rights of way or non-crop areas), golf courses, parks, playgrounds, cemeteries, athletic fields

Pests:

Controls or suppresses weeds such as but not limited to:

- Black medic (*Medicago lupulina*)[§]
- Broadleaf plantain (*Plantago major*)*
- Bull thistle (*Cirsium vulgare*)
- Canada thistle (*Cirsium arvense*)*

- Common chickweed (*Stellaria media*)[§]
- Crabgrass (*Digitaria spp.*)- pre-emergent control only
- Creeping buttercup (*Ranunculus repens*)
- Creeping Charlie/Ground ivy (*Glechoma hederacea*)*
- Dandelion (*Taraxicum officinale*)
- Dovefoot geranium (*Geranium molle*)*
- English daisy (*Bellis perennis*)[§]
- False dandelion (*Hypochoeris radicata*)*
- Heal-all (*Prunella vulgaris*)*
- Lawn burweed (*Soliva ptero-sperma*)*
- Narrowleaf plantain (*Plantago lanceolata*)*
- Oxalis (*Oxalis spp.*)*
- Prostrate knotweed (*Polygonum aviculare*)*
- Shepherd's-purse (*Capsella bursa-pastoris*)*
- Silverweed cinquefoil (*Potentilla anserina*)*
- *Speedwell spp.*(including Persian Speedwell (*Veronica persica*) and Slender speedwell (*Veronica filiformis*)*)
- White clover (*Trifolium repens*)
- Wild chamomile (*Matricaria chamomilla*)*
- Wild geranium (*Geranium robertianum*)*
- moss, algae*, lichens*, liverworts*

Controls or suppresses diseases such as but not limited to:

- Dollar Spot*
- Rust*
- Snow Mold*

* Not for use in California

§ only approved for suppression in California

Application Directions: Apply at a rate of 3-8lbs /1000 ft² (0.5 -1.15 oz /yard²), or 5-8 lbs/1000 ft² (0.72 -1.15 oz /yard²) in California. {Optional:} [The standard rate for this product is 5 lbs /1000 ft² (0.72 oz /yard²); use the higher rate of 8 lbs /1000 ft² (1.15 oz /yard²) on tough perennial weeds, such as common chickweed[§], buttercup and broadleaf plantain*. Use at 3 lbs /1000 ft² † (0.5 oz /yard²) on young moss, and weed seedlings of false dandelion*, English daisy*, dandelion* or white clover*.] Repeat treatment in 3 to 4 weeks for best results.

For Spot Treatment, apply 5 lbs/1000 ft². Repeat treatment in 3 to 4 weeks for best results.

For Pre-emergent control of broadleaf weeds and crabgrass, begin application when soil temperatures reach 50 to 55 °F. Apply 5-8 lbs/1000 ft², depending on weed pressure. Repeat treatment in 3 to 6 weeks for best results.

{Optional table}

Application rate	Weeds	Diseases	{Optional column} [Area Treated
3 lbs/ 1000 ft ² (0.43 oz/yd ²) †	Weed seedlings of English daisy* (<i>Bellis perennis</i>), dandelion* (<i>Taraxacum officinale</i>) false dandelion* (<i>Hypochaeris radicata</i>), young moss	Dollar spot *, snow mold*, and rust*. Use the higher rate when disease pressure is high.	xxxx ft ² /xxx lb container
4 lbs/ 1000 ft ² (0.58 oz/yd ²) †	Moss, speedwell species, including Persian speedwell, (<i>Veronica persica</i>) and slender speedwell* (<i>Veronica filiformis</i>), wild geranium* (<i>Geranium robertianum</i>)		xxxx ft ² /xxx lb container
5 lbs/ 1000 ft ² (0.72 oz/yd ²)	Black medic (<i>Medicago lupulina</i>) §, bull thistle (<i>Cirsium vulgare</i>), Canada thistle* (<i>Cirsium arvense</i>), creeping charlie/ground ivy* (<i>Glechoma hederacea</i>), dandelion (<i>Taraxacum officinale</i>), dove's foot geranium* (<i>Geranium molle</i>), English daisy (<i>Bellis perennis</i>), false dandelion* (<i>Hypochaeris radicata</i>), heal-all* (<i>Prunella vulgaris</i>), lawn burweed* (<i>Soliva pterosperma</i>), narrowleaf plantain* (<i>Plantago lanceolata</i>), oxalis* (<i>Oxalis</i> spp.), prostrate knotweed* (<i>Polygonum aviculare</i>), Shepherd's-purse* (<i>Capsella bursa-pastoris</i>), silverweed cinquefoil* (<i>Potentilla anserina</i>), algae*, lichens*, liverworts*. Preemergent control of Crabgrass (<i>Digitaria</i> spp.)		xxxx ft ² /xxx lb container
8 lbs/ 1000 ft ² (1.15 oz/yd ²)	Common chickweed [§] (<i>Stellaria media</i>), creeping buttercup (<i>Ranunculus repens</i>), broadleaf plantain* (<i>Plantago major</i>), white clover* (<i>Trifolium repens</i>).		xxxx ft ² /xxx lb container]

† Lower application rate not for use in California. Minimum application rate in California is 5 lbs/ 1000 ft²

* Not for use in California

§ only approved for suppression in California

]

How to Apply:

Contents may have settled during shipment; invert bag several times to ensure product is mixed.

For best results, apply a few days after mowing when temperature is between 50°F and 85°F. If temperature is below 80°F ensure turf is wet from dew, or water turf lightly before application. If temperature is above 80°F do not pre-wet grass and apply product in the late afternoon or evening. Do not exceed the 5 lbs/1000 ft² rate when temperatures are above 85°F. Do not apply when daytime temperature exceeds 90°F.

Apply using a broadcast/rotary, drop, or handheld spreader set to the correct application rate. Uniform coverage is important to prevent turf damage. See table below for spreader settings. Wait at least 24 hours after application to water turf. Turf areas can be re-sown one day after treatment.

Apply product only on turf. Application on hard surfaces can cause staining. If product gets on surface, sweep off if the surface is dry, rinse off immediately if the surface is wet. If product gets on clothing or footwear, wash immediately.

[Spreader settings:

Weed Species	Rate	{Optional column} Scotts® Broadcast and Rotary Spreader Setting (All Models)	{Optional column} Scotts® Drop Spreader Setting (All Models)	{Optional column} Scotts® Hand-Held Spreader Setting (All models)
Weed seedlings of English daisy* (<i>Bellis perennis</i>), dandelion* (<i>Taraxacum officinale</i>) false dandelion* (<i>Hypochaeris radicata</i>), young moss	3 lbs per 1000 square feet	2.5	5	3
Black medic (<i>Medicago lupulina</i>) [§] , bull thistle (<i>Cirsium vulgare</i>), Canada thistle* (<i>Cirsium arvense</i>), creeping charlie/ground ivy* (<i>Glechoma hederacea</i>), dandelion (<i>Taraxacum officinale</i>), dove's foot geranium* (<i>Geranium molle</i>), English daisy (<i>Bellis perennis</i>), false dandelion* (<i>Hypochaeris radicata</i>), heal-all* (<i>Prunella vulgaris</i>), lawn burweed* (<i>Soliva pterosperma</i>), narrowleaf plantain* (<i>Plantago lanceolata</i>), oxalis* (<i>Oxalis</i> spp.), prostrate knotweed* (<i>Polygonum aviculare</i>), Shepherd's-purse* (<i>Capsella bursa-pastoris</i>), silverweed cinquefoil* (<i>Potentilla anserina</i>), speedwell species, including Persian speedwell, (<i>Veronica persica</i>) and slender speedwell* (<i>Veronica filiformis</i>), wild geranium* (<i>Geranium robertianum</i>), moss, algae*, lichens*, liverworts*. Preemergent control of	5 lbs/1000 square feet	3	6	3.5

crabgrass (<i>Digitaria spp.</i>).				
Common chickweed [§] (<i>Stellaria media</i>), creeping buttercup (<i>Ranunculus repens</i>), broadleaf plantain* (<i>Plantago major</i>), white clover* (<i>Trifolium repens</i>).	8 lbs/1000 square feet	4.5	7.5	5

† Lower application rate not for use in California. Minimum application rate in California is 5 lbs/1000 ft²

* Not for use in California

§ only approved for suppression in California

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions. Spreaders other than those listed in this table may be used, as long as they are calibrated prior to use to deliver the product within the use rates on this label.]

What will I see?

Works quickly; dying weeds will be seen within 24 hours of application. Treated weeds, moss, or algae* will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however, the grass will recover within a few days to weeks. May stain thatch (dead grass under turf); this does not affect the health of the grass. If turf is damaged, irrigation 24 hours after application, followed by mowing to remove the turf tip burn can help turf recover faster.

* Not for use in California

Use Precautions:

Do not apply to drought stressed grass; ensure turf is well watered prior to application. Do not apply when the daytime temperature will exceed 90°F. Do not apply to bentgrass. If unsure of grass sensitivity, make a test application to a small area of the turf. Avoid applying to desirable plants. Do not apply if rainfall is expected within 24 hours of application. Do not irrigate within 24 hours of application.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep this product in its original container and in a secured area that is inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

Plastic containers: Triple rinse or equivalent. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Paper and plastic bags: Completely empty into application equipment. Then dispose of bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BATCH CODE:

{Optional table}

[

Guaranteed Fertilizer Analysis XX-XX-XX {insert N-P-K ratio}	
Total nitrogen (N) X% Ammoniacal nitrogen X% Urea nitrogen X% Other water soluble nitrogen X% Water insoluble nitrogen	X%
Available phosphate (P ₂ O ₅)	X%
Iron (Fe)	X%
Sulfur (S)	X%
Magnesium (Mg) (Total) X% water soluble magnesium X% chelated magnesium	X%
Sulfur X% combined sulfur X% free sulfur	X%
Soluble potash (K ₂ O)	X%
Calcium (Ca)	X%
Derived From: {Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer.}	

]

{Note to reviewer: Spanish label text may also be included in addition to English label text.}

{The following claims and product information, may or may not be presented on the product's label and labeling:}

[

{For products being sold into states requiring a metals internet statement on fertilizer products, the following statement will be used:}

-Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

{For products being sold into Florida, the following statements will be used:}

-FL License No.: FXXXXXX

-Florida Applications:

We recommend that you follow the Florida Green Industries BMP's at:

<http://www.flrules.org/Gateway/reference.asp?No=Ref-04706>

We recommend that you follow the Florida Golf Course BMP's at:

<http://www.flrules.org/Gateway/reference.asp?No=Ref-04702>

{For states with nitrogen application restrictions, including but not limited to, MD, NH, NJ, PA and VA, the following statement will be used:}

-Nitrogen Notice: Do not apply more than X.XX lbs of this product per 1,000 sq.ft.

-Phosphate Notice: This lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.

-www.neudorff.com



- [Moss], [weed], [algae*], [lichen*], [liverwort*] and [disease*] killer [for turf]
- Turf [Moss], [weed], [algae*], [lichen*], [liverwort*] and [disease*] [Killer] [Control]
- [Selective] [Broadleaf] [Weed], [algae*], [lichen*], [liverwort*] and [disease*] [Killer][Control]
- [Selective Control of][Controls] [Broadleaf] Weeds in Turf
- Kills [moss], [weed], [algae*], [lichen*], [liverwort*] and [diseases*], not [turf][grass]
- Kills [moss], [weed], [algae*], [lichen*], [liverwort*] and [diseases*] without harming turf
- Kills [weeds], [algae*], [moss], [liverworts*] and [lichens*]
- Easy-to-use [moss], [weed], [algae*], [lichen*], [liverwort*] and [disease*] killer for the turf
- Cure your [weed], [moss] and [disease*] problems fast
- Ideal [weed], [moss] & [disease*] control for turf
- Iron-based [disease], [moss], [disease*] & [weed] control for turf
- Controls dollar spot*, snow mold*, and rust* [on turf]
- Turf [Fungicide*], [Herbicide] and [Fertilizer]
- Kills [dandelions], [clover] and [most] other [major][broadleaf] turf weeds* *as listed
- Prevents and controls [Moss], [weed], [algae*], [lichen*], [liverwort*], and [turf diseases*]
- [Controls] [kills] [dandelions] [buttercup] [English daisy] [daisy] [clover]
- Controls disease* in the turf, kills the weeds.
- [Disease*], [moss] and [Weed] Killer in one!
- Tough on [moss] [weeds] and [disease*], easy on turf
- Prevent disease* and kill the weeds!
- Kills most common turf diseases* fast!
- Kills [weeds][dandelions], down to the root
- NEU1572H is derived from iron that is used to selectively control/ suppress [broadleaf weeds], [moss], and [diseases*] growing in turf
- Herbicide and Fungicide* in 1 for the Turf!

-NEU1572H is formulated to feed your turf and control [broadleaf weeds], [moss], [algae*], [diseases*] commonly found in turf.

-Pre- and Post-emergent activity of broadleaf weeds with one application

-Prevents crabgrass and broadleaf weeds from germinating

-Pre and post emergent herbicide

-Pre and post emergent weed control for turf

-Controls weeds before they sprout

-Provides pre-emergent control of weeds in turf

-Pre-emergent control of [broadleaf weeds] and [crabgrass] [in turf]

-Controls weeds before you see them (!)

-Apply pre-emergence for extended weed control

-Controls Weeds and Feeds your Turf.

-Builds thick, green turf from the roots up

-Builds thick, green turf while killing listed weeds

-Strengthens and protects your turf

-Greens up turf in 7 days

-Thickens turf to crowd out weeds

-[Fast] [quick] green-up

-Greens your turf fast

-Two [2] in One [1] fertilizer and [weed][moss] killer

-Kills [weeds] and [moss], feeds your turf

-Provides fast green-up of turf

-Contains Iron, a plant nutrient

-[Guaranteed Analysis]: xx-xx-xx {insert N-P-K ratio} [Fe 1.82%]

-[X-X-X {insert N-P-K ratio}] [1.82% Fe] Nitrogen, Phosphorus, Potash / Potassium, [Iron]

-[N] Nitrogen [and [Fe] Iron] for greening [P] Phosphorus helps develop a stronger, hardier root system [K] Potassium / Potash improves plant hardiness and drought resistance

-Healthier, stronger and greener turf

-WEED AND FEED

-All you need for a thicker, greener turf without dandelions

-[6 in 1!] [5 in 1!] [4 in 1!] [3 in 1!] [2 in 1!] [for use with the corresponding number of the following pests]: Controls [Disease*], [Weeds], [Moss], [Algae] and [Lichens] in your turf [while also fertilizing!]

-6 in 1! Fertilizes your turf while controlling weeds, moss, disease*, algae* and lichens!

-3 in 1- Weeds, feeds, and controls diseases*

-Treats up to XXXX ft² [sq. ft] [of turf]

-Value [package]. Treats xxx square feet

-xx lb container covers up to xxxx sq. ft.

-Covers X,XXX square feet/sq. ft.

-Treats [and]/[Feeds][Fertilizes] XXXX sq. ft.

-Tip: 1,000 sq. ft. = 20 ft. x 50 ft

-Works in temperatures [between] [from] 50°F and 90°F [10°C and 32°C]

-Works in cool weather [down to 50°F]

-Low temperature weed control [down to 50°F]

-[Even][also] works in cool weather

- Works fast
- See results in 24 hours
- Weeds start dying within 24 hours
- Fast and effective
- Eliminates unwanted moss, algae* and lichens* quickly
- Kills [weeds], [moss], [disease*], [algae*], and [lichens*] [quickly and effectively] [fast]
- Fast Acting!
- Can be used on new turf prior to [grass emergence] [germination]
- Can be used on newly [reseeded][overseeded][seeded][planted] turf [prior to germination]
- Can apply [1] [one] day after [planting][seeding][reseeding][overseeding] turf
- Can apply [1] [one] day before [planting][seeding][reseeding][overseeding] turf
- Turf areas can be [re-sown] [overseeded] one day after treatment
- Can [re-seed] [seed][overseed][plant] turf 1 day after applying
- Apply, and [reseed][overseed][plant][seed] turf the next day
- Weed killer for newly planted turf
- Spring and fall weed control
- For a healthy turf
- Patent Pending
- Rainfast in 24 hours
- No unpleasant odor
- Derived from iron [Fe]
- For non-selective [moss] weed and grass control, use [H01 or NEU1171H – or Finalsan Grass & Weed Killer or Fireworxx 80]
- Can be used to spot treat problem areas [in turf]
- [Helpful Hints:] Best results are obtained when applied in the Spring or Fall when environmental conditions are cool and there is ample moisture
- [Helpful Hints:] The best times to treat weeds are in the Spring when they are small and growing and in the Fall to eliminate many of next year's weeds
- People and pets can enter treated area after granules are no longer visible (approximately 6 hours)
- Clears out [clover], [dandelion] and other broadleaf weeds and helps prepare turf for winter
- Promotes healthy, weed- and disease*-free turf
- Broadcast application recommended
- Apply in early spring for crabgrass prevention
- Prevents crabgrass for up to 6 weeks
- [In warm winter climates,] a fall application can prevent fall and winter weeds from germinating
- Builds strong, deep roots
- Before/After [application] **-READ THE LABEL FIRST!**
- *not for use in California {to be used with any marketing claim that has an asterisk}
- [Open box] [Box contains 2 [containers][bags][bottles] of product]
- Refer to leaflet for complete [First Aid], [Directions for Use], [Precautionary Statements], and [Storage and Disposal Instructions]

- [See][refer to] [inside][side][back][panel][attached booklet][container label] for [additional] [complete] [precautionary statements], [directions for use], [application instructions], [first aid statements], [storage and disposal], [use precautions] {Note to reviewer - When the first aid statement will appear on the side or back panel, either the claim "See side panel for First Aid" or "See back panel for First Aid" will appear on the front panel}
- For questions or comments, call toll free 1-XXX-XXX-XXXX [or visit us online at www.neudorff.com]
- For emergency medical information, you may call toll free 1-XXX-XXX-XXXX. [or visit us online at www.neudorff.com]
- Satisfaction guaranteed
- Easy-to-use (ready-to-use) (RTU) dry formulation
- Easy to apply (dry product) (granules).
- The product is easy to use, clean to handle and economical.
- Cut Here To Open
- NOTE:** This package is sold by weight. Contents may have settled during shipment.
- This container is made from XX% recycled materials.
- Convenient. Easy-to-use. Requires no mixing or spraying.

{Optional warranty statements. Registrant or distributor has choice of using any or none.}

-NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

{OR}

-To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

{OR}

-Unconditionally Guaranteed by (company name). **OUR COMMITMENT TO YOUR SATISFACTION:** If you are not happy with the way this product works in your garden, please let us know. We will be happy to replace the product, refund your money, or substitute another (brand name) product. Simply send your name, address and proof of purchase to (Company name & address).

{OR}

(company name) guarantees if for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

{OR}

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

]

Registrant: W. Neudorff GmbH KG
An der Muhle 3,
31860 Emmerthal, Germany
www.neudorff.com