

PM 92

67727-1

6/29/99

1 of 3

Master Label 1,4SIGHT®

Aerosol Grade - Potato Dormancy Enhancer

Product Benefits:

- Naturally occurring plant hormone delays sprouting, so smaller buds on potatoes; [and/or]
- Enhances the potato's ability to heal cuts and bruises; [and/or]
- Reduces respiration, resulting in less water loss; [and/or]
- Reduces Refrigeration and ventilation needs; [and/or]
- Reduces pressure bruising; [and/or]
- Firmer potatoes with improved visual appearance.

Active Ingredient: 1,4-Dimethylnaphthalene*	By Weight
Inert Ingredients:	94.7%
TOTAL	5.3%
	100.0%

* Contains 7.9 pounds active ingredient per gallon.

Keep Out of Reach of Children

WARNING

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Wear protective eyewear (goggles or face shield). Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Statement of Practical Treatment

- If in Eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.
- If Swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol.
- If on the Skin: Wash with plenty of soap and water. Get medical attention.

Emergency Information

For spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300



Environmental Hazards

This product is highly toxic to freshwater fish and aquatic invertebrates. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Net Contents _____

Manufactured For:
D-I-1-4, Inc.
P.O. Box 860
Meridian, ID 83680 USA

EPA Reg. Number 67727-1
EPA Est. Number 68738-JPN-001
Made in Japan
Revision D-3-S



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

- Keep container closed. This product temporarily inhibits germination of seed potatoes.

PESTICIDE DISPOSAL

- Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

- Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not reuse empty container.

Continue Directions for Use on Reverse

NOTICE


- 1,4SIGHT® is used as an aerosol to enhance the dormancy of potatoes during the storage phase.
- Applicators and other handlers must wear: long-sleeved shirt, long pants, shoes, socks, protective eyewear (goggles or face shield), and chemical resistant (such as nitrile or butyl) gloves.
- For reentry into treated areas during application and prior to ventilation or settling of aerosol fog, workers must wear: coveralls over long-sleeved shirt, long pants, shoes plus socks, and chemical resistant (such as nitrile or butyl) gloves; face-sealing goggles, unless a full-face respirator is worn; and a respirator with an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE prefilter.
- 1,4SIGHT® must not be applied to potatoes in the field.
- 1,4SIGHT® may be applied anytime after the potatoes are placed into the storage area.
- 1,4SIGHT® is a short term dormancy enhancer which should not adversely affect eventual germination of seed potatoes.
- Let 60 days elapse before using a treated storage area as a temporary holding area for seed potatoes. Air system components (including ducts) and building must be thoroughly ventilated before area is used for storage of seed potatoes to prevent dormancy effects.
- Do not allow vapors to come in contact with storage areas used for seed potatoes within 60 days of their planting.

FORCED AIR DISTRIBUTION METHOD

1. Prepare aerosol generating equipment. The discharge from the vaporizer should be located to provide optimum distribution in the storage area. Take precautions to prevent the spray from coming into direct contact with the potatoes.
2. Running fans during the application may aid disbursement of vapors, but is not required.
3. It is advantageous to keep the storage area closed as long as practical to allow for maximum absorption into the potatoes.

TREATMENT OF STORAGE AREAS THAT DO NOT HAVE RECIRCULATING AIR SYSTEMS

- These storage areas are more difficult to treat. Additional steps must be taken to treat these storage areas.
1. Prepare aerosol generating equipment. The discharge from the vaporizer should be located to provide



optimum distribution in the storage area. Take precautions to prevent the spray from coming into direct contact with the potatoes.

2. Completely close off the area to be treated. This may include vent doors in the attic and walk doors around the perimeter. The large truck doors may need to be sealed to prevent excess leakage.
3. Begin treatment by following the listed steps:
 - A. Start the aerosol generator.
 - B. Adjust the location of the vaporizer so all the storage area receives an equal amount of aerosol fog. It may be necessary to move the vaporizer during the application or make the application in multiple phases. For reentry into treated areas during application and prior to ventilation or settling of the aerosol fog, workers must wear: coveralls over long-sleeved shirt, long pants, shoes plus socks, and chemical resistant (such as nitrile or butyl) gloves; face-sealing goggles, unless a full-face respirator is worn; and a respirator with an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE prefilter.
4. Wait at least 12 hours before fresh air is introduced into the storage area, then return the storage area to "normal" operation.
5. It is advantageous to keep the storage area closed as long as practical to allow for maximum absorption into the potatoes.

APPLICATION TIMING AND RATES

Application of 1,4SIGHT® can be made anytime after the potatoes are placed in the storage area.

1. Apply at rates of up to 1 pound of active ingredient per 50,000 lbs (500 cwt), which is equivalent to 20 ppm on a product to potato basis or 1 gallon per 4,000 cwt.
2. While potatoes may be retreated as necessary for effective control, do not exceed the maximum application of 80 ppm during the storage season.

1 cwt. = 100 lbs = 2.5 cubic feet; 1 ton = 20 cwt.

CONDITIONS OF SALE

D-I-1-4, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to manufacturer, and buyer assumes the risk for any such use.

