

FYFANON 6% DUST
FOR GRAIN PROTECTION
For Use on Stored Grains
Wheat - Corn - Oats - Rye - Barley

ACCEPTED
DEC 16 1997
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **67760-26**

ACTIVE INGREDIENT:

Malathion (0-0-dimethyl phosphorodithioate of diethyl mercaptosuccinate).....	6.0%
INERT INGREDIENTS	94.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

IN CASE OF MEDICAL EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 800-228-5635 EXT. 153

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.

NOTE TO PHYSICIAN: Malathion, upon repeated, prolonged and careless exposure, may cause cholinesterase inhibition. Atropine is antidotal.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Do not take internally. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling. Wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in areas near application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

EPA Reg. No. 67760-26

96 AUG -7 A10:46

REC'D EPA/OPP/DPD1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in original labeled container, in cool, dry, locked place out of the reach of children. In case of spills, remove spillage with an industrial type vacuum cleaner and wash area with household bleach.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: This product may be used on selected food grain. Use of this product as directed will be within the tolerance of 8 parts per million for Malathion. Treated grain is not subject to down-grading and can be milled into flour or feed using appropriate industry methods.

CLEAN ALL EMPTY BINS: Empty bins should be treated with a suitable residual insecticide before they are filled with new grain. Refuse, trash and old grain should be swept up and the entire area cleaned and made ready for new storage.

GRAIN PROTECTION: For use on stored grains (Wheat, Corn, Oats, Rye and Barley) for protection against the following insects: Granary weevil, rice weevil, maize weevil, flat grain beetle, saw-toothed grain beetle, confused flour beetle, red flour beetle, rusty grain beetle, foreign grain beetle, khapra beetle, lesser grain borer, larger grain borer, Indian meal moth larvae, Mediterranean flour moth larvae; and suppresses cadelles, grain mites and Angoumous grain moth larvae.

WHEN AND HOW TO APPLY

TREATMENT AT THE ELEVATOR: Use suitable powder application equipment to meter the product onto the grain as it enters the final storage at a rate of 10 lbs. per 1,000 bushels. Mount the device near the auger outlet to treat the flow of grain going onto the final storage bin. Wear a dust mask.

FOR TREATMENT OF GRAIN IN SHIPS HOLDS OR OTHER TRANSPORTATION FACILITIES: For intercontinental shipments of grain by boat or barge, apply this product in the same manner as for treatment of grain going into an elevator bin at the rate of 10 lbs. per 1,000 bushels of grain. Use appropriate application equipment.

GRAIN SURFACE TREATMENT AGAINST INDIAN MEAL MOTH LARVAE AND OTHER INSECTS:

To protect stored grain from attack by Indian meal moth larvae and other stored grain insects in the top 8 to 10 inches of grain, apply this product to the surface as a preventative or curative treatment at the rate of 5 lbs. per 1,000 square feet of grain surface. Apply evenly over the surface of grain after it has been loaded into final storage. Begin application at the far end of the bin. Apply at the recommended rate and rake into the grain surface as you backup towards the exit. Do not walk through treated grain. Check grain in 3 to 4 weeks to determine presence of insects. Repeat treatment as needed. Wear a dust mask during application.