

67760-68

06/25/2009

1/33



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

AND

OFFICE OF
PREVENTION, PESTICIDES
TOXIC SUBSTANCES

JUN 25 2009

Ms. Carrie M. Tackema
Cheminova, Inc
P.O. Box 110566
One Park Drive, Suite 150
Research Triangle Park, NC 27709

Dear Ms. Tackema:

Subject: Accurate Herbicide (Update to Agree with Similar Products)
EPA Registration No. 67760-68
Application Dated April 2, 2009

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable provided you make the following changes before you release the product for shipment.

1. On page 4, under General Information, delete the word "recommended" from the first and second sentence. The sentences may be modified as necessary.
2. On page 5, revise the first sentence to read "If those **instructions** conflict with this label do not tank mix that product with Accurate.
3. There are several areas of the label where the phrase "recommended rate" appears. Revise these phrases to read "**labeled rate**".
4. On page 9, under Tank Mixtures in Cereals (Wheat, Barley and Triticale), revise the first and second sentences to read "Read and follow all manufacturer's **instructions** for any companion herbicides, fungicides, and/or insecticides. If those **instructions** conflict with this label, do not tank mix that product with Accurate.
5. There are several areas of the label where the phrase "local recommendations" appears. Revise these phrases to read "local **guidance**".
6. On pages 11 and 15, under With Liquid Nitrogen Solution Fertilizer, revise the last sentence in the fourth paragraph to read "Consult your agricultural dealer...for specific **information** before adding an adjuvant to these tank mixtures."

Page 2

EPA Registration No. 67760-68

7. There are several places where a phrase similar to "to rotate to a (rotational) crop at a shorter interval than recommended appears. Revise those phrases to read "To rotate...at a shorter interval than **specified**".
8. On page 13, under Grain Sorghum, delete the word "recommended" from the first sentence. The sentence may be modified as necessary.
9. On page 16, under checking soil pH, delete the word "recommended" from the last sentence.
10. On page 22, under Spray Equipment, revise the first sentence to read "For specific application equipment, refer to manufacturer's **instructions** for additional information..."
11. On page 22, under After Spraying Accurate and Before Spraying Crops Other Than Wheat, Barley, Fallow, Pasture, or Rangeland revise the first sentence in bullet 6 to read "If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) **listed** on this label."
12. In the supplemental label for Weed Control in Suagrcane in the State of Hawaii, delete the word "recommended" from the sentence under General Information. You may revise the sentence as necessary.
13. In the supplemental concerning the Conservation Reserve Program, delete the word "recommended" from the second sentence under Directions for Use.
14. In the supplemental concerning the Conservation Reserve Program, under Weed Control, revise the second and third sentences to read "Read and follow all manufacturer's **instructions** for the 2, 4-D used. If those **instructions** conflict with this label, do not tank mix the 2,4-D with Accurate."

Submit one copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling supersedes all previously approved ones. A stamped copy of labeling enclosed for your records.

Sincerely,

Vickie K. Walters for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

Accurate®

Herbicide

Dry Flowable

For Use on Wheat, Barley, Fallow, Pastures and Rangeland

ACTIVE INGREDIENT:

Metsulfuron Methyl

Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate... 60.0%

OTHER INGREDIENTS 40.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

IN CASE OF A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL TOLL FREE, DAY OR NIGHT 1-866-303-6950.

Read the entire label before using this product.

Use only according to label instructions.

Read "DISCLAIMER" before buying or using.

If terms are not acceptable, return product unopened without delay.

SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND USE DIRECTIONS


EPA Reg. No. 67760-68

EPA Est. No.

NET CONTENTS: 8 oz.

Cheminova, Inc.
One Park Drive, Suite 150
P.O. Box 110566
Research Triangle Park, NC 27709

Accurate® is a registered trademark of Cheminova A/S



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN
CAUTION**

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID

IF ON SKIN

OR CLOTHING: Take off contaminated clothing. Rinse skin with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-303-6950 for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

WPS USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard [40CFR Part 170] must wear: long sleeved shirt and long pants, shoes plus socks, and chemical resistant gloves from category A such as, butyl rubber, natural rubber, neoprene rubber, or nitrile rubber ≥14 mils.

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

IMPORTANT INFORMATION

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, shoes plus socks, and chemical resistant gloves made of any waterproof material.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

Weed control in pastures and rangeland is not within the scope of the Worker Protection Standard.

FOR MORE INFORMATION, CALL TOLL-FREE 1-800-548-6113

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

PRODUCT DISPOSAL:

Nonrefillable containers equal to or less than 5 gallons:

Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable containers greater than 5 gallons:

Do not reuse or refill this container. Offer for recycling if available. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Repeat this procedure two more times.

GENERAL INFORMATION

Accurate® is recommended for use on land primarily dedicated to the production of wheat, barley, fallow, pasture and rangeland.

Accurate is recommended for use on wheat, barley, fallow, pasture and rangeland in most states. Check with your state extension or Department of Agriculture before use to be certain **Accurate** is registered in your state.

Accurate is not registered for use in Alamosa, Conejos, Costilla, RioGrande, and Saquache counties of Colorado.

Accurate is a dry-flowable granule that controls weeds in wheat (including durum), barley, pasture, rangeland grasses, and fallow. **Accurate** is mixed in water or can be preslurried in water and added to liquid nitrogen carrier solutions and applied as a uniform broadcast spray. A surfactant should be used in the spray mix unless otherwise specified on this label. **Accurate** is noncorrosive, nonflammable, nonvolatile, and does not freeze.

Accurate controls weeds by postemergence activity. For best results, apply **Accurate** to young, actively growing weeds. The use rate depends upon the weed spectrum and size of weeds at application. The degree and duration of control may depend on the following factors:

- weed spectrum and infestation intensity
- weed size at application
- environmental condition at and following treatment.

Environmental Conditions and Biological Activity

Accurate is absorbed through the foliage of broadleaf weeds, rapidly inhibiting their growth. Leaves of susceptible plants appear chlorotic from 1 to 3 weeks after application and the growing point subsequently dies.

Application of **Accurate** provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

Accurate may injure crops that are stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, different varieties of the crop may be sensitive to treatment with **Accurate** under otherwise normal conditions. Treatment of such varieties may injure crops.

In warm, moist conditions, the expression of herbicide symptoms is accelerated in weeds; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to **Accurate**.

Weed control may be reduced if rainfall or snowfall occurs soon after application.

APPLICATION INFORMATION

FALLOW

Use Rates

Apply 1/10 oz. **Accurate** per acre.

Application Timing

Accurate may be used as a fallow treatment, in the spring or fall when the majority of weeds have emerged and are actively growing.

Tank Mixtures in Fallow

Accurate may be used as a fallow treatment, and may be tank mixed with other herbicides that are registered

for use in fallow. If those recommendations conflict with this label do not tank mix that product with **Accurate**. Read and follow all label instructions on timing, precautions, and warnings for any companion products before using these tank mixtures. Follow the most restrictive labeling.

WHEAT, BARLEY, and TRITICALE

Use Rates

Wheat (including durum), Barley and Triticale

Apply 1/10 oz. **Accurate** per acre to wheat, barley or triticale once per use season.

Pasture and Rangeland

Apply 1/10 to 4/10 oz. **Accurate** per acre as a broadcast treatment to pasture and rangeland. For spot applications, use 1 oz. per 100 gal. of water. Do not exceed 3/4 oz. **Accurate** per acre.

Harvest Aid

Apply 1/10 oz. **Accurate** per acre in combination with 2,4-D or Glyfos® to aid in dry down of many broadleaved weeds, thereby aiding grain harvest.

Application Timing

Dryland Wheat, Barley and Triticale

(Except Durum or Wampum Variety)

Make applications after the crop is in the 2-leaf stage but before boot once per use season.

Durum and Wampum Variety Spring Wheat

Make applications after the crop is tillering but before boot once per use season. Applications to durum and wampum varieties should be made in combination with 2,4-D.

Irrigated Wheat and Barley

Make applications after the crop begins tillering but before boot. First post-treatment irrigation should be delayed for at least 3 days after treatment and should not exceed 1 in. of water.

Wheat and Barley – Harvest Aid

Make applications after the crop has reached the hard dough stage, but no later than 10 days before harvest.

See section on Harvest Aid tank mixtures.

Fallow

Accurate may be used as a fallow treatment, in the spring or fall when the majority of weeds have emerged and are actively growing.

Do not apply during boot or early heading, as crop injury may result.

APPLICATION TIMING – PASTURE GRASSES

Accurate may be used on some native grasses such as bluestems and grama, and on other pasture grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy. Specific application information on several of these pasture grasses follows:

Pasture Grass	Minimum time from Grass establishment to Accurate application
Bermudagrass	2 months
Bluegrass, bromegrass and orchardgrass	6 months
Timothy	12 months
Fescue	24 months

Fescue Precautions:

Note that **Accurate** may temporarily stunt fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- tank mix **Accurate** with 2,4-D.
- use the lowest recommended rate for target weeds
- use surfactant at ½ to 1 pt per 100 gal of spray solution (1/16 to 1/8% v/v)
- make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall
- Do not use surfactant when liquid nitrogen is used as a carrier.

The first cutting yields may be reduced due to seedhead suppression resulting from treatment with **Accurate**.

Timothy Precautions:

Timothy should be at least 6" tall at application and be actively growing. Applications of **Accurate** to timothy under any other conditions may cause crop yellowing and/or stunting. To minimize these symptoms, take the following precautions:

- tank mix **Accurate** with 2,4-D
- use the lowest recommended rate for target weeds
- use surfactant at ½ pt per 100 gal (1/16% v/v)
- make applications in the late summer or fall
- do not use surfactant when liquid nitrogen is used as a carrier.

Ryegrass Pastures (Italian or perennial): Do not apply Accurate as injury to or loss of the pasture may result.

Other Pastures: Varieties and species of pasture grasses differ in their tolerance to herbicides. When using **Accurate** on a particular grass for the first time, limit use to one container. If no injury occurs throughout the season, larger acreage may be treated the following season.

Broadleaf pasture species, such as alfalfa and clover, are highly sensitive to **Accurate** and will be severely stunted or injured by **Accurate**

WEEDS CONTROLLED

Unless otherwise directed, treat when weeds are less than 4" tall or 4" across and are actively growing. Effectiveness may be reduced if rainfall occurs within 4 hours after application.

Cereals, Pasture, Rangeland, and Fallow**1/10 oz. per acre**

Blue/purple mustard*	Lambsquarters
Bur buttercup (testiculate)	(common, slimleaf)
Coast fiddleneck	Mayweed chamomile
(tarweed)	Miners lettuce
Common chickweed	Pigweed (redroot, smooth, tumble)
Common purslane	Plains coreopsis
Conical catchfly	Prickly lettuce*
Cowcockle	Russian thistle*
False chamomile	Shepherd's purse
Field pennycress	Smallseed falseflax
(fanweed)	Smartweed (green, lady's thumb, pale)
Filaree	Snow speedwell
Flixweed*	Tansymustard*
Groundsel (common)	Treacle mustard
Henbit	(Bushy Wallflower)
Kochia*	Tumble/Jim Hill mustard

Volunteer sunflower
Waterpod
Wild mustard

Additional Weeds In Pasture/Rangeland Only

1/10 to 2/10 oz. per acre

Bitter sneezeweed	Dandelion
Buttercup	Marestail
Carolina geranium	Plantain
Common broomweed	Wild garlic*
Common mullein	Woolly croton*
Curly dock	

2/10 to 3/10 oz. per acre

Annual marshelder	Horsemint (beebalm)
Blackeyed-Susan	Musk thistle*
Buckbrush [‡]	Pensacola bahiagrass
Burclover	Purple scabious
Common yarrow	Western snowberry [‡]
Dogfennel	Wild carrot

4/10 oz. per acre

Serecia lespedeza*

Weeds Suppressed[‡]*

Cereals, Pasture, Rangeland, and Fallow

1/10 oz. per acre

Canada thistle*	Knotweed (prostrate)*
Common sunflower*	Sowthistle (annual)*
Corn gromwell*	Wild buckwheat*

Brush Suppressed[‡]

3/10 oz. per acre

Blackberry	Multiflora rose*
Dewberry	

Weeds/Brush Suppressed with Spot Application (Pasture/Rangeland only)

1 oz. per 100 gal of water

Blackberry*	Dewberry*
Canada thistle*	Multiflora rose*

* See the **Specific Weed Problems** section

[‡] Weed suppression is a reduction in weed competition (reduced population and/or vigor) as visually compared to an untreated area. The degree of suppression varies with the rate used, the size of the weeds, and the environmental conditions following treatment.

Specific Weed Problems

NOTE: Thorough spray coverage of all weed species listed below is very important.

Blue Mustard, Flixweed, and Tansymustard: For best results, apply **Accurate** tank mixtures with 2,4-D

or MCPA postemergence to mustards, but before bloom.

Canada Thistle and Sowthistle: Apply either **Accurate** plus surfactant or **Accurate** plus 2,4-D or MCPA in the spring after the majority of thistles have emerged and are small (rosette stage to 6" elongating stems) and actively growing. The application will inhibit the ability of emerged thistles to compete with the crop.

For spot applications to Canada Thistle in pasture and rangeland, apply as a foliar spray once plant is fully leafed. Apply to runoff and include a surfactant in the spray mix at 1 to 2 qt per 100 gal of spray solution. Complete coverage of all foliage and stems is required for control. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage.

Corn Gromwell and Prostrate Knotweed: Apply **Accurate** plus surfactant when weeds are actively growing, are no larger than 2" tall, and when crop canopy will allow thorough coverage. Tank mixing 2,4-D or MCPA with **Accurate** can improve results.

Kochia, Russian thistle, Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use **Accurate** in a tank mix with "Banvel"/"Banvel" SGF and 2,4-D, or bromoxynil and 2,4-D (such as $\frac{3}{4}$ - 1 pt "Buctril" + $\frac{1}{4}$ - $\frac{3}{8}$ lb active 2,4-D ester). **Accurate** should be applied in the spring when kochia, Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing (refer to the Tank Mixtures section of this label for additional details).

Sunflower (common/volunteer): Apply either **Accurate** plus surfactant or **Accurate** plus 2,4-D or MCPA after the majority of sunflowers have emerged, are 2" to 4" tall and are actively growing. Use spray volumes of at least 3 gal by air or 5 gal by ground (10 gal by ground in pastures).

Wild Buckwheat: For best results, apply **Accurate** plus 2,4-D or MCPA when plants have no more than 3 true leaves (not counting the cotyledons). If plants are not actively growing, delay treatment until environmental conditions favor active weed growth.

Musk Thistle: Apply **Accurate** at 2/10 to 3/10 oz per acre in the spring or early summer prior to flowering or in the fall after newly emerged plants have reached the rosette stage of growth. Fall applications should be made before the soil freezes.

Multiflora Rose: For best control, apply **Accurate** as a broadcast application when multiflora rose is less than 3' tall. Application should be made in the spring, soon after multiflora rose is fully leafed.

For spot applications in pasture and rangeland, apply as a foliar spray once plant is fully leafed. Apply to runoff and include a surfactant in the spray mix at 1 to 2 qt per 100 gal of spray solution. Complete coverage of all foliage and stems is required for control. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage.

Blackberry and Dewberry: For spot applications in pasture and rangeland, apply as a foliar spray once plant is fully leafed. Apply to runoff and include a surfactant in the spray mix at 1 to 2 qt per 100 gal of spray solution. Complete coverage of all foliage and stems is required for control. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage.

Pensacola bahiagrass control in established Bermudagrass pasture: Apply **Accurate** at 3/10 oz per acre plus surfactant. Apply after green-up in the spring but before bahiagrass seedhead formation. Application should be made when moisture is sufficient to enhance grass growth.

Accurate is very effective for removal of bahiagrass from bermudagrass pastures. In highly infested pastures, the use of **Accurate** can clear the areas of useful forage until the bermudagrass has time to cover the area. Therefore, **Accurate** treatments should be spread out over a period of years. Do not apply to an entire farm or ranch in one year. Fertilization (particularly with nitrogen and potassium) and/or replanting may accelerate the process of reestablishment of bermudagrass.

Under heavy bahiagrass pressure, grazing pressure, or adverse weather conditions (heat and drought), bahiagrass regrowth may occur.

NOTE: **Accurate** should not be used for the control of common or Argentine bahiagrass. Also, **Accurate** should not be applied in liquid fertilizer solutions for Pensacola bahiagrass control, as poor control and/or regrowth may occur.

Serecia lespedeza: Apply **Accurate** at 4/10 oz per acre plus a surfactant at 1 to 2 qt per 100 gal of total spray solution. For best results, make applications to serecia lespedeza beginning at flower bud initiation through the full bloom stage of growth.

NOTE: Do not make applications if drought conditions exist at intended time of application.

Wild Garlic: Apply 1/10 to 2/10 oz per acre of **Accurate** in the early spring when wild garlic is less than 12" tall with 2" to 4" of new growth.

Woolly Croton: Apply 1/10 to 2/10 oz per acre of **Accurate** in the late spring or early summer at preemergence through 2 true leaf stage.

TANK MIXTURES IN CEREALS (WHEAT, BARLEY AND TRITICALE)

Read and follow all manufacturers' label recommendations for any companion herbicides, fungicides, and/or insecticides. If those recommendations conflict with this label, do not tank mix that product with **Accurate**. Read and follow all label instructions on timing, precautions, and warnings for any companion products before using these tank mixtures. Follow the most restrictive labeling.

Accurate may be tank mixed with other suitable registered herbicides to control weeds listed under **Weeds Suppressed**, weeds resistant to **Accurate**, or weeds not listed under **Weeds Controlled**.

With 2,4-D (amine or ester) or MCPA (amine or ester)

Accurate can be used as a tank mix treatment with 2,4-D or MCPA (ester formulations provide best results) herbicides after weeds have emerged. For best results, use 1/10 oz. of **Accurate** per acre: add 2,4-D or MCPA herbicides to the tank at ¼ to ½ lb. active ingredient. Surfactant may be added to the mixture at ½ to 1 qt. per 100 gallons of spray solution; however, adding surfactant may increase the potential for crop injury.

Apply **Accurate** plus MCPA after the 3 to 5-leaf stage but before boot (with Durum and Wampum varieties do not apply before tillering). Apply **Accurate** plus 2,4-D after tillering (refer to appropriate 2,4-D manufacturer's label), but before boot.

With Dicamba

For best results, apply **Accurate** at 1/10 oz. per acre: add 1/16 to 1/8 lb. active ingredient dicamba. Surfactant may be added to the mixture at ½ to 1 qt. per 100 gallons of spray solution; however, adding surfactant may increase the potential for crop injury. Also refer to dicamba labels for application timing and restrictions.

With 2,4-D (amine or ester) and Dicamba

Accurate may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D. Observe all applicable directions, restrictions and precautions on labels of all products used.

Make applications at 1/10 oz. of **Accurate** + 1/16-1/12 lb. active ingredient dicamba + 4-6 oz. active 2,4-D Ester or Amine per acre. Use higher rates when weed infestation is heavy. Add 1-2 pts. of surfactant to the 3 way mixture, where necessary, as deemed by local recommendations. Use of additional surfactant may not be needed with the higher phenoxy rates and ester phenoxy formulations. Consult the specific 2,4-D or dicamba label, or local recommendations for more information.

Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum wheat) apply after the crop is tillering and before it exceeds the 5-leaf stage.

Do not apply this 3-way mixture at high rates more than once a year or more than twice per year at the low rates.

With Bromoxynil (such as "Buctril", "Bronate")

Accurate may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, or fallow. For best results, add bromoxynil containing herbicides to the tank at 3 to 6 oz. active ingredient per acre (such as "Bronate" or "Buctril" at $\frac{3}{4}$ - $1\frac{1}{2}$ pts. per acre).

With "Starane"

For improved control of Kochia (2-4" tall), Russian thistle, mustard species and wild buckwheat, **Accurate** may be tank mixed with $\frac{1}{3}$ to $1\frac{1}{3}$ pts. per acre of "Starane."

With "Starane" + "Salvo"

For improved control of Kochia (2-4" tall) Russian thistle, mustard species and wild buckwheat, **Accurate** may be tank mixed with $\frac{2}{3}$ to $2\frac{2}{3}$ pts. per acre of "Starane" + "Sword."

With "Starane" + "Sword"

For improved control of Kochia (2-4" tall) Russian thistle, mustard species and wild buckwheat, **Accurate** may be tank mixed with $\frac{3}{4}$ to $2\frac{3}{4}$ pts. per acre of "Starane" + "Sword."

With "Maverick"

Accurate can be tank mixed with "Maverick" herbicide for improved control of weeds in wheat and barley.

With "Stinger", "Curtail", or "Curtail M" or "Widematch"

Accurate can be tank mixed with "Stinger", "Curtail", or "Curtail M" herbicides for improved control of weeds in wheat and barley.

With NUANCE™ Herbicide

Accurate may be tank mixed with NUANCE Herbicide based on local recommendations.

With NIMBLE™ Herbicide

Accurate may be tank mixed with NIMBLE Herbicide based on local recommendations.

With Grass Control Products

Tank mixtures of **Accurate** and grass control products may result in poor grass control. Cheminova recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or Cheminova representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of **Accurate** and the grass product to a small area. Do not tank mix **Accurate** with "Hoelon" 3EC, as grass control may be reduced.

With "Assert" herbicide or "Avenge" herbicide

Accurate can be tank mixed with "Avenge" or "Assert". When tank mixing **Accurate** with "Assert", always include another broadleaf weed herbicide with a different mode of action (for example: 2,4-D ester, MCPA ester, "Buctril", or "Bronate"). Tank mixed applications of NUANCE™ plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

With "Puma"

Accurate can be tank mixed with "Puma" herbicide for improved control of weeds in wheat and barley.

With "Discover NG"

Accurate can be tank mixed with "Discover NG" herbicide for improved control of weeds in spring wheat.

With "Everest"

Accurate can be tank mixed with "Everest" herbicide for improved control of weeds in spring wheat.

With Insecticides and Fungicides

Accurate may be tank mixed or used sequentially with insecticides and fungicides registered for use on cereal grains.

However, under certain conditions (drought stress, cold weather, or if the crop is in the 2 – 4 leaf stage), tank mixes or sequential applications of **Accurate** with organophosphate insecticides (such as parathion, "Di-Syston") may product temporary crop yellowing or, in severe cases, crop injury.

The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application.

Test these mixtures in a small area before treating large areas.

Do not apply **Accurate** within 60 days of crop emergence where an organophosphate insecticide (such as "Di-Syston") has been applied as an in-furrow treatment as crop injury may result.

Do not use **Accurate** plus malathion, as crop injury will result.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing **Accurate** in fertilizer solution.

Accurate must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the **Accurate** is added. Use of this mixture may result in temporary crop yellowing or stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at ½ pt – 1 qt per 100 gal of spray solution (0.06 – 0.25% v/v) based on local recommendations.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or Cheminova representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with **Accurate** and fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Do not add surfactant when using **Accurate** in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

**ACCURATE HERBICIDE WITH MCPA, 2,4-D AND/OR
DICAMBA FOR SUPPRESSION OF WINTER ANNUAL BROADLEAF
WEEDS IN WINTER WHEAT TO BE GRAZED OUT IN THE STATE OF
TEXAS, OKLAHOMA, NEW MEXICO, and KANSAS**

GENERAL INFORMATION

Accurate herbicide can be tank mixed with MCPA, 2,4-D and/or dicamba for suppression of winter annual broadleaf weeds in winter wheat to be grazed out and not harvested for grain, in the states of Texas, Oklahoma, New Mexico and Kansas.

DIRECTIONS FOR USE

For the suppression of winter annual broadleaf weeds (such as henbit and mustards) in winter wheat in the states of Texas, Oklahoma, New Mexico and Kansas, **Accurate** at 0.05 (1/20) oz. per acre should be tank mixed with MCPA, 2,4-D and/or dicamba at label rates. Winter annual broadleaf weeds should be less than 1" tall or in the rosette stage for suppression. Add a Cheminova recommended nonionic surfactant having at least 80% active ingredient at 1 to 2 qts. per 100 gallons of spray solution (0.25 to 0.5% v/v).

Rotation Intervals For Crops in Non-Irrigated Land Following Use of **ACCURATE** at 0.05 (1/20) oz. per acre on Wheat That Will be Grazed Out

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Sorghum, Grain	7.9 or lower	No restrictions	4
Cotton	7.9 or lower	No restrictions	10
Alfalfa	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Beans, Dry	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22

Rotation intervals for crops not covered above following the use of **Accurate** at 0.05 (1/2) oz. per acre on wheat that will be grazed out.

The minimum rotation interval is 22 months with at least 18" of cumulative precipitation during the period:

- to any crop not listed on the rotation intervals table above
- if the soil pH is not in the specified range

To rotate to a crop at an interval shorter than recommended, a field bioassay must be successfully completed to rotate to that crop. See section on Field Bioassay for further information.

IMPORTANT RESTRICTIONS

This treatment is for use on winter wheat that will be grazed out and will not be harvested for grain

IMPORTANT PRECAUTIONS

Accurate suppresses weeds by postemergence activity. For best results, apply **Accurate** to young, actively growing weeds. The degree and duration of suppression at 1/20 oz. per acre may depend upon the following factors:

- weed spectrum and infestation intensity
- weed size at application
- environmental condition at and following treatment

WHEAT, BARLEY AND TRITICALE – HARVEST AID

Use Rates

Apply 1/10 oz. of **Accurate** per acre in combination with 2,4-D or Glyphos to aid in dry down of many broadleaf weeds, thereby aiding grain harvest.

Application Timing

Make applications after the crop has reached the hard dough stage, but no later than 10 days before harvest.

TANK MIXTURES IN HARVEST AID

A tank mix of **Accurate** plus 2,4-D and surfactant, or Glyphos, will typically aid in dry down of many broadleaved weeds, thereby aiding grain harvest. Postemergence application should be made to actively

growing weeds after the crop is in the hard dough stage. If weeds are not dry within 10 days after application, delay harvest until weeds are dry.

See weeds listed in Weeds Controlled chart of this label.

With 2,4-D

Use 1/10 oz **Accurate** plus ¼ to ½ lb active ingredient 2,4-D per acre on moderate weed infestations; higher rates of 2,4-D may be used on large weeds if permitted by the 2,4-D brand labeling. Include 1 to 2 qt surfactant per 100 gal spray solution.

In addition to the weeds listed in Weeds Controlled chart of this label, the 2,4-D combination will also dry down common cocklebur, marehail, puncturevine and common and wild sunflower. In areas where 2,4-D use is restricted, apply **Accurate** with surfactant only; however, this treatment may be less effective.

With Glyphos

Use 1/10 oz **Accurate** plus the locally recommended rate of Glyphos (see Glyphos label for maximum seasonal rate). **Accurate** requires the use of an adjuvant for optimum activity – Consult the Glyphos label or local recommendations for the amount of adjuvant to include.

GRAIN SORGHUM

General Information

Accurate is recommended for use on irrigated or dryland grain sorghum in Colorado, Kansas, Nebraska, Oklahoma and Texas (North of I-20).

Use Rates: Apply **Accurate** at 1/20 oz. per acre plus ¼ lb. active ingredient 2,4-D amine per acre. Do not use surfactant or crop oil.

Crop Stage: For optimum performance and crop safety, apply **Accurate** plus 2,4-D amine when grain sorghum is 3 to 15 inches in height. If sorghum is taller than 10 inches to the top of the canopy, use drop nozzles and keep spray off the foliage. Apply only before the boot stage. Read and follow all other use instructions, warnings and precautions on companion herbicide labels.

Sorghum varieties vary in sensitivity to 2,4-D amine. Spray only varieties known to be tolerant to 2,4-D amine. Contact seed company and Local County Extension Service for this information.

Pest Stage: Application of **Accurate** plus 2,4-D amine should be made when all or a majority of the weeds have germinated and emerged. For best results, spray when weeds are less than 6 inches tall.

Weeds Controlled with Tank Mix of **Accurate plus 2,4-D amine:**

- Pigweed species
- Puncture vine
- Velvetleaf

Application Information

Accurate may be applied to grain sorghum by properly calibrated ground or aerial equipment. **Accurate** can be used on either dryland or irrigated grain sorghum. If application is made to irrigated sorghum, delay first post-treatment irrigation for at least 3 days after treatment. The first post-treatment irrigation should not exceed 1".

Use cultivation prior to **Accurate** + 2,4-D amine treatment to cover exposed brace roots of grain sorghum to minimize injury from 2,4-D amine.

Precautionary Statements

- Temporary crop yellowing and/or stunting may occur soon after application, especially when crop is under stress conditions.

- Do not use on grain sorghum grown for seed production or syrup. Do not use on forage sorghum.
- Do not use for forage or silage within 30 days of application.
- Do not include a surfactant or crop oil to the tank mix.
- Do not apply this treatment under cold, wet weather conditions or to grain sorghum growing under stress caused by weather, insects or disease as crop injury may result.
- Do not apply to long season grain sorghum varieties or grain sorghum that is planted after July 1, as crop injury or delayed maturity may occur.
- Do not exceed (1) one application per year.
- **Accurate** must be used with 2,4-D; in areas where 2,4-D use is restricted, follow requirement of the restriction. If 2,4-D use is prohibited, do not use **Accurate** on grain sorghum.

SURFACTANTS

Spray Adjuvants

Applications of **Accurate** must include either a nonionic surfactant or a crop oil concentrate. In addition, an ammonium nitrogen fertilizer may be used. If another herbicide is tank mixed with **Accurate** select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

Antifoaming agents may be needed. Consult your Ag dealer or applicator for a listing of recommended surfactants.

Nonionic Surfactants (NIS)

- Apply 0.06 to 0.50% v/v (1/2 to 4 pts. per 100 gallons of spray solution) – see Tank Mixtures section for additional information.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Exceptions: On all spring wheat and spring or winter barely use ½ to 1qt. per 100 gallons.

Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Ammonium Nitrogen Fertilizer

- Use 2 qts./acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lbs./acre of spray grade ammonium sulfate (AMS). Use 4 qts./acre UAN or 4 lbs./acre AMS under arid conditions.
- Do not use liquid nitrogen fertilizer as the total carrier solution.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality.

Antifoaming agents may be used if needed.

Do not use low rates of liquid fertilizer as a substitute for surfactant.

GROUND APPLICATION

To obtain optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flood nozzles on 30" spacings, use at least 10 gallons per acre (GPA), flood nozzles no larger than TK10 (or equivalent), and a pressure of at least 30 pounds per square inch (psi). For 40" nozzle spacings,

use at least 13 GPA; for 60" spacings, use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

With "Raindrop RA" nozzles, use at least 30 GPA and ensure that nozzle spray patterns overlap 100%.

For flat-fan nozzles, use at least 3 GPA for applications to wheat or barley. Use at least 10 GPA for applications to pasture or rangeland.

Use 50-mesh screens or larger.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

Wheat, Barley and Fallow – use 1 to 5 GPA. Use at least 3 GPA in Idaho, Oregon or Utah.
Pasture and Rangeland – Use 2 to 5 GPA.

When applying **Accurate** by air in areas adjacent to sensitive crops, use solid stream nozzles oriented straight back. Adjust the swath to avoid spray drift damage to sensitive crops downwind and/or use ground equipment to treat the border edge of fields. See the **Spray Drift Management** section of this label.

Product Measurement

Accurate is measured using the **Accurate** volumetric measuring cup. The degree of accuracy of this cup varies by +/- 7.5%. For more precise measurements, use scales calibrated in ounces.

WITH LIQUID NITROGEN FERTILIZER

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing **Accurate** in fertilizer solution.

Accurate must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the **Accurate** is added. Use of this mixture may result in temporary crop yellowing or stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at ½ pt – 1 qt per 100 gal of spray solution (0.06 – 0.25% v/v) based on local recommendations.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or Cheminova representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with **Accurate** and fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Do not add surfactant when using **Accurate** in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Note: In certain areas east of the Mississippi river unacceptable crop response may occur with the use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist. In these areas consult your agricultural dealer, consultant or field advisor for a specific recommendation before using nitrogen fertilizer carriers.

Liquid nitrogen fertilizer solutions that contain sulfur can increase crop response.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

CROP ROTATION

Before using **Accurate**, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your wheat, barley, fallow, pasture, or rangeland acres at the same time.

Minimum Rotational Intervals

Minimum rotational intervals* are determined by the rate of breakdown of **Accurate** applied. **Accurate** breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase **Accurate** breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow **Accurate** breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

Accurate should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, **Accurate** could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of **Accurate**.

Checking Soil pH

Before using **Accurate**, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop not listed (see Rotation Intervals table), or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table, or if the minimum cumulative precipitation has not occurred since application.

Field Bioassay

To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with **Accurate**. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips.

If a field bioassay is planned, check with your local Agricultural dealer or Cheminova representative for information detailing the field bioassay procedure.

ROTATIONAL INTERVALS FOR CEREALS

All Areas – Following Use of **Accurate** at 1/10 oz per Acre

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Winter and spring wheat	7.9 or lower	No restrictions	1
Durum wheat, barley, spring/winter oat	7.9 or lower	No restrictions	10

19/33

ROTATION INTERVALS FOR CROPS IN NON-IRRIGATED LAND

Following Use of Accurate at 1/10 oz per Acre on Wheat, Barley, Fallow or Pasture

Location		Crop	Soil pH	Minimum Cumulative Precipitation	Minimum Rotation Interval
State	County or Area			(inches)	(months)
Colorado	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
		Field corn	7.9 or lower	15	12
		IR Corn	7.9 or lower	No restrictions	4
		STS Soybeans	7.9 or lower	No restrictions	4
Idaho	Southern Idaho	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas	7.3 or lower	10	10
		Chickpeas	7.4 or higher	28	34
Kansas	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Central and Western Kansas (West of the Flint Hills)	Field corn	7.9 or lower	15	12
		IR Corn	7.9 or lower	15	4
	Western Kansas W. of Hwy. 183	Soybeans	7.5 or lower 7.6-7.9	22 33	22 34
	Central Kansas; Generally E. of Hwy. 183 and W. of the Flint Hills	Soybeans	7.9 or lower	15	12
		STS Soybeans	7.9 or lower	15	4
Montana	Statewide	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22
		Alfalfa (hay only)	7.6-7.9	No restrictions	34
			7.5 or lower	No restrictions	22
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22

Location				Minimum Cumulative Precipitation	Minimum Rotation Interval
State	County or Area	Crop	Soil pH	(inches)	(months)
Nebraska	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
		IR Corn STS Soybeans	7.9 or lower	No restrictions	4
	Generally W. of Hwy. 77 and E. of the Panhandle	Field corn	7.9 or lower	15	12
		Soybeans	7.5 or lower	22	22
			7.6-7.9	33	34
New Mexico	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Eastern New Mexico	Cotton (dryland only)	7.9 or lower	30	22
North Dakota	W. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower, Sunflower	7.9 or lower	22	22
	E. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower, Sunflower	7.9 or lower	34	34
Oklahoma	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
		Field corn	7.9 or lower	15	12
		IR Corn STS Soybean	7.9 or lower	No restrictions	4
	Panhandle	Cotton (dryland only)	7.9 or lower	30	22
	E. of the Panhandle	Cotton (dryland only)	7.9 or lower	25	14

21/33

Location				Minimum Cumulative Precipitation	Minimum Rotation Interval
State	County or Area	Crop	Soil pH	(inches)	(months)
Oregon	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas	7.3 or lower	10	10
		Chickpeas	7.4 or higher	28	34
South Dakota	Statewide	Flax, Safflower, Soybean Sunflower	7.9 or lower	No restrictions	22
	S. of Hwy. 212 & E. of the Missouri River, & S. of Hwy. 34 & W. of the Missouri River	Grain sorghum, Proso millet	7.9 or lower	13	12
	Generally E. of Missouri River & S. of Hwy. 14, & W. of Missouri River	Field corn	7.9 or lower	15	12
Texas	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Soybean Sunflower	7.9 or lower	No restrictions	22
	Panhandle	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	30	22
	N. of Central Texas*	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	25	14
	* The counties of N. Central Texas are: Archer, Baylor, Bell, Bosque, Bowie, Callahan, Camp, Cass, Clay, Collin, Cooke, Coryell, Dallas, Delta, Denton, Eastland, Ellis, Falls, Fannin, Foard, Franklin, Grayson, Hardeman, Haskell, Hill, Hood, Hopkins, Hunt, Jack, Johnson, Kaufman, Knox, Lamar, Limestone, McLennan, Milam, Montague, Morris, Nafarro, Palo Pinto, Parker, Rains, Red River, Robertson, Rockwall, Shackelford, Somervell, Stephens, Tarrant, Throckmorton, Titus, Upshur, Van Zandt, Wilbarger, Wichita, Williamson, Wise, Wood, Young.				
Washington	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas	7.3 or lower	10	10
		Chickpeas	7.4 or higher	28	34

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Utah	Statewide	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
Wyoming	Statewide	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Southern Wyoming	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
	Southern Wyoming (Goshen, Laramie, and Platte counties only)	Field corn	7.9 or lower	15	12
	Northern Wyoming	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22

Rotation Intervals not covered above – The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (see the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than recommended, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

ROTATION INTERVALS IN PASTURE OR RANGELAND FOR OVERSEEDING AND RENOVATION

Location	Crop	Maximum Rate on Pasture (oz. per acre)	Minimum Rotation Interval (months)
AL, AR, FL, GAL, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV	alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, orchardgrass, bromegrass, ryegrass, fescue, timothy	1/10 to 3/10	4
	wheat (except Durum)	1/10 to 3/10	1
	Durum, barley, oat	1/10 to 3/10	10
All areas not included above*	red clover, white clover, sweet clover	1/10 to 2/10	12
	Bermudagrass, bluegrass, orchardgrass, bromegrass, ryegrass, timothy	1/10 to 2/10	6
	Fescue	1/10 to 2/10	18
	wheat (except Durum)	1/10 to 2/10	1
	Durum, barley, oat	1/10 to 2/10	10

Rotation Intervals not covered above – The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (see the Rotation Intervals table)
- if the use rate applied is not specified in the table

To rotate to a major field crop at an interval shorter than recommended, a field bioassay must be successfully

completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

RECROPPING INTERVALS FOR GRASSES ON CONSERVATION RESERVE PROGRAM (CRP)

Whenever **Accurate** has previously been used in wheat, barley, triticale or fallow, the following grasses may be planted after the intervals specified in the tables below. The planting of grass and legume mixtures is not recommended as injury to the legume may occur.

- Bentgrasses
- Blue grama
- Bluestems – Big, Little, Plains, Sand, WW Spar
- Buffalograss
- Galleta
- Green needlegrass
- Green sprangletop
- Indian ricegrass
- Lovegrasses – Sand, Weeping
- Orchardgrassnot (excluding Piaute)
- Prairie sandreed
- Sand dropseed
- Sheep fescue
- Sideoats grama
- Switchgrass
- Wild-ryegrasses – Beardless, Russian
- Wheatgrasses – Crested, Intermediate, Pubescent, Slender, Streamback, Tall, Thickspike, Western

ROTATIONAL INTERVALS

MN, MT, ND, SD, and Northern WY:

Soil pH	Use Rate (ounces/acre)	Minimum Interval for Planting Grasses
7.5 or lower	1/10	4 months (all grasses)
7.6 to 7.9	1/10	4 months (Wheatgrasses only)

AR, CO, ID, KS, LA, NE, NM, OK, OR, TX, UT, WA, Southern WY:

Soil pH	Use Rate (ounces/acre)	Minimum Interval for Planting Grasses
7.9 or lower	1/10	2 months (all grasses)

GRAZING

There are no grazing restrictions on **Accurate**.

IMPORTANT PRECAUTIONS

Treated vegetation may be cut for forage or hay. Coveralls, shoes plus socks and chemical resistant gloves made of any waterproof material must be worn if cutting within 4 hours of treatment.

MIXING INSTRUCTIONS

1. Fill the tank $\frac{1}{4}$ to $\frac{1}{3}$ full of water (If using liquid nitrogen fertilizer solution in place of water, see Tank Mixtures sections for additional details).
2. While agitating, add the required amount of **Accurate**.
3. Continue agitation until the **Accurate** is fully dispersed, at least 5 minutes.
4. Once the **Accurate** is fully dispersed, maintain agitation and continue filling tank with water. **Accurate** should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.

7. Apply **Accurate** spray mixture within 24 hours of mixing to avoid product degradation.
8. If **Accurate** and a tank mix partner are to be applied in multiple loads, pre-slurry the **Accurate** in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the **Accurate**.

Do not use **Accurate** with spray additives that reduce the pH of the spray solution to below 3.0.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when the crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping to avoid crop injury.

Do not make applications using equipment and/or spray volumes or under weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift, refer to the Spray Drift Management section of the label.

Continuous agitation is required to keep **Accurate** in suspension.

SPRAYER CLEANUP

Spray equipment must be cleaned before **Accurate** is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined in the **After Spraying Accurate** section of this label.

At the End of the Day

When multiple loads of **Accurate** are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

After Spraying Accurate and Before Spraying Crops Other Than Wheat, Barley, Fallow, Pasture, or Rangeland

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of **Accurate** as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or a Cheminova-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your agricultural dealer, applicator, or Cheminova representative for a listing of approved cleaners.

NOTES:

1. **Attention:** Do not use chlorine bleach with ammonia, as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When **Accurate** is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of **Accurate** and applications of other pesticides to **Accurate**-sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to **Accurate** to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**IMPORTANCE OF DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets (>150 – 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label.

Controlling Droplet Size – General Techniques

- **Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size – Aircraft

- **Number of Nozzles** – Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** – Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** – The boom length should not exceed $\frac{3}{4}$ of the wing or rotor length – longer booms increase drift potential.
- **Application Height** – Application more than 10 ft. above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

NOTE: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

To better manage weed resistance when using **Accurate**, use a combination of tillage, and tank-mix partners or sequential herbicide applications that have a different mode of action than **Accurate**, to control escaped weeds. Do not let weed escapes go to seed.

Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative herbicide recommendations available in your area.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes.

PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain, or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in location where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
 - Do not use on grasses grown for seed.
- Do not apply to irrigated land where tailwater will be used to irrigate crops other than wheat and barley.
- Do not apply to frozen ground as surface runoff may occur.
- Do not apply to snow-covered ground.
- Wheat and barley varieties may differ in their response to various herbicides. Cheminova recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of **Accurate** to a small area.
- Under certain conditions, such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after **Accurate** application, temporary discoloration and/or crop injury may occur. **Accurate** should not be applied to wheat or barley that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5-leaf stage. Severe winter stress, drought, disease or insect damage following application also may result in crop injury.
- The combined treatment effects of **Accurate** postemergence preceded by preemergence wild oat herbicides may cause crop injury to spring wheat when crop stress (soil crusting, planting too deep, prolonged cold weather, or drought) causes poor seedling vigor.
- In the Pacific Northwest, to prevent cold weather-related crop injury, avoid making applications during winter months when weather conditions are unpredictable and can be severe.
- Do not apply to wheat, barley or pastures undersown with legumes, as injury to the forage may result.
- To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery dry or light sandy soils until they have been stabilized by rainfall, trashy mulch, reduced tillage, or other cultural practices. Injury to immediately adjacent crops may occur when treated soil is blown onto land used to produce crops other than cereal grains or pasture/rangeland.
- For ground applications applied to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced. The addition of 2,4-D or MCPA should improve weed control under these conditions.

WARRANTY DISCLAIMER

Cheminova warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CHEMINOVA MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Cheminova or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Cheminova's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Cheminova shall not be liable for losses or damages resulting from handling or use of this product unless Cheminova is promptly notified of such loss or damage in writing. In no case shall Cheminova be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Cheminova or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Accurate and Glyfos are registered trademarks of Cheminova A/S.

Raindrop RA is a trademark of Delavan Corp.

Assert is a registered trademark of Nufarm, Inc.

Avenge is a registered trademark of Amvac Chemical Corporation

Banvel is a registered trademark of Micro Flo Company, LLC

Buctril, Bronate, Hoelon, Puma and Di-Syston are trademarks of Bayer CropScience

Discover NG is a registered trademark of Syngenta Crop Protection

Sword, Salvo are trademarks or registered trademarks of Loveland Products, Inc.

Everest is a registered trademark of Arysta LifeScience North America

Starane, Stinger, Widematch and Curtail are registered trademarks of Dow AgroSciences, LLC

Maverick is a registered trademark of Monsanto Technology, LLC

SUPPLEMENTAL LABELING

Accurate® Herbicide

EPA Reg. No. 67760-68

FOR WEED CONTROL IN SUGARCANE IN THE STATE OF HAWAII

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state, consult the agency responsible for pesticide regulation.

GENERAL INFORMATION

Accurate® is recommended for use on sugarcane in Hawaii.

INFORMATION ON RESISTANT WEEDS

Naturally-occurring weed biotypes resistant to this product are known to exist in Hawaii sugarcane. To delay the development of resistant biotypes, refer to the Weed Resistance section on the Federal Section 3 label.

SUGARCANE MAXIMUM USE RATE: Do not apply more than 6.25 ounces of product per acre per crop.

WEED CONTROL IN SUGARCANE: For best weed control or weed suppression, apply **Accurate®** as a preplant/preemergence broadcast application or as postemergence direct spray application to weeds when environmental conditions favor active growth of broadleaf weeds. Unless otherwise directed, always include a surfactant at 0.25 to 0.50% V/V for postemergence mixture.

PREPLANT/PREEMERGENCE APPLICATION: Apply as a broadcast application at a rate of 0.83 to 3.33 ounces per acre with ground equipment or air. Application should be made before crop emerges as crop injury to some varieties may occur.

POSTEMERGENCE APPLICATION: Apply as directed spray application to emerged weeds at rates of 0.83 to 3.33 ounces per acre with ground equipment or by hand equipment. Avoid contact with the sugarcane foliage as injury may occur.

WEEDS CONTROLLED: **Accurate®** will control a number of broadleaf weeds including:

Amaranthus sp.
Ipomoea triloba (Aiea Morningglory)
Portulaca sp. (Purslane)
Euphorbia sp. (Spurge)
Momordica charantia (Peria)
Erigeron canadensis (Fireweed)
Crotalaria sp.
Ageratum sp.
Crassocephalum crepidioides (Thick head)

PARTIALLY CONTROLLED:

Chloris inflata (Swollen fingergrass)

NOTE: Do not apply **Accurate®** closer than 16 months before harvest.

PRECAUTIONS:

- Do not use liquid fertilizer in addition or as a substitute for a surfactant.
- Do not use on soils with pH greater than 7.9 (for example, highly calcareous soils) if the following crop is sensitive to **Accurate®**. Extended soil residual activity could adversely affect minimum rotation intervals for all crops.
- Do not apply to irrigated land where tailwater will be used to irrigate crops other than sugarcane. Varieties of sugarcane may differ in their tolerance to herbicides and some varieties can be adversely affected with post emergence applications. Therefore it is recommended that a small test block be installed prior to large scale applications.
- Do not apply **Accurate®** to sugarcane that is stressed by severe weather conditions, drought, low fertility, water saturated soil, disease or insect damage, as crop injury may result. Under certain conditions such as heavy rainfall, temporary discoloration and/or crop injury may occur. To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery dry or light sandy soils until they have been stabilized by rainfall, trashy mulch, reduced tillage or other cultural practices. Injury to adjacent crops may occur when treated soil is blown onto land used to produce other crops. For ground applications applied to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced.

Accurate® must be cleaned from application equipment according to cleanup procedures described in the SPRAYER CLEANUP section of the Federal Section 3 label.

IMPORTANT NOTICE

BEFORE BUYING OR USING Accurate®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This label contains supplemental instructions for use of the referenced Cheminova product. Follow these instructions carefully. This label must be in the possession of the user at the time of application.

Read the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies on the Section 3 Federal product label.

Accurate® is a registered trademark of Cheminova A/S

Cheminova, Inc.
One Park Drive, Suite 150
P.O. Box 110566
Research Triangle Park, NC 27709

SUPPLEMENTAL LABELING

Accurate® Herbicide

EPA Reg. No. 67760-68

FOR SELECTIVE WEEDING TO AID IN THE ESTABLISHMENT AND MAINTENANCE OF GRASSES IN THE CONSERVATION RESERVE PROGRAM

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Accurate® is recommended for the control or suppression of certain broadleaf weeds (refer to EPA approved label for weeds controlled and suppressed) in the following perennial native or improved grasses grown on land enrolled in the Conservation Reserve Program (CRP):

Blue Grama	Orchardgrass
Bluestems -	Sideoats grama
big	Switchgrass - blackwell
little	Wheatgrasses -
plains	bluebunch
sand	crested
WW Spar	intermediate
Buffalograss	pubescent
Green sprangletop	Siberian
Indiangrass	slender
Kleingrass	streambank
Lovegrasses -	tall
atherstone	thickspike
sand	western
weeping	Wildrye grass - Russian
wilman	

GENERAL INFORMATION

Accurate® provides postemergence activity and offers short-term residual control **or suppression** of labeled weeds. Degree of control and duration of effect depend on weed spectrum and density; weed size and variability; amount of cover canopy; growing conditions prior to, at and following application; amount and timing of precipitation; and spray coverage.

Maximize potential for grass establishment by consulting with the Soil Conservation Service or other local experts concerning planting techniques and other cultural practices.

APPLICATION TIMING

Preplant (prior to planting)

Accurate® may be applied prior to planting at not more than 1/10 oz/A on all labeled grasses except orchardgrass and Russian wildrye grass.

Preemergence (after planting but before grass emergence)

Accurate® may be applied after planting, but before grass emergence, at not more than 1/10 oz/A on all labeled grasses except orchardgrass and Russian wildrye grass.

Early postemergence to new plantings

Accurate® may be applied at not more than 1/10 oz/A on all labeled grasses anytime after grass emergence. Because grass species differ in time of emergence, apply only after majority of grasses are in the 3-to 4-leaf stage.

Early postemergence applications to stands planted the previous season

Accurate® may be applied at not more than 1/10 oz/A on all labeled grasses if the majority of the grasses have one or more leaves.

Late postemergence (tillered stands) to stands planted the previous season

Accurate® may be applied at not more than 1/10 oz/A to all labeled grasses in all labeled states.

WEED CONTROL

For best weed control performance, use **Accurate®** in a tank mix with 2,4-D (ester formulations perform best). Read and follow all manufacturer's label recommendations for the 2,4-D used. If those recommendations conflict with this label, do not tank mix the 2,4-D with **Accurate®**.

Apply **Accurate®** preemergence to weeds or postemergence to small actively growing weeds before they exceed 4" in height or diameter. For all postemergence applications, add a surfactant at the rate of 1 to 2 quarts/100 gallons of spray solution.

One to two inches of rainfall (enough to wet the top 2-3 inches of soil profile) may be needed to move the herbicide into the weed root zone before the next flush of weeds emerge. The amount of moisture required for sufficient activation increases with crop or weed residue and for finer textured soils. Without sufficient rainfall to move **Accurate®** into the weed root zone, weeds that germinate after treatment will not be controlled.

Avoid postemergence applications to weeds which are not actively growing due to adverse weather conditions. Weeds hardened off by cold weather or drought stress may not be adequately controlled.

Performance: Because newly planted CRP grass stands do not sufficiently compete with weeds and because weed pressure in CRP fields is often severe, performance from **Accurate®** may not always be satisfactory. An additional herbicide application or mowing may be needed.

TANK MIXTURES**Preplant**

Accurate® may be tank mixed with Glyphos® or "Landmaster" II as a preplanting treatment to control broadleaf and grassy weeds. When using "Landmaster" II tank mix, allow at least 7 days after application before planting grasses. Refer to Glyphos® and "Landmaster" II labels for all use instructions, label rates, weed control, claims, warnings and precautions.

Postemergence

Accurate® can be tank mixed with 2,4-D at ¼ lb AI/A for all labeled grasses larger than the 5-leaf stage. For fully tillered stands up to ½ lb AI/A of 2,4-D may be used. Surfactant may be added at ½ to 1 qt per 100 gallons of spray solution. However, the addition of surfactant may increase the chance of grass injury.

Accurate® can also be tank mixed with "Banvel". Use not more than 1/8 to ¼ lb AI/A "Banvel" plus **Accurate®** after majority of grasses are in the 3-leaf stage. In established grasses (2nd year stands), use not more than ¼ to ½ lb AI/A "Banvel" plus **Accurate®**. Surfactant may be added at ½ to 1 qt per 100 gallons of spray solution. However, the addition of surfactant may increase the chance of grass injury.

Read and follow all use instructions, label rates, weed control claims, warnings and precautions for the companion herbicide(s).

PRECAUTIONS

Under certain conditions such as high soil pH, heavy rainfall, prolonged cool weather, or frost conditions just prior to or soon after application, temporary discoloration or stunting of the grasses may occur.

Legumes in a seeding mixture may be severely injured or killed following an application of Accurate®. Do not apply to frozen ground where surface runoff may occur. Do not apply when snow covers the ground. Do not use more than 2/10 ounce per acre per year.

Because cultivars of perennial grass differ in their tolerance to herbicides, limit the first use of **Accurate®** to a small area prior to adoption as a field practice. Likewise, components in a seed mixture will vary in tolerance to **Accurate®** so that the final stand may not reflect seed ratio.

Do not treat stands showing winter stress or lack of vigor symptoms as grass injury may occur.

IMPORTANT NOTICE

BEFORE BUYING OR USING Accurate®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This label contains supplemental instructions for use of the referenced Cheminova product. Follow these instructions carefully. This label must be in the possession of the user at the time of application.

Read the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies on the Section 3 Federal product label.

Accurate® is a registered trademark of Cheminova A/S
"Banvel" is a registered trademark of BASF Corporation
"Landmaster" is a registered trademark of Albaugh, Inc.

Cheminova, Inc.
One Park Drive, Suite 150
P.O. Box 110566
Research Triangle Park, NC 27709