

67760-116

03/07/2012

1/36



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Chemical Safety and Pollution Prevention
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

67760-116

Date of Issuance:

MAR 07 2012

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Couraze 4 Insecticide

Name and Address of Registrant (include ZIP Code):

Cheminova, Inc.
Washington Office
1600 Wilson Boulevard
Suite 700
Arlington, VA 22209

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No 67760-116."
 - On page 6, under "Pollinator Information:" add the subheading "RESTRICTION:" in front of the sentence "Do not use Couraze 4 Insecticide on crops requiring bee pollination during bloom and a minimum of 10 days prior to bloom."

Signature of Approving Official:

Venus Eagle, Product Manager 01
Insecticide-Rodenticide Branch, Registration Division (7505P)

Date:

MAR 07 2012

- In every crop box that refers to "unless otherwise directed/allowed by supplemental labeling" revise to read "unless otherwise directed by state specific supplemental labeling."

3. Storage stability (830.6317) and corrosion characteristics (830.6320) data must be submitted within 18 months from the date of this registration notice.

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Please note that, should you wish to add/retain a reference to the company's website on your label, the website then becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions, please contact Autumn Metzger at 703-305-5314 or metzger.autumn@epa.gov.

A stamped copy of the label is enclosed for your records.

Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

coGROUP **4A** INSECTICIDE

COURAZE 4 INSECTICIDE

A systemic, soil and foliar applied insecticide for use on listed crops including cotton, peanuts, potatoes, soybeans and tobacco; in citrus, tree nut and fruit orchards; on field and listed greenhouse vegetables; on berry, bush and vine crops; and on other listed crops. Also for use in commercial poultry facilities.

ACTIVE INGREDIENT	% BY WT.
Imidacloprid; 1-[6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	40.4%
OTHER INGREDIENTS:	59.6%
TOTAL:	100.0%

Contains 4 lbs. of active ingredient per gallon

EPA Reg. No. 67760-

EPA Est. No.

Manufactured for:
Cheminova, Inc.
P. O. Box 110566
One Park Drive
Research Triangle Park, NC 27709
www.cheminova.us.com

ACCEPTED
With COMMENTS
In EPA Letter Dated:
MAR 07 2012
Under the Federal Insecticide, Fungicide
and Rodenticide Act, As amended, for the
pesticide Registered under EPA Reg. No:
67760-116

NET CONTENTS: 1 GALLON

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
IF INHALED:	<ul style="list-style-type: none"> • Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice:
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Cheminova, Inc. 1-866-303-6950 for emergency medical treatment information.	

4/36

NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid contact with eyes or clothing. Wear protective eye wear. Wear long sleeved shirt and long pants, socks, shoes, and chemical resistant gloves made of waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and' long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC), or Viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4.6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gels inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

TAKE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

For Aerial Applications

For aerial applications, mount the spray boom on the aircraft so to minimize drift caused by wing tip vortices. Use the minimum practical boom length. Do not exceed 75% of the wing span or rotor diameter. Release the spray at the lowest possible height consistent with good pest control and flight safety. Do not use applications more than 10 feet above the crop canopy.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the airstream as much as possible and by avoiding excessive spray boom pressure.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy, and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions

Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Mixing and loading Requirements

To avoid potential contamination of groundwater, use a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading area and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes, or field drains.

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. Follow specific drift management practices:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows).
- Only spray inward, toward the orchard or vineyard, for application to the outside rows.

No-Spray Zone Requirements for Soil and Foliar Applications

Do not apply by ground, within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Runoff Management

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When using Couraze 4 Insecticide on erodible soils, employ the best management practice for minimizing runoff. Consult your local Natural Resources Conservation Service for recommendations in your use area.

Endangered Species Notice

Under the Endangered Species Act, it is a federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

Resistance Management

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

Couraze 4 Insecticide contains a Group 4A insecticide. Insect biotypes with acquired or inherent tolerance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by Couraze 4 Insecticide and to other Group 4A insecticides.

The active ingredient in Couraze 4 Insecticide is a member of the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to Couraze 4 Insecticide. In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season: 1) only a single, soil application of Couraze 4 Insecticide be made; 2) foliar applications of products from the same class not be made following a long residual, soil application of Couraze 4 Insecticide, or other neonicotinoid products.

If a soil application of Couraze 4 Insecticide has not been made during a crop season and foliar applications are to be made, avoid using a block of more than three consecutive applications of Couraze 4 Insecticide and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Cheminova strongly encourages the rotation to a block of applications with effective products with a different mode of action before using additional applications of neonicotinoid products. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect's ability to develop resistance to this class of chemistry.

Foliar applications of this product or other products of Group 4A insecticides must not be used on crops previously treated with a long-residual, soil-applied product from this same Group.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Clutch, Couraze, Galian, Impulse, Intruder, Leverage, Nuprid; Pasada, Provado, Trimax Pro, and Venom.

Other Group 4A, neonicotinoid products used as soil/seed treatments include: Admire Pro, Advise, Alias, Belay, Couraze, Cruiser, Gaucho, Macho, Macho Max, Nuprid, Platinum, Venom, and Widow.

Contact your Cooperative Extension specialist, certified crop advisor, and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil injected or soil incorporated, the worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC), or Viton
- Shoes plus socks

APPLICATION DIRECTIONS

Soil Applications:

For soil applications of Couraze 4 Insecticide, direct product into the seed or root zone of crop. Failure to place Couraze 4 Insecticide into root zone may result in loss of control or delay in onset of activity. Couraze 4 Insecticide may be applied with ground or chemigation application equipment.

When applied as a soil application, optimum activity of Couraze 4 Insecticide results from applications to the root-zone of plants to be protected. The earlier Couraze 4 Insecticide is available to a developing plant, the earlier the protection begins. The systemic nature of Couraze 4 Insecticide allows movement from roots through the xylem tissue to all vegetative parts of the plant. This results in extended residual activity of Couraze 4 Insecticide, the control of insects, and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate of Couraze 4 Insecticide applied affects the length of the plant protection period. Higher listed rates within the specified rate range can be used when infestations occur later in crop development or where pest pressure is continuous. Couraze 4 Insecticide will generally not control insects infesting flowers, blooms, or fruit. Additional crop protection may be required for insects feeding in, or on these plant parts, and for insects not listed in the crop-specific, pests-controlled sections of this label. Additionally, specific application information for Couraze 4 Insecticide is also provided in the crop-specific sections of this label.

Foliar Applications:

Apply foliar applications of Couraze 4 Insecticide as a directed or broadcast foliar spray. Thorough coverage of foliage is necessary without runoff for optimum insecticidal efficacy. Use adequate spray volumes, properly calibrated application equipment, and spray adjuvant if necessary to obtain thorough coverage. Failure to provide adequate coverage and retention of Couraze 4 Insecticide on leaves and fruit may result in loss of insect control or delay in onset of activity. Couraze 4 Insecticide may be applied with properly calibrated ground or aerial application equipment. Minimum spray volumes unless otherwise specified on crop specific application sections are 10 gallons per acre by ground and 5 gallons per acre by air. Couraze 4 Insecticide may also be applied by overhead chemigation (see additional information in "Chemigation" section of this label below) if allowed in crop specific application section.

RESTRICTIONS:

Do not apply Couraze 4 Insecticide in enclosed structures such as planthouses or greenhouses except as specified in the TOBACCO, CUCURBIT VEGETABLES, FRUITING VEGETABLES and GREENHOUSE VEGETABLES, (Mature plants in production greenhouses): Cucumber, Tomato only sections of this label.

Make applications only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool, or other soil-less media. Couraze 4 Insecticide should be pre-mixed with water or other appropriate diluents prior to application. Keep Couraze 4 Insecticide and water suspension agitated to avoid settling.

Plants Grown for Seed:

Use of Couraze 4 Insecticide on crops grown for production of true seed intended for private or commercial planting is not permitted unless allowed under state specific, supplemental labeling.

Supplemental Control Measures:

Suppression or less than complete control of insect pests including reduced feeding may also result from application of Couraze 4 Insecticide. Control of these pests/diseases may require supplemental control measures.

Pollinator Information:

As with any insecticide, care must be taken to minimize exposure of Couraze 4 Insecticide to honey bees and other pollinators. Do not use Couraze 4 Insecticide on crops requiring bee pollination during bloom and a minimum of 10 days prior to bloom.

Additional Information:

Additional information on uses of Couraze 4 Insecticide for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants, or local Cheminova, Inc. representatives.

Application Rate Restriction:

Do not apply more than 0.5 lb. active ingredient per acre per year regardless of formulation or method of application.

MIXING INSTRUCTIONS

To prepare the application mixture, add a portion of the required amount of water to the spray tank and with agitation, add Couraze 4 Insecticide. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application. Couraze 4 Insecticide may also be used with other pesticides and/or fertilizer solutions. Please see "Compatibility" section of this label. When tank mixtures of Couraze 4 Insecticide and other pesticides are involved, prepare the tank mixture as specified above and follow suggested "Mixing Order" below.

Mixing Order

When pesticide mixtures are needed, add wettable powders or wettable granules first; Couraze 4 Insecticide and other suspension concentrate (flowable) products second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer/pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

Compatibility

Test compatibility of the intended mixture before adding Couraze 4 Insecticide to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Do not use if poor mixing or formation of precipitates that do not readily redisperse occur, as it indicates an incompatible mixture.

CHEMIGATION

Types of Irrigation Systems: Foliar chemigation applications of Couraze 4 Insecticide may be made to crops through overhead sprinkler systems if specified in crop specific application sections. Soil chemigation applications of Couraze 4 Insecticide may only be made to crops through chemigation as specified in crop specific application sections and only through low pressure systems specified for a given crop. Do not apply Couraze 4 Insecticide through any other type of irrigation system.

Make foliar chemigation applications of Couraze 4 Insecticide as concentrated as possible. Retention of Couraze 4 Insecticide on target site of insect infestation is necessary for optimum activity. Chemigation of Couraze 4 Insecticide in water volumes exceeding .10 inches per acre are not recommended. See crop specific application sections of the label for more information.

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact Cooperative Extension Service specialists, equipment manufacturers, or other experts.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional reduced pressure zone, back flow preventer (RPZ), or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump

motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROTATIONAL CROPS*

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient; as soon as practical, following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval is required.

IMMEDIATE PLANT-BACK:

All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop, and sweet), rapeseed, sorghum, and wheat.

30 DAY PLANT-BACK:

Cereals (including buckwheat, millet, oats, rice, rye, and triticale), and safflower.

12 MONTH PLANT-BACK: All Other Crops

* Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

FIELD CROPS

Cotton

Soil Application

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Cotton aphid, Plant bugs, Thrips, Whiteflies	0.65	8.5 – 10.6 (depending on row-spacing)
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 14 days • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 10.55 fluid ounces per acre (0.33 lb active ingredient per acre). • Regardless of formulation or method of application, apply no more than 0.5 lb. active ingredient per acre per season, including seed treatment, soil, and foliar uses. • Do not apply more than a total of 6 applications of the active ingredient per season. • Do not graze treated fields after any application of Couraze 4 Insecticide. Applications: Apply specified dosage in one of the following methods: In-furrow spray during planting directed on or below seed. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. Chemigation into root zone through low pressure drip or trickle irrigation. See Resistance Management Section of this label.		

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Cotton aphid, Cotton flea hopper, Bandedwinged whitefly, Plant bugs (excludes <i>Lygus hesperus</i>), Green stink bug, Southern green stink bug, Bollworm/budworm (ovicidal effect)	1 - 2
Pest Suppressed	
Lygus bug (<i>Lygus hesperus</i>), Whiteflies (other than bandedwinged whitefly)	2

Restrictions:

- Pre-harvest interval (PHI): 14 days
- Minimum interval between applications: 7 days
- Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 10 fluid ounces per acre (0.31 lb. AI per acre)
- Regardless of formulation or method of application, apply no more than 0.5 lb. active ingredient per acre per season, including seed, soil, and foliar uses.
- Do not graze treated fields after any application of Couraze 4 Insecticide.
- Couraze 4 Insecticide may only be applied through properly calibrated ground, aerial, or chemigation application equipment.

Applications: Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.

Tank Mix Applications

Pests Controlled (In addition to pests listed above)	Couraze 4 Insecticide Rate fluid ounces per acre	Bidrin® 8* Rate fluid ounces per acre
For early season control of: Thrips	1 – 1.5	1.6 – 3.2
For mid to late season control of: Plant bugs, Stink bugs (including Brown stink bug), Grasshoppers, Saltmarsh caterpillar, Cotton leaf perforator	1 – 1.5	4.0 – 8.0
Restrictions (In addition to Restrictions listed above): * Refer to the Bidrin 8 product label for specific use instructions. Observe all restrictions and precautions that appear on the label.		

PEANUT¹

Soil Application

Pests Controlled	Rate: Fluid ounces per acre*
Aphids Leafhoppers Whiteflies	8.0 – 12.0
Pests Suppressed	
Thrips	8.0 – 12.0
Restrictions <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 14 days • Maximum COURAZE allowed per season: 12.0 fluid ounces/Acre (0.38 lb AI/Acre) Applications Apply specified dosage in one of the following methods: In-furrow spray during planting directed on or below seed or by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	
Plant virus management Increases in Tomato spotted wilt virus (TSWV) incidence have been observed with applications of COURAZE on multiple varieties of peanut. This may also be the case with other tospoviruses, or other viruses transmitted by various thrips species or perhaps, other pests. Prior to applying COURAZE to peanuts, consult with the State, Cooperative Extension Service, or Cheminova representative, for instructions. Growers are advised to weigh insect control benefits against potential increase in viral disease levels. In areas where TSWV or other tospovirus are endemic, growers are encouraged to use virus resistant varieties and consult the University of Georgia, Tomato spotted wilt virus index, before applying COURAZE.	
^{1/} Use not permitted in California unless otherwise directed by supplemental labeling.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre*
Aphids Leafhoppers Whiteflies	1.4

Restrictions

- Pre-Harvest Interval (PHI): 14 days
- Minimum interval between applications: 5 days
- Maximum COURAZE allowed between crop season: 4.2 fluid ounces/Acre (0.13 lb AI/Acre)
- 1/ Use not permitted in California unless otherwise directed by supplemental labeling.

POTATO**Soil Application**

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Potato psyllid	0.45 – 0.65	6.5 – 10.0
Pests/Diseases Suppressed	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Symptoms of: Potato leaf roll virus (PLRV), Potato yellows, Net necrosis, Wireworms (with in-furrow spray at-planting)	0.45 – 0.65	6.5 – 10.0
Restrictions: <ul style="list-style-type: none"> • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 10.0 fluid ounces per acre (0.31 lb active ingredient per acre) Application: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • In-furrow spray during planting directed on seed pieces or seed potatoes. • Subsurface side-dress on both sides of the row covered with 3 or more inches of soil. • Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil. • Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. For effective pest control or suppression, Couraze 4 Insecticide applications must be placed below soil surface and in contact with seed piece or within root zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of Couraze 4 Insecticide may be made in a 2 to 4 inch band (width of planter-shoe opening) and completely covered. 		

Seed Piece Application

Pests Controlled	Rate: Fluid ounces per 100 lbs. of seed	Rate: Fluid ounces per acre*
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Potato psyllid, Wireworms (seed-piece protection)	0.2-0.4	4.0-8.0
Pests/Diseases Suppressed	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Symptoms of: Potato leaf roll virus (PLRV), Potato yellows, Net necrosis	0.4	8.0
Restrictions: <ul style="list-style-type: none"> • Maximum Couraze 4 Insecticide allowed per crop season when making seed piece treatment applications: 10.0 fluid ounces per acre (0.31 lb. active ingredient per acre) • Do not use treated seed-pieces for food feed, or fodder. • Do not apply any subsequent application of Couraze 4 Insecticide (in-furrow), Gaucho, Leverage, Provado, or similar Group 4A insecticide following a seed piece treatment with Couraze 4 Insecticide. Application: Apply specified dosage as a diluted spray onto seed pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part Couraze 4 Insecticide. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after Couraze 4 Insecticide application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed pieces as soon as possible after treating avoiding prolonged exposure of Couraze 4 Insecticide treated seed pieces to sunlight and in accordance with the instruction of your local Extension service: .		
* Based on a seeding rate of 2000 lbs. per acre.		

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre*
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Psyllids	1.5

Restrictions:

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 7 days
- Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 6.4 fluid ounces per acre (0.2 lb AI per acre)

Applications: Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.

SOYBEAN

Foliar Application¹

Pests Controlled	Rate: Fluid ounces per acre*
Aphids, Bean leaf beetle, Cucumber beetles/Rootworm adults, Japanese beetle (adults), Leafhoppers, Whiteflies	1.5
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 7 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 3.65 fluid ounces per acre (0.13 lb AI per acre) 	
¹ Use not permitted in California unless otherwise directed by supplemental labeling.	
Applications: Apply specified rate per acre as a broadcast or directed foliar spray through properly calibrated ground, aerial or chemigation application equipment. Thorough uniform coverage is necessary to achieve optimum control.	

TOBACCO

Soil Application

Pests Controlled	Rate: Fluid Ounces per 1000 plants (as seedling tray drench)	Rate: Fluid ounces per 1000 plants (in-furrow or transplant-water)
Aphids, Flea beetles	0.5	0.7
Mole crickets, Whiteflies, Wireworms	0.7 – 1.4	0.9 – 1.4
Pests/Disease Suppressed		
Cutworms Symptoms of: Tomato spotted wilt virus (TSWV)	1.4	1.4
Restrictions: <ul style="list-style-type: none"> • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications or foliar sprays to seedlings: 16.0 fluid ounces per acre (0.5 lb. active ingredient per acre) • Pre-Harvest Interval (PHI): 14 days 		
Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Uniform, broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting followed immediately by overhead irrigation to wash Couraze 4 Insecticide from foliage into potting media. Failure to wash Couraze 4 Insecticide from foliage may result in a reduction in pest control. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots. • In-furrow spray or transplant water drench during setting. • Carefully handle transplants during setting to avoid dislodging treated potting media from roots. 		
Application Efficacy: Proper tray drench applications of Couraze 4 Insecticide have been shown to be the most efficacious method of application. However, the specified rate of Couraze 4 Insecticide may be applied as combination of the tray drench in the planthouse and/or transplant water drench in field. Adverse growing conditions may cause a delay in uptake of Couraze 4 Insecticide into the plant and a delay in control.		

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids	0.8 – 1.6

Flea beetles, Japanese beetles	1.6
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 14 days • Minimum interval between applications: 7 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 9.0 fluid ounces per acre (0.28 lb. AI per acre) Applications: Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.	

VEGETABLE AND SMALL FRUIT CROPS

CUCURBIT VEGETABLES¹ : Crops of Crop Group 9: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber; Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis* melon including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

Soil Application

Field Application Rates. See details below for additional planthouse rates.	
Pests Controlled	Rate: Fluid ounces per acre.
Aphids, Cucumber beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	8.0 – 12.0
Pests/Diseases Suppressed	
Bacterial wilt (as vectored by various cucumber beetles), Leaf silvering resulting from whitefly feeding	12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days • Maximum Couraze 4 Insecticide allowed per application when making soil applications: 12.0 fluid ounces per acre (0.38 lb/active ingredient per acre) Applications: Apply the specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. • In-furrow spray directed on or below seed. • Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2 inches with sufficient irrigation within 24 hours of application. • Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting. • Post-seeding drench, transplant water drench, or hill drench. • Subsurface side-dress on both sides of each row. Couraze 4 Insecticide must be incorporated into root zone. 	

Planthouse Application Rates	
Pest Controlled	Rate: Fluid ounces per 1000 plants
Aphids, Whiteflies	0.05
Restrictions: <ul style="list-style-type: none"> • Maximum amount Couraze 4 Insecticide applied in the planthouse: 0.05 fluid ounces (0.001568 lb. active ingredient per 1000 plants) • Maximum number Couraze 4 Insecticide applications in planthouse: 1 Applications: Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following methods: <ul style="list-style-type: none"> • Uniform, broadcast high volume foliar spray, followed immediately by sufficient overhead irrigation to wash Couraze 4 Insecticide from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash Couraze 4 Insecticide from foliage may result in reduced pest control. 	

<ul style="list-style-type: none"> • Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray. <p>The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher listed rates or increased number of applications in planthouse may result in significant plant injury. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots.</p> <p>Crop Tolerance Confirmation: Not all varieties of cucurbit vegetables have been tested for tolerance to Couraze 4 Insecticide applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.</p> <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p> <p>* Use not permitted in CA unless otherwise directed by supplemental labeling.</p>
--

GREENHOUSE VEGETABLES; (Mature plants in production greenhouses ONLY): Cucumber, Tomato only

Soil Application

Pests Controlled	Rate: Fluid ounces per 1000 plants
Aphids, Whiteflies	0.7
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 0 day • Maximum number Couraze 4 Insecticide applications per crop season when making soil applications: 1 • Maximum amount of product allowed per crop season: 0.7 fluid ounce (0.022 lb. AI)/1,000 plants. <p>Applications: Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Make application only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool or other soil-less media, or plants growing hydroponically. Do not apply to immature plants since phytotoxicity may occur. Make applications when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (<i>Orius</i> spp.) can occur when Couraze 4 Insecticide is applied.</p> <p>Many varieties of vegetables have been tested for tolerance to Couraze 4 Insecticide and show good safety. However, certain varieties may show more sensitivity to Couraze 4 Insecticide. Therefore, treat a few plants before treating the whole greenhouse.</p> <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p>	

FRUITING VEGETABLES¹: Crops of Crop Group 8 plus Okra: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento, and sweet), Tomato, Pepinos, Tomatillo

Soil Application

Field Application Rates. See details below for additional planthouse rates.	
Pests Controlled	Rate: Fluid ounces per Acre
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	Okra and Pepper; 8.0-16.0 Other Crops; 8.0-12.0
Diseases Suppressed	
Symptoms of: Tomato mottle virus, Tomato spotted wilt virus, Tomato yellow leaf curl virus	Okra and Pepper; 8.0-16.0 Other Crops; 8.0-12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days • Maximum Couraze 4 Insecticide allowed on pepper and okra crops per application when making soil applications: 16.0 fluid ounces/Acre (0.5 lb. A.I. per acre) • Maximum Couraze 4 Insecticide allowed on other fruiting vegetable crops per application when making soil applications: 12.0 fluid ounces/Acre (0.38 lb. A.I. per acre) • DO NOT use on crops grown for seed unless allowed by state-specific supplemental labeling. <p>Applications: Apply specified dosage in one of the following methods:</p> <ul style="list-style-type: none"> • Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. • In-furrow spray directed on or below seed. • Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation within 24 hours of application. • Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting. • Post-seeding drench, transplant-water drench, or hill drench. • Subsurface side-dress on both sides of each row. Couraze 4 Insecticide must be incorporated into root-zone. 	

Planthouse Application Rates	
Pests Controlled	Rate: Fluid ounces per 1000 plants
Aphids, Whiteflies	0.05
Restrictions: <ul style="list-style-type: none"> Maximum amount Couraze 4 Insecticide applied in the planthouse: 0.05 fluid ounces (0.001568lb.A.I. per 1000 plants. Maximum number Couraze 4 Insecticide applications in planthouse: 1 Applications: Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners: <ul style="list-style-type: none"> Uniform, broadcast high-volume foliar spray followed immediately by sufficient overhead irrigation to wash Couraze 4 Insecticide from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash Couraze 4 Insecticide from foliage may result in reduced pest control. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray. <p>The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher listed rates or increased number of applications in planthouse may result in significant plant injury. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots.</p> Crop Tolerance Confirmation: Not all varieties of fruiting vegetables have been tested for tolerance to Couraze 4 Insecticide applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse. <p>Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p> <p>* Use not permitted in CA unless otherwise directed by supplemental labeling.</p>	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Colorado potato beetle, Leafhoppers, Whiteflies	1.5 – 2.5
Pepper weevil	2.5
Restrictions: <ul style="list-style-type: none"> Pre-harvest interval (PHI): 0 day Minimum interval between applications: 5 days Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 7.6 fluid ounces per acre (0.24 lb. AI per acre) Applications: <p>Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest Crop Groups.</p> <p>For pepper weevil, apply specified dosage of Couraze 4 Insecticide by ground equipment only, timing applications prior to a damaging population becoming established. Good coverage of foliage and fruit is necessary for optimum control. Applications of Couraze 4 Insecticide must be incorporated into a full-season program where alternations of effective products from multiple classes of chemistry and different modes of action are utilized in a blocked or windowed approach.</p> <p>For additional information, please contact your Cheminova, Inc. representative, Extension Specialist or crop advisor.</p> <p>When targeting adult whiteflies, use the higher listed rates.</p> <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p>	

GLOBE ARTICHOKE ¹

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> Pre-harvest interval (PHI): 7 days Maximum Couraze 4 Insecticide allowed per crop season; 16.0 fluid ounces/Acre (0.5 lb AI/Acre) Applications: <p>Apply specified dosage in the following method:</p> <ol style="list-style-type: none"> Chemigation into root zone through low pressure drip, trickle, micro sprinkler or equivalent equipment. In-furrow spray of planting directed on or below seed. <p>¹ Use not permitted in California unless otherwise directed by Supplemental labeling.</p>	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers	1.6 – 4.0
Restrictions: <ul style="list-style-type: none"> • Pre-harvest interval (PHI): 7 days • Minimum interval between applications: 14 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16 fluid ounces per acre (0.5lb AI per acre) Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.	

HERBS¹ : Crops of Crop Subgroup 19A: Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Chamomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Cilantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

Soil Application

Pests Controlled	Rate: Fluid ounces per Acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	8.0 – 12.0
Pests Suppressed	
Thrips (foliage feeding thrips only)	8.0 – 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 14 days • Maximum Couraze 4 Insecticide per season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI/Acre) Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • In-furrow spray during planting directed on or below seed. • In-furrow spray or transplant water drench during selling or transplanting. • Shankd into or below eventual seed line. • Chemigation into root zone through low pressure drip, trickle micro-sprinkler, or equivalent equipment. Crop Tolerance Confirmation: Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat and evaluate only small areas or numbers of plants of each prior to commercial use.	
¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per Acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.4
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 5 days • Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 4.2 fluid ounces per acre (0.13 lb. AI per acre) Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.	
Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov)	

for latest Crop Groups.

Couraze 4 Insecticide may be applied through properly calibrated ground and aerial application equipment. Thorough coverage with direct contact of the spray material to the target pests is required for optimum control. The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage and control.

Crop Tolerance Confirmation: Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat and evaluate only small areas or numbers of plants of each prior to commercial use.

¹Not permitted for use in California unless otherwise directed by Supplemental label.

BRASSICA (COLE) LEAFY VEGETABLES¹ : Crops of Crop Group 5: Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens Mustard spinach, Rape greens, Turnip tops (leaves).

Soil Application

Pests Controlled	Rate: Fluid ounces per acre (on 36 inch rows)
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	5.0 – 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days • Maximum Couraze 4 Insecticide allowed per application when making soil applications: 12.0 fluid ounces/Acre (0.38lb AI per acre) Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler or equivalent equipment. • In-furrow spray directed on or below seed. • Narrow (2 inches or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1-1/2 inches with sufficient irrigation within 24 hours of application. • Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting. • Post-seeding drench, transplant water drench, or hill drench. • Subsurface side-dress on both sides of each row. Couraze 4 Insecticide must be incorporated into root zone. 	
¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per Acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.5
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 5 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 7.7 fluid ounces/Acre (0.24 lb AI per acre) Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest Crop Groups.	
¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

LEAFY VEGETABLES¹ - Crops of Crop Subgroup 4A plus Watercress: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Rocket), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Com salad, Cress (garden). Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water). Watercress (upland)

Soil Application

Pests controlled	Rate: Fluid ounces per acre (on 36 inch rows)
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	5.0 – 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days 	

- Maximum Couraze 4 Insecticide allowed per application when making soil applications: 12.0 fluid ounces per acre (0.38 lb AI per Acre)
 - For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following application.
 - Applications must be made to fully leafed-up canopies only.
 - DO NOT apply to native cress growing in streams or other bodies of water.
- Applications:** Apply specified dosage in one of the following methods:
- Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment.
 - In-furrow spray directed on or below seed.
 - Narrow (2 inches or less) surface band spray over seed line during planting incorporated to a depth of 1 to 1-1/2 inches with sufficient irrigation within 24 hours of application.
 - Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting.
 - Post-seeding drench, transplant water drench or hill drench.
 - Subsurface side-dress on both sides of each row. Couraze 4 Insecticide must be incorporated into root zone.
- ¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.5
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 5 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 7.7 fluid ounces per acre (0.24lb AI per acre) • For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the applications. Applications must be made to fully leafed-up canopies only. <p>Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.</p> <p>Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest Crop Groups.</p> <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p>	

LEAFY PETIOLE VEGETABLES¹ — Crops of **Crop Subgroup 4B:** Cardoon, Celery, Lettuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss chard \

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, thrips (foliage feeding thrips only), Whiteflies	5.0 – 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 45 days • Maximum Couraze 4 Insecticide allowed per application when making soil applications: 12.0 fluid ounces/Acre (0.38lb AI per acre) <p>Applications: Apply specified dosage in one of the following methods:</p> <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. • In-furrow spray directed on or below seed. • Narrow. (2 inches or less) surface band spray over seed line during planting incorporated to a depth of 1 to 1-1/2 inches with sufficient irrigation within 24 hours of application. • Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting. • Post-seeding drench, transplant water drench, or hill drench. • Subsurface side-dress on both sides of each row. Couraze 4 Insecticide must be incorporated into root zone. <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p>	

LEGUME VEGETABLES¹ (except soybean, dry): Crops of **Crop Group 6:** Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean

Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white-lupin, and white sweet lupin)

Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea; garden pea, green pea, snow pea, sugar snap pea)
Other Beans and Peas [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	8.0 – 12.0
Diseases suppressed	
Symptoms of: Bean common mosaic virus (BCMV), Bean golden mosaic virus (BGMV), Beet curly top hybrigeminivirus (BCTV)	8.0 – 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre) Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. • In-furrow spray at planting directed on or below seed. • In a narrow (2 inches or less) surface band over seed-line during planting incorporated to a depth of 1 to 1-1/2 inches with sufficient irrigation with 24-hours following application. • In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. • As a post-seeding drench, transplant drench, or hill drench. ¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Whiteflies	1.4
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 7 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 4.2 fluid ounces per acre (0.13lb AI per acre) Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest Crop Groups.	
¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

ROOT VEGETABLES¹ - Crops of Crop Subgroup 1B (except Sugarbeet) and Kava: Beet (garden)², Burdock (edible)², Carrot², Celeriac², Chervil (turnip-rooted)², Chicory², Ginseng, Horseradish, Kava2/3, Parsley (turnip-rooted), Parsnip², Radish², Oriental radish (diakon)², Rutabaga², Salsify (oyster plant), Salsify (black)², Salsify (Spanish), Skirret, and Turnip²

Soil Application

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	0.35 - 0.85	5.0 - 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre) • Maximum Couraze 4 Insecticide soil applications per crop season: 1 Application: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. • In-furrow spray (rate specified per 1000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting. • In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting. Length of Control and Use Rates: The rate applied affects the length of control. Use higher listed rates where infestations occur later in crop development, or where pest pressure is continuous. Couraze 4 Insecticide rates less than 0.7 fluid ounces/1000 row-feet will not provide adequate residual pest control. Couraze 4 Insecticide treated crops grown on very high organic matter soils (muck) may also require additional pest management control. <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p> <p>² Tops or greens from these crops may be utilized for food or feed.</p> <p>³ Not permitted for use in California unless otherwise directed by Supplemental labeling.</p>		

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.4
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 5 days • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 1.4 fluid ounces per acre (0.044 lb AI per acre) on Radish, 4.2 fluid ounces per acre (0.13 lb AI per acre) on other crops. • Maximum Couraze 4 Insecticide soil application(s) per crop season: 1 on radish, 3 on all other crops Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest Crop Groups. <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p> <p>² Tops or greens from these crops may be utilized for food or feed.</p> <p>³ Not permitted for use in California unless otherwise directed by Supplemental labeling.</p>	

TUBEROUS and CORM VEGETABLES¹ - Crops of Crop Subgroup 1C: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet)², Chayote (root), Chufa, Dasheen (taro)², Ginger, Leren, Sweet potato, Tanier (cocoyam)², Turmeric, Yam bean (jicama, manioc pea). Yam (true)² (For applications on potato, see Field Crops section)

Soil Application

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	0.35 - 0.85	5.0 - 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI) from planting application: 3 days (leaves); 125 days (corms) • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 12.0 fluid ounces/Acre (0.38lb AI per acre) • Maximum Couraze 4 Insecticide soil applications per crop season: 1 Applications: Apply specified dosage in one of the following methods: <p>¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.</p> <p>² Tops or greens from these crops may be utilized for food or feed.</p> <p>³ Not permitted for use in California unless otherwise directed by Supplemental labeling.</p>		

- In-furrow spray (rate specified per 1000 row-feet) over planting materials (hulis) or shanked-in 1 to 2 inches below hulis depth at planting.
 - Side-dress not more than 0.3 fluid ounces/1000 row-feet no later than 45 days after planting. Observe the same PHI as above.
- Length of Control and Use Rates:** The rate applied affects the length of control. Use higher listed rates where infestations occur late in crop development, or where pest pressure is continuous. Couraze 4 Insecticide rates less than 0.35 fluid ounces/1000 row-feet may not provide adequate residual pest control. Couraze 4 Insecticide treated crops grown on very high organic matter soils (muck) may also require additional pest management control.
- ¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.
- ² Tops or greens from these crops may be utilized for food or feed.

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.4
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 5 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 1.4 fluid ounces per acre (0.044 lb. AI per acre) on Radish, 4.2 fluid ounces per acre (0.13 lb. AI per acre) on other crops. • Maximum Couraze 4 Insecticide application(s) per crop season: 3 on all crops Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest Crop Groups.	
¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
² Tops or greens from these crops may be utilized for food or feed.	

STRAWBERRY¹

Soil Application

Annual and Perennial Crops	
Pests Controlled	Rate: Fluid ounces per acre
Aphids, Whiteflies	12.0 – 16.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 14 days • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 16.0 fluid ounces/Acre (0.50 lb AI per acre). Do not apply immediately prior to bud opening or during bloom or when bees are actively foraging.	
Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening. • As a plant material or plant hole treatment just prior to, or during transplanting. • As a band spray over-the-row in a minimum of 20 gallons of water per acre, followed immediately by overhead irrigation to incorporate product into root zone. Do not use plastic or other mulches that limit movement of Couraze 4 Insecticide into root zone. The rate applied affects the length of control. Use higher listed rates where infestations may occur later in crop development or where pest pressure is continuous.	
Post-harvest Use on Perennial Crops	
Pests Controlled	Rate: Fluid ounces per acre
White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	8.0 – 12.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 14 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 12.0 fluid ounces per acre (0.38lb A.I. per acre) • DO NOT use both application methods on the same crop in the same season. Applications: Apply a single application post harvest to coincide with renovation of strawberry fields and during active egg laying period of beetles. Apply specified dosage of Couraze 4 Insecticide in one of the following methods: <ul style="list-style-type: none"> • As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre. • As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed. • As a chemigation application with 600 to 1000 gallons of water followed by 0.1 to 0.25 inches irrigation. Incorporation Requirement: All soil-surface applications must be followed by 0.25 inches of rainfall or overhead irrigation water	

per acre within 2 hours of application. Failure to adequately incorporate Couraze 4 Insecticide into egg-deposition zone may result in decreased activity.

Foliar Application

Annual and Perennial Crops	
Pests Controlled	Rate: Fluid ounces per acre
Aphids, Spittlebugs, Whiteflies	1.5
Restrictions: <ul style="list-style-type: none"> • Pre-harvest Interval (PHI): 7 days • Minimum interval between applications: 5 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 4.6 fluid ounces per acre (0.14 lb. AI per acre). • Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging. • DO NOT make both a soil and foliar application on the same crop in the same season. Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.	

SUGARBEET

Soil Application-For use only in CA

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Whiteflies, Flea beetles	3.0 – 6.0
Diseases Suppressed	
Symptoms of: Western yellows/Beet curly tophybrigeminivirus (BCTV)	3.0 – 6.0
Restrictions: <ul style="list-style-type: none"> • Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 6.0 fluid ounces/Acre (0.18 lb AI per acre) • Do not apply immediately prior to bud opening or during bloom or when bees are actively foraging. Applications: Apply specified dosage in the following method: Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting. The low rate may be applied to aid establishment of stands in Whitefly areas, or for early season control of the other pests listed.	
¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

Rate: Fluid ounces/ Acre	Rate: Fluid ounces/1000 row-feet									
	Based on average row spacing (in inches):									
	10	15	20	25	30	34	36	38	40	45
5.0	0.11	0.16	0.22	0.28	0.33	0.38	0.39	0.41	0.44	0.49
5.5	0.13	0.18	0.24	0.30	0.37	0.41	0.44	0.46	0.48	0.54
6.0	0.13	0.20	0.26	0.33	0.39	0.45	0.47	0.51	0.53	0.60
6.5	0.14	0.22	0.29	0.36	0.43	0.48	0.52	0.54	0.57	0.64
7.0	0.15	0.23	0.31	0.38	0.46	0.53	0.55	0.59	0.62	0.69
7.5	0.16	0.25	0.33	0.41	0.49	0.56	0.60	0.63	0.66	0.75
8.0	0.17	0.26	0.36	0.44	0.53	0.60	0.63	0.67	0.70	0.79
8.5	0.18	0.28	0.38	0.47	0.56	0.63	0.68	0.71	0.75	0.84
9.0	0.20	0.30	0.39	0.49	0.60	0.68	0.71	0.75	0.79	0.89
9.5	0.21	0.31	0.41	0.52	0.63	0.71	0.75	0.79	0.84	0.94
10.0	0.22	0.33	0.44	0.55	0.66	0.75	0.79	0.84	0.89	0.99
10.5	0.23	0.34	0.46	0.57	0.69	0.78	0.83	0.87	0.92	1.03
11.0	0.24	0.37	0.48	0.61	0.72	0.83	0.87	0.92	0.97	1.09

11.5	0.25	0.38	0.51	0.63	0.76	0.86	0.91	0.97	1.01	1.14
12.0	0.26	0.39	0.53	0.66	0.79	0.90	0.95	1.00	1.06	1.18
12.5	0.28	0.41	0.55	0.69	0.83	0.93	0.99	1.05	1.10	1.24
13.0	0.29	0.43	0.57	0.71	0.86	0.98	1.03	1.09	1.14	1.29
13.5	0.30	0.45	0.60	0.75	0.89	1.01	1.07	1.13	1.18	1.33
14.0	0.31	0.46	0.62	0.77	0.92	1.05	1.10	1.17	1.23	1.39

Length of Control and Application Rates: The Couraze 4 Insecticide rate applied affects the length of control and to a considerable extent, the degree of control or effect. Row spacing X Couraze 4 Insecticide rate combinations in shaded blocks may not provide adequate residual pest control and should not be used for long-term, residual control. Use higher labeled rates where infestation may occur later in crop development or where pest pressure is continuous. Cheminova, Inc. offers no warranty for use of Couraze 4 Insecticide at rates below 0.36 fluid ounces/1000 row-feet.

TREE, BUSH, and VINE CROPS

BANANA and PLANTAIN¹

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers	8.0 – 16.0
Pests Suppressed	
Scales	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> Pre-Harvest Interval (PHI): 0 day Maximum Couraze 4 Insecticide allowed per crop season when making soil applications: 16.0 fluid ounces per Acre (0.5 lb AI per A) Applications: Apply specified dosage in the following method: <ul style="list-style-type: none"> Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. ¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Thrips	3.2
Restrictions: <ul style="list-style-type: none"> Pre-Harvest Interval (PHI): 0 day Minimum interval between applications: 14 days Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16.0 fluid ounces Per acre (0.5lb AI per acre) Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Apply specified dosage as a broadcast or directed spray to infested area insuring thorough coverage. Couraze 4 Insecticide may be applied through properly calibrated ground or aerial application equipment. Aerial applications of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. Addition of an organosilicone adjuvant at a rate not to exceed 2.0 fluid ounces per 100 gallons, finished spray solution may improve coverage and pest control.	
¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

BUSHBERRY : Crops of Crop Subgroup 13B: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Saial

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Japanese beetle: (adults, feeding on foliage) White grub complex: (grubs of Asiatic garden beetle, European and Masked chafer. Japanese beetle and Oriental beetle)	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> Pre-Harvest Interval (PHI): 7 days 	

- Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.50 lb AI per acre)
- Do not apply pre-bloom or during bloom or when bees are actively foraging.

Applications:

Apply specified dosage in one of the following methods:

- Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment.
- 18-inch band on each side of the row followed by irrigation immediately after application.

For optimal grub control, apply Couraze 4 Insecticide to control 1st or 2nd instar larvae. Application may be made post-bloom up to 7 days prior to harvest or post-harvest until October 1st. For optimum control of Japanese beetle larvae, make applications from June 1 to July 15. Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root zone will help protect berry plant roots from grub feeding.

Apply Couraze 4 Insecticide to moist soil. If necessary, apply one hour of irrigation water immediately before application of Couraze 4 Insecticide. To ensure maximum efficacy of soil surface spray, 1/2 to 1 inch of irrigation water or rainfall should be applied or received within 24 hours of application of Couraze 4 Insecticide to facilitate movement into the soil and into the root zone.

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters	1.2-1.6
Japanese beetle (adults), Thrips (foliage feeding thrips only)	2.4-3.2
Blueberry maggot	3.2

Restrictions:

- Pre-Harvest Interval (PHI): 3 days
- Minimum interval between applications: 7 days
- Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16.0 fluid ounces per acre (0.5lb AI per acre)
- Maximum number of Couraze 4 Insecticide applications per crop season when making foliar applications: 5
- Minimum application volume (water): 20.0 GPA-ground, 5.0 GPA-aerial.
- Do not apply pre-bloom or during bloom or when bees are actively foraging.

Applications: Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (www.epa.gov) for latest crop groups.

CANEBERRY - Crops of Crop Subgroup 13A: Blackberry (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marion berry, nectarberry, olallieberry, Oregon evergreen berry, phenomenal berry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these) **Raspberry** (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*)

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Whiteflies	8.0 – 16.0
Rednecked cane borer	12.0 – 16.0
Pests Suppressed	
Thrips (foliage feeding thrips only)	16.0

Restrictions:

- Pre-Harvest Interval (PHI): 7 days
- Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per acre)
- Do not apply pre-bloom or during bloom or when bees are actively foraging.

Soil Application: Apply specified dosage in one of the following methods:

- Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment.
- Basal, soil drench in a minimum of 500 gallons solution per acre.

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphid Leafhoppers Whiteflies	3.2
Restrictions:	

- Pre-Harvest Interval (PHI): 3 days
- Minimum interval between applications: 7 days
- Maximum Couraze allowed per crop season: 9.6 fluid ounces per Acre (0.3 lb AI per acre).
- Do not apply pre-bloom or during bloom or when bees are actively foraging.

¹ Use not permitted in California unless otherwise directed by supplemental labeling.

CITRUS (Containerized) - Crops of Crop Group 10: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, Tangelo, White sapote (*Casimiroa* spp.) and other cultivars and/or hybrids of these.

Soil Application

Pests Controlled	Rate ml/ft ³ container media
Aphid, Asian citrus psyllid, Blackfly, Citrus leafminer, Leafhoppers/Sharpshooters, Mealybugs, Scales, Whiteflies	0.38
Citrus root weevil (larval complex)	0.63 – 1.25
Pests Suppressed	
Citrus thrips (foliage feeding thrips only)	1.25
Application: Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of Couraze 4 Insecticide per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, treatment should be made at planting prior to insect infestation. Retreat if necessary. For control of larvae of the citrus root weevil complex, application should be made prior to neonate larvae entering potting media. Utilize higher dosage for heavy infestations.	

CITRUS (Field): Crops of Crop Group 10: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, Tangelo, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these.

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Asian citrus psyllid, Blackfly, Citrus leafminer, Leafhoppers/Sharpshooters, Mealybugs, Scales, Termites (FL only), Whiteflies	8.0 – 16.0
Pests/Diseases Suppressed	
Citrus nematode, Symptoms of: Citrus tristeza virus (CTV) through vector control, Citrus yellows, Thrips (foliage feeding thrips only)	16.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 0 day • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per Acre) Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil should be lightly pre-wetted to break soil surface tension prior to applications of Couraze 4 Insecticide. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move Couraze 4 Insecticide into root zone. Allow 24 hours before initiating subsequent irrigations. • Soil surface band spray on both sides of the tree. Bands should overlap at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root zone. This method is suitable for very coarse soils with 0.75% organic matter or less. • Drench to base of tree not exceeding one quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only for trees up to 8 feet tall. • For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. • For suppression of citrus nematode, apply specified dosage through low pressure chemigation or soil surface band spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of Couraze 4 Insecticide over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response. 	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Asian citrus psyllid, Blackfly, Leafhoppers/Sharpshooters, Leafminers, Mealybugs, Scales, Whiteflies	4.0 – 8.0 (depending on tree size, target pest and infestation pressure)
Pests Suppressed	
Thrips (foliage feeding thrips only)	8.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 0 day • Minimum interval between applications: 10 days • Maximum COURAZE 4 INSECTICIDE allowed per season when making foliar applications: 16 fluid ounces per acre (0.5 lb AI per acre) • Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging. Applications: Scales - time applications to the crawler stage. Treat each generation. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (www.epa.gov) for latest crop groups.	

COFFEE¹

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Leafminers	8.0 – 16.0
Pests Suppressed	
Scales	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces per Acre (0.5 lb AI per acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. • Subsurface side-dress shanked into the root-zone on both side of the plants followed by irrigation. • Basal, soil drench in sufficient water to insure incorporation into the root zone followed by irrigation. ¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Whiteflies	3.2
Pest suppressed	
Scales	3.2
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 7 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16 fluid ounces per Acre (0.5lb AI/Acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging. Applications: Apply specified dosage as a broadcast or directed spray to infested area insuring thorough coverage. Couraze 4 Insecticide may be applied through properly calibrated ground or aerial application equipment. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. COURAZE 4 INSECTICIDE may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. ¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

CRANBERRY

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Rootgrubs (Scarabaeidae), Rootworms (Chrysomelidae)	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 30 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5lb AI/Acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging Applications: Apply Couraze 4 Insecticide to moist soil. Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre. • As a chemigation application with 600 to 1000 gal water. Immediately upon application, Couraze 4 Insecticide must be incorporated into root-zone by 0.1-0.3 inches water/Acre: either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.	
Rootgrubs and Rootworms Best control may be achieved when application is made post-bloom immediately after bees are removed. Applications should target early instar larvae. Couraze 4 Insecticide has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of the Couraze 4 Insecticide and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.	

GRAPE - American bunch grape, Muscadine grape and Vinifera grape

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
European fruit lecanium, Leafhoppers/Sharpshooters, Mealybugs, <i>Phylloxera</i> * spp	8.0 – 16.0
Pest/Disease Suppressed	
Grapeleaf skeletonizer, Nematodes, Pierce's disease	16.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 30 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5lb AI/Acre) Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. • Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation. • Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation. • For suppression of nematodes, apply 7 fluid ounces in a single application or two 3.5 fluid ounce applications on a 30 to 45 day interval. Treatment(s) should be applied only by 1) chemigation into root zone through above ground low pressure drip, trickle, micro-sprinkler, or equivalent equipment; or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root zone of the plant. Repeated and regular use of Couraze 4 Insecticide over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response. For optimum results, make application(s) between bud-break and the pea-berry stage. Apply a total of 14 7 fluid ounces/Acre under any of the following conditions: <ol style="list-style-type: none"> 1. Where vigorous vine growth is expected; 2. In warmer growing areas; 3. Where mealybug and European fruit lecanium populations are expected to be heavy; 4. Where vine populations exceed 600 per acre, or 5. For suppression of nematodes. * Repeated and regular use of Couraze 4 Insecticide over several, consecutive growing seasons controls existing <i>Phylloxera</i> infestations over time or prevents <i>Phylloxera</i> from becoming established.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Leafhoppers, Sharpshooters, Mealybugs	1.2 – 1.6
Grapeleaf skeletonizer	1.6
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 0 days • Minimum interval between applications: 14 days 	

- Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 3.2 fluid ounces/Acre (0.1 lb AI/Acre)

Couraze 4 Insecticide may be applied by ground application only.

Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application, of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines.

HOP¹

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids	9.6
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 60 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 9.6 fluid ounces/Acre (0.3 lb AI/Acre) Applications: Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> • Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. • Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation. • Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation. Apply a higher dosage where extended residual control is desired or for treating larger vines or vines with dense foliage volume.	
¹ Use not permitted in California unless otherwise directed by supplemental labeling.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids	3.2
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 28 days • Minimum interval between applications: 21 days • Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 9.6 fluid ounces/Acre (0.3 lb AI/Acre) Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines.	

POME FRUIT – Crops of Crop Group 11: Apple, Crabapple, Loquat, Mayhew, Pear (including Oriental pear), Quince

Soil application

Pests Controlled	Rate: Fluid ounces per acre
Aphids (including Woolly apple aphid), Leafhoppers	8.0 – 12.0
Restrictions: Pre-harvest interval (PHI): 21 days <ul style="list-style-type: none"> • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI/Acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging Applications: Apply specified dosage in the following method: <ul style="list-style-type: none"> • Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. 	

Foliar application

Pests Controlled	Rate: Fluid ounces per acre
Leafhoppers	1.6 – 3.2
Aphids (except Woolly apple aphid), Apple maggot, Leafminers, San Jose scale	3.2
FOR PEAR ONLY: Mealybugs, Pear psylla	8

Restrictions:

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 10 days
- Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 16 fluid ounces/Acre (0.5 lb AI/Acre)
- Do not apply pre-bloom or during bloom or when bees are actively foraging

Applications: Applications targeting apple maggot should be combined with manufacturer's application rate of a sticker.

Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (www.epa.gov) for latest crop groups.

POMEGRANATE¹**Soil Application**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Whiteflies	8.0 – 16.0
Restrictions:	
<ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 0 day • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging. 	
Applications: Apply specified dosage in the following method:	
<ul style="list-style-type: none"> • Chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. 	
¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Whiteflies	3.2
Pests suppressed	
Scales	3.2
Restrictions:	
<ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 7 days • Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 9.6 fluid ounces/Acre (0.3 lb AI/Acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging. 	
Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines.	
¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

STONE FRUIT - Crops of Crop Group 12: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh and dried)

Soil Application

In-field, Soil Application	
Pests Controlled	Rate: Fluid ounces per acre
Aphids (including Woolly apple aphid), Leafhoppers	8.0 – 12.0
Restrictions:	
<ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 21 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging. 	
Applications: Apply specified dosage in the following method:	
<ul style="list-style-type: none"> • Chemigation into the root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. 	
Preplant, Root Dip Application	

Pests Controlled	Rate: Fluid ounces per 10 gallons root-dip solution
Black peach aphid (infesting roots)	1.0
Mix Couraze 4 Insecticide at 1.0 fluid ounces per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in the Couraze 4 Insecticide solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Green June beetle, Japanese beetle, Leafhoppers/Sharpshooters, Plant bugs, Rose chafer, San Jose scale	1.6 – 3.2
Cherry Fruit fly	2.4 – 3.2
Pests Suppressed	
Plum curculio, Stink bugs	3.2
Restrictions for Apricot, Nectarine, Peach: <ul style="list-style-type: none"> Pre-Harvest Interval (PHI): 0 day Minimum interval between applications: 7 days Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 9.6 fluid ounces/Acre (0.3lb AI/Acre) Minimum application volume (water): 50 GPA - ground application, 25 GPA- aerial application. Do not apply pre-bloom or during bloom or when bees are actively foraging. Restrictions for Cherries, Plums, Plumcot, Prune: <ul style="list-style-type: none"> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum Couraze 4 Insecticide allowed per season when making foliar applications: 16.0 fluid ounces/Acre (0.5lb AI/Acre) Minimum application volume (water): 50 GPA - ground application, 25 GPA- aerial application. Do not apply pre-bloom or during bloom or when bees are actively foraging. Applications: <p>Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (www.epa.gov) for latest crop groups.</p>	

TREE NUTS¹ - Crops of Crop Group 14: Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Mealybugs, Spittlebugs, Termites, Whiteflies	8.0 – 16.0
Pests/Diseases Suppressed	
Pecan scab (from reduction in honeydew deposition)	16.0
Thrips (foliage-feeding thrips only)	16.0
Restrictions: <p>Pre-Harvest Interval (PHI): 7 days</p> <ul style="list-style-type: none"> Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. Do not apply to Almonds. Applications: Apply specified dosage prior to or at onset of pest infestation in one of the following methods: <ul style="list-style-type: none"> Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent irrigation equipment. Pre-wet soil prior to applications of Couraze 4 Insecticide and allow soil to dry following application and prior to subsequent irrigation. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site. Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Product should be applied in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area should follow within 48 hours to promote uptake by root system. For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18-24 inches to obtain optimum control. Allow soil to dry following treatment and prior to applying any irrigation. Remarks: Use the higher listed rates when applied by shank or subsurface sidedress, used on larger trees, soils with high clay content, for high plant populations, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.	

¹Not permitted for use in California unless otherwise directed by Supplemental label.

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids (except black pecan aphid), Leafhoppers/Sharpshooters, <i>Phylloxera</i> spp. (leaf infestations), Spittlebugs, Whiteflies	1.4 – 2.8
Black pecan aphid, Mealybugs, San Jose scale	3.2
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 6 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 11.5 fluid ounces/Acre (0.36lb AI/Acre) • Minimum application volume (water): 50 GPA-ground application, 25 GPA- aerial application • Do not apply pre-bloom or during bloom or when bees are actively foraging. Applications: Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation two applications on a 10 to 14 day interval may be required to achieve control. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (www.epa.gov) for latest crop groups.	

¹Not permitted for use in California unless otherwise directed by Supplemental label.

TROPICAL FRUIT- Acerola, Atemoya¹, Avocado, Birida¹, Black sapote, Canistel, Cherimoya¹, Custard apple¹, Feijoa, Jaboticaba, Guava, Llama¹, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop¹, Spanish lime, Star apple, Starfruit, Sugar apple¹, Wax jambu

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Avocado lacebug, Leafhoppers, Whiteflies	12.0 – 16.0
Pests Suppressed	
Scales, Thrips (foliage feeding thrips only)	16.0
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 6 days • Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/A). • Do not apply pre-bloom or during bloom or when bees are actively foraging. Applications: Apply specified dosage in the following method: <ul style="list-style-type: none"> • Chemigation through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. Not permitted for use in California unless otherwise directed by Supplemental label.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Mealybugs, Thrips (foliage feeding thrips only), Whiteflies	3.2
Pests Suppressed	
Scales	3.2
Restrictions: <ul style="list-style-type: none"> • Pre-Harvest Interval (PHI): 7 days • Minimum interval between applications: 10 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16.0 fluid ounces per acre (0.5 lb AI/A). • Do not apply pre-bloom or during bloom or when bees are actively foraging. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or from improved control of other pests. Aerial application of Couraze 4 Insecticide may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines.	

Not permitted for use in California unless otherwise directed by Supplemental label.

OTHER CROPS

CHRISTMAS TREE¹

Soil Application

Pests Controlled	Rate: Fluid ounces per acre
White grub complex (damage from grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> Maximum Couraze 4 Insecticide allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre) Applications: Soil incorporation and movement of Couraze 4 Insecticide to the root zone is required for activity. Couraze 4 Insecticide can be incorporated most readily when applied to moist soil. Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> Chemigation into root zone through low pressure drip, trickle, micro-sprinkler, or equivalent equipment. 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25-1 inch of irrigation within 12 hours after application. For optimal grub control, apply Couraze 4 Insecticide during adult flight activity, or up to mid-July, when 1 st instar larvae are present. ¹ Not permitted for use in California unless otherwise directed by Supplemental label.	

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Adelgids, Sawflies	1.6 – 3.2
Restrictions: <ul style="list-style-type: none"> Minimum interval between applications: 7 days Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre) Applications: Gall-forming adelgids - time applications to coincide with full bud-swell of earliest bud-breaking trees. Once galls form spraying will be ineffective. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.	

POPLAR/COTTONWOOD¹ – (includes members of the genus *Populus* grown for pulp or timber)

Soil Application

Field Application Rates. See details below for Cuttings/Whips Application rates.	
Pests Controlled	Rate: Fluid ounces per acre
Aphids, Cottonwood leaf beetle	8.0 – 16.0
Pest Suppressed	
<i>Phylloxera popularia</i>	8.0 – 16.0
Restrictions: <ul style="list-style-type: none"> Maximum Couraze 4 Insecticide allowed at-plant per crop season: 16.0 fluid ounces/Acre (0.5 lb AI/A). Do not apply pre-bloom or during bloom or when bees are actively foraging. Applications: Apply specified dosage in the following method: <ul style="list-style-type: none"> Chemigation through low-pressure drip irrigation. For narrow row, cutting orchards/nurseries used for plant propagation, shank into root zone followed by adequate irrigation to promote uptake. (Adequate irrigation depends on soil moisture level at application. Under dry conditions, apply 0.25 inches/Acre). For Cottonwood leaf beetle, protection against damage will occur when application is made early season, when the beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake. For <i>Phylloxera</i> , apply early in the year from break of dormancy through May.	

Cutting/Whip Application Rates. See details above for Field Application Rates.	
Pests Controlled	Cutting/Whip Soaking Solution fluid ounces Couraze 4 Insecticide Needed per 100 gallons
Cottonwood leaf beetle	6.65 – 13.3 (unhydrated cuttings/whips) 13.3 – 20.0 (partially hydrated cuttings/whips)
Pest Suppressed	
Aphids, <i>Phylloxera popularia</i>	13.3 (unhydrated cuttings/whips) 20.0 (partially hydrated cuttings/whips)
Restrictions:	

- Maximum Couraze 4 Insecticide allowed at-plant per crop season: 16.0 fluid ounces/Acre (0.5 lb AI/A).

Applications: Moisture content of cuttings/whips prior to application, the solution concentration, and the length of soaking interval interact to affect the amount of product absorbed into plant material. For a constant soaking interval of 24 hours, drier cuttings/whips absorb a higher quantity of solution and require a lower concentration. Conversely, more hydrated cuttings/whips absorb less solution and require a higher concentration. Soaking of cuttings/whips should occur in a covered container in absence of UV light. Not all *Populus* spp. clones/varieties/hybrids have been tested for crop safety. Without specific knowledge about a particular *Populus* spp. clone/variety/hybrid, treat and evaluate small numbers of cuttings/whips prior to commercial use.

Apply Couraze 4 Insecticide in one of the following cuttings/whips soaking methods:

- For freshly cut (unhydrated) cuttings/whips, soak plant material in specified solution concentration for 24 hours prior to cold storage. After removal from cold storage, plant as needed.
- For previously hydrated cuttings/whips removed from cold storage, allow plant material to reach room temperature and soak in specified solution concentration for 24 hours prior to planting.

Proper care must be taken in disposal of any residual soaking solution. Solution may be applied to existing trees or other registered crops as long as all product label precautions and restrictions are observed.

Use not permitted in California unless otherwise directed by supplemental labeling.

Foliar Application

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leaf beetles	1.6 – 3.2
Restrictions: <ul style="list-style-type: none"> • Minimum interval between applications: 10 days • Maximum Couraze 4 Insecticide allowed per crop season when making foliar applications: 16 fluid ounces per acre (0.51b AI per acre) • Do not apply pre-bloom or during bloom or when bees are actively foraging. <p>Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Couraze 4 Insecticide may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Couraze 4 Insecticide may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.</p> <p>Use not permitted in California unless otherwise directed by supplemental labeling.</p>	

COMMERCIAL POULTRY FACILITIES

POULTRY HOUSING STRUCTURES

Pests Controlled	Rate: Fluid ounces per acre
Darkling beetles, Hide beetles (Dermestids)	3.0 (90 ml)
Restrictions: <ul style="list-style-type: none"> • Do not apply when poultry are present or within 7 days of poultry placement. • Do not allow food or feed to be contacted by spray mixture. • Remove all feed and water from the treatment area before applying spray mixture. • Keep people and pets out of treated areas until sprays have dried. <p>Applications: Apply between flocks. De-caking and sanitation procedures must be completed prior to application. Apply as a spot, crack and crevice, or surface spray on floors, walls, and support beams of poultry structure. Use a minimum of ½ to 2 gallons of spray mixture per 1000 square feet. Prepare spray mixture by filling the spray tank with ½ the water required, then add the specified volume of product. Add the remaining water with agitating or mixing. Maintain constant agitation while applying.</p> <p>Apply spray mixture to the entire footing including 1 foot up the wall above the footing, and in 3 to 4 foot wide bands directly under all feed lines. Areas under feed lines typically harbor large numbers of adults and larvae of either of these pests when an infestation occurs. Pre-measure these areas to determine the amount of spray mixture to apply. For poultry structures prone to large infestations, treat all footings, including 1 foot up the wall and the entire floor area.</p> <p>Cracks and crevices are prone to large infestations of these pests. Apply as a crack and crevice treatment around wall insulation or other areas that may harbor them. If structures have support beams, treat the floor with a 1 foot band around each beam and apply 2 feet up the beam.</p> <p>For poultry structures prone to extreme infestation, treat the entire structure with a broadcast application. Apply 3.0 fluid ounces in 2 gallons of water per 1000 square feet of surface. Apply broadcast spray to areas where litter has accumulated (floor, underneath feed and water lines, lower sections of walls, and corners).</p> <p>Resistance Management, Mode of Action Rotation: To avoid problems with pest resistance to imidacloprid, rotate to another insecticide which provides a different mode of action every 2-3 flocks. Rotate between 3 different insecticide mode of action classes labeled for control of these pests over the course of a calendar year.</p>	
Pests Controlled	Rate: Fluid ounces per acre
Nuisance ants	0.125 – 0.25 (3/4 – 1.5 TSP)
Restrictions:	

- Do not use for control of fire ants (native or imported), carpenter ants, harvester ants or pharaoh ants.
- Keep people and pets out of all treated areas until sprays have dried.

Applications: Apply as a crack and crevice or wall void treatment inside poultry structures. Apply to all cracks, crevices, drilled holes, onto walls, around structure entry points such as doors, windows, vents, eaves, soffits, and utility access openings. If nests are present in voids, apply into the void wherever possible. Apply evenly to all treatment surfaces, but not to the point of runoff. Apply exterior areas around the structure where ants may be present (soil, turf, ornamental shrubs and plantings, and groundcover in close to or touching the structure). For above-ground nests, such as those found in wooden posts, decks, fences, or trees, spray into openings where ants are traveling and on the wood surface.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment. Open dumping is prohibited.

PESTICIDE STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Do not store diluted spray.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable containers 5 gallons or less:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable containers 5 gallons or larger:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Repeat this procedure two more times.

WARRANTY DISCLAIMER

Cheminova warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CHEMINOVA MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions cited on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Cheminova or the Seller. All such risks shall be assumed by Buyer and User. Buyer and User agree to hold Cheminova and the Seller harmless for any claims related to such factors.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to one of the following, at Cheminova's election:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, Cheminova shall not be liable for consequential, incidental, or special damages or losses in any matter.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Cheminova or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Couraze® 4 is a registered trademark of Cheminova, Inc.
Admire, Calypso, Gaucho, Leverage, Provado and Trimax are registered trademarks of Bayer CropSciences
Actara, Centric, Cruiser and Platinum are registered trademarks of a Syngenta Group Company.
Assail and Intruder are registered trademarks of Nippon Soda Company, LTD.
Belay and Clutch are registered trademarks of Arysta LifeScience
Venom is a trademark of Valent USA Corporation
Advise and Gallent are trademarks of Agrilance, LLC
Impulse and Macho are trademarks of Albaugh, Inc.
Widow is a trademark of Loveland Products, Inc.
Nuprid is a trademark of NuFarm America, Inc.
Pasada and Alias are trademarks of Makhteshim Agan of North America, Inc.

Cheminova, Inc.
P. O. Box 110566
One Park Drive
Research Triangle Park, NC 27709
www.cheminova.us.com