



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

67760-129

Date of Issuance:

10/22/2014

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Azoxystrobin 296 g/L + Flutriafol
218 g/L SC

Name and Address of Registrant (include ZIP Code):

Paula Bodey
Cheminova, Inc.
P.O. Box 110566
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Date:

10/22/14

2. You are required to comply with the DCI identified below:

a. Azoxystrobin GDCI-128810-892, issued on 11/09/2011

You must comply with all of the data requirements within the deadlines established by the order. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: http://www.epa.gov/oppsrrd1/contacts_prd.htm

3. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, “EPA Reg. No. 67760-129.”

4. Submit one copy of the final printed label for the record before you release the product for shipment.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:


- Basic CSF dated 11/27/2013
- Alternate CSF #1 dated 10/02/2014

If you have any questions, please contact Shaja Joyner at (703) 308-3194 or joyner.shaja@epa.gov.

Shaja B. Joyner, Product Manager 20
Fungicide and Herbicide Branch
Registration Division (7505P)

Enclosures:

Acute Toxicity Review for EPA File Symbol 67760-REO dated 8/28/14 {DP417376}
End Use Product Chemistry Review dated 8/27/2014 {DP417375}



Group 1 & 3 Fungicide

Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide

Alternate brand name: TOPGUARD® EQ Fungicide]
[When alternate brand name is used, it will also be used in the body of the label.]

ACTIVE INGREDIENTS:	By Weight
Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate*	25.3%
Flutriafol	18.63%
OTHER INGREDIENTS:	<u>56.07%</u>
TOTAL:	100.00%

Contains 1.82 lb. of flutriafol and 2.47 lb. of azoxystrobin active ingredient per gallon.
Suspension Concentrate.
*IUPAC

Keep Out Of Reach of Children
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

**IN CASE OF A MEDICAL EMERGENCY INVOLVING THIS PRODUCT,
CALL TOLL FREE, DAY OR NIGHT, 1-866-303-6950**

– OR –

For MEDICAL EMERGENCY 866-303-6950	For SPILLS CHEMTREC 800-424-9300
---	---

- Note to PM**, the following bracketed statements are individually optional depending on the packaging configuration and whether a booklet label design is used:
- a. [See First Aid statement on back panel of booklet.]
 - b. [See First Aid statement on back panel.]
 - c. [See additional precautionary statements and Directions for Use in booklet.]
 - d. [Read the entire label before using this product. See First Aid, Precautionary Statements, Directions for Use on individual packages.]

Read the entire label before using this product. Use only according to label instructions. Read the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES before buying or using. If terms are unacceptable, return product unopened without delay.

EPA Reg. No. 67760-XXX

EPA Est. No. _____

Net Contents: _____ gal.

Manufactured For:
CHEMINOVA INC.
PO Box 110566
Research Triangle Park, NC 27709
1-800-548-6113

TOPGUARD is a registered trademark of Cheminova.
[Optional marketing language] [Cheminova] [-] [Helping you grow.]

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if absorbed through skin. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently skin contact may cause allergic reactions in some individuals.

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

If on skin:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-303-6950 for emergency medical treatment information.

Note to Physician: No specific antidote. Treat symptomatically.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators must wear: long-sleeved shirt and long pants, and shoes plus socks.

Mixers and Loaders must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves (such as Natural Rubber, Selection Category A), and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, mammals, and aquatic invertebrates. Azoxystrobin and flutriafol can be persistent for several months or longer. For terrestrial uses: Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and run off may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Azoxystrobin, a degradate of azoxystrobin, and flutriafol are known to leach through soil to groundwater under certain conditions as a result of label use. These chemicals may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Run off of this product will be reduced by avoiding applications when rainfall or irrigation is expected within 48 hours.

Notify State and/or Federal authorities and Cheminova immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONARY STATEMENTS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY AND/OR ILLEGAL RESIDUES.

[Optional state restriction language for the state of NY: Not for sale, distribution, or use in Nassau and Suffolk Counties of New York State.]

[Optional state restriction language for the state of PA: Use of this product through airblast application equipment on grapes is prohibited in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, MillCreek, Fairview, Girard and Springfield. his prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.]

[Optional language for CA: In California, this product may only be applied to fields that are not irrigated, or are irrigated using pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems.]

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is located in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves, shoes plus socks, and protective eyewear.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store unused product in original container in a cool, dry, secure area.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling:

Nonrefillable containers equal to or less than 5 gallons:

Do not reuse or refill this container. Triple rinse container promptly after emptying. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable containers greater than 5 gallons and less than 260 gallons:

Do not reuse or refill this container. Triple rinse or pressure rinse container promptly after emptying. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Triple rinse or Pressure rinse as follows:]

Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times, or use water pressurized to at least 40 PSI to rinse all interior portions. Empty the rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and

continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Note to PM, IBC's may be nonrefillable or refillable, the container handling section will state either Nonrefillable or Refillable.]

Nonrefillable container. {Or, } Refillable container.

Bottom discharge IBC (Intermediate Bulk Container): [Only for refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the re-filler.] Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or longer until rinsate becomes clear. Replace the lid and close bottom valve.

PRODUCT DIRECTIONS

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Product Restrictions:

Chemigation: Do not apply this product through any type of irrigation.

Greenhouses: Do not use for disease control in food crops grown in greenhouses.

Crop Rotation Interval: Treated areas may be rotated to the following crops based on plant back intervals outlined in the table below. Rotation to any other crop not listed in the table is not allowed.

Crop	Plant Back Interval
Corn (field, field corn grown for seed, and popcorn), peanuts, soybeans, sugar beets, grapes, pome fruit, and stone fruit.	0 Days
Sweet corn, cotton	180 Days

Compatibility Jar Test:

Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide is compatible with most products, however not all have been tested. Use the following compatibility test to ensure physical compatibility.

It is recommended before adding Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide along with other additives or pesticide products to a spray tank that a compatibility jar test is conducted.

Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add wettable powders and water dispersible granular products first, next liquid flowables, then emulsifiable concentrates, and last liquid soluble products. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. Use tank mix combinations on a small number of plants before treating larger areas. When tank mixing, follow more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on

Use of Adjuvants: An adjuvant may be used with Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide unless otherwise specified in the crop use directions. Please note that some phytotoxic effects have been demonstrated when tank mixed with adjuvants that contain some form of silicone.

Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide is compatible with most products; however, not all have been tested. Use the compatibility jar test to ensure physical compatibility.

Tank Mixing: Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide has demonstrated some phytotoxic effects when tank mixed with emulsifiable concentrate (EC) products. These effects are enhanced if applications are made under cool, cloudy conditions that exist for several days following application.

Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide may be tank mixed with most fungicides, herbicides, insecticides, and/or other additives unless prohibited on the label of the tank mix partner. Follow more restrictive labeling of any tank mix partner. Although Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide is compatible with most products, not all combinations have been tested. Use the compatibility jar test to ensure physical compatibility. Before applying any tank mixture not specifically recommended on this label, the crop safety of the target crop should be confirmed by applying the mixture to a small area of the target crop in accordance to the label instructions.

Resistance Management: Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide is a mixture that consists of two modes of fungicide action, a Group 3 (flutriafol) and a Group 11 (azoxystrobin). Repeated use of the same group of fungicides for a targeted disease may lead to the selection of resistant strains of fungi and result in reduced disease control. Utilizing premixes such as Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide and/or rotating to fungicides with different modes of action are reliable disease resistance management practices.

For assistance on a particular crop and disease control situation, consult your local agricultural dealer, consultant, applicator, or state extension personnel for specific practices or recommendations in your area. Cheminova encourages proper use rates, timing, and alternation with other effective fungicides to ensure long-term control of the fungal diseases on this label.

Water Volumes: Use the following spray volume guidelines, unless otherwise specified in the specific Use Directions on this label:

For ground application, use a minimum of 10 gallons of spray solution per acre by ground sprayer.

For aerial application use a minimum of 10 gallons of spray solution per acre for tree and vine crops, and a minimum of 2 gallons of spray solution for row crops. Higher spray volumes will result in better coverage and thus improved disease control.

High Disease Pressure: Where a product rate range is listed, use the higher rate of Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide and/or reduced spray interval when disease pressure is high and/or conditions are favorable for disease development.

Phytotoxicity to Apples: Any product containing azoxystrobin (including Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide) is extremely phytotoxic to certain apple and crabapple varieties. Extreme caution must be used to prevent injury to apple trees (and apple fruit) from spray drift. AVOID SPRAY DRIFT that may reach apple trees. See Spray Drift Management section in this label for ways to reduce spray drift or contact your State extension agent for spray drift prevention guidelines in your area.

Even trace amounts of azoxystrobin may cause phytotoxicity to certain apple and crabapple varieties. DO NOT use spray equipment which has been previously used to apply azoxystrobin to spray apple trees.

THE APPLICATOR AND GROWER ARE RESPONSIBLE FOR SPRAY DRIFT MANAGEMENT.

SPRAY DRIFT MANAGEMENT

To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of equipment and weather related factors determine the potential for spray drift.

DO NOT spray when conditions favor drift beyond the area intended for application. Conditions that may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

APPLICATION METHODS AND INFORMATION

Spray Equipment/Volume:

Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide can be applied with equipment commonly used for ground or aerial application of pesticides. Choice of method must be in accordance with this label.

Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide must be applied in sufficient water volume to ensure thorough coverage and penetration for good disease control. Avoid overlap of spray solution as crop injury may occur. (Please see Water Volumes section and specific crop use directions.)

Mixing Order instructions:

1. Equipment must be clean before preparing spray solution.
2. Fill spray tank with clean water to reach $\frac{3}{4}$ of the final spray solution required.
3. Do not prepare more spray solution than required for immediate operation.
4. Begin agitation. Continue agitation during mixing and application of the spray solution.
5. Add tank mix products in the following order:
 - a. Products in PVA bags. If using a product that is in PVA bags, add this to the spray tank first, ensuring the bags are completely dissolved before adding the next product.
 - b. Water dispersible products (such as Azoxystrobin 296 g/L & Flutriafol 218 g/L SC Fungicide, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) should be added to the tank next.
 - c. Water soluble products.
 - d. Emulsifiable concentrates (such as oil concentrates, when applicable).
 - e. Water soluble additives (such as ammonium sulfate or urea ammonium nitrate, when applicable)
6. Add remaining water to fill the tank to 100% of spray solution required.

Ensure that each product added to the spray tank is thoroughly mixed and suspended prior to adding the next product. Thoroughly clean spray tank after each day's use and dispose of pesticide rinsate by application to an already treated area.

RATE EQUIVALENCY TABLE

Product Rate (fl.oz./acre)	Equivalent Flutriafol Active Ingredient Rate (lb.AI/acre)	Equivalent Azoxystrobin Active Ingredient Rate (lb.AI/acre)
4.0	0.057	0.077
5.0	0.071	0.096
6.0	0.085	0.116
7.0	0.10	0.135
8.0	0.114	0.154

Use Directions for Cherry (Sweet and Tart) and Plum

Target Diseases	Product Rate (fl oz/ac)	Application Directions
Alternaria Spot and Fruit Rot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum prunicola</i>) Brown Rot Blossom Blight and Fruit Rot (<i>Monilinia spp.</i>) Cherry Leaf Spot (<i>Blumeriella jaapii</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Powdery Mildew (<i>Sphaerotheca pannosa, Podosphaera clandestina</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	6.0 – 8.0	For Brown Rot Blossom Blight, start applications at early bloom and continue through petal fall. For Fruit Rot apply one to two weeks prior to harvest. For Cherry Leaf Spot start applications at petal fall. Repeat as necessary if disease conditions are favorable for disease development. For all other diseases apply preventatively or when conditions are favorable for disease development and continue on a 7 – 10 day application interval. For multiple applications refer to the guidelines under Resistance Management.
Restrictions: <ul style="list-style-type: none"> Do not exceed 8 fl.oz/A for a single application. Do not apply more than 4 applications of product per year. Do not apply more than 32 fl oz/ac of product per acre per year Do not apply more than 0.456 lb a.i. of flutriafol per acre per year Do not apply more than 1.5 lb a.i. of azoxystrobin per acre per year Restricted Entry Interval (REI): The REI is 12 hours Preharvest Interval: Do not apply within 7 days of harvest 		

Use Directions for Grapes [Optional language: (fresh, table, wine, raisin)]

Target Diseases	Product Rate (fl oz/ac)	Application Directions
Black Rot (<i>Guignardia bidwellii</i>) Phomopsis Cane and Leaf Spot (<i>Phomopsis viticola</i>) Powdery Mildew (<i>Uncinula necator</i>) Suppression Only: Botrytis Bunch Rot (<i>Botrytis cinerea</i>)	5.0 – 6.0	For powdery mildew, begin application at 6 to 10 inch shoots. For all other diseases, apply preventatively or when conditions are favorable for disease development. Repeat on 14 – 21 day intervals throughout the season. For multiple applications refer to the guidelines under Resistance Management. An adjuvant may be added at recommended rates. Azoxystrobin is extremely phytotoxic to certain apple varieties even in trace amounts. Avoid spray drift. Please see Phytotoxicity to Apples section for management guidance.
Downy Mildew (<i>Plasmopara viticola</i>)	8.0	Apply preventatively or when conditions are favorable for disease development. Repeat on 14 – 21 day intervals throughout the season.
Restrictions: <ul style="list-style-type: none"> • No single application should exceed 8 fl.oz/A. • Do not apply more than 6 applications of product per acre per year • Do not apply more than 34 fl oz/ac of product per acre per year • Do not apply more than 0.488 lb a.i. of flutriafol per acre per year • Do not apply more than 1.5 lb a.i. of azoxystrobin per acre per year • Restricted Entry Interval (REI): The REI is 5 days for girdling or turning of grapes. The REI for all other activities is 12 hours. • Preharvest Interval: Do not apply within 14 days of harvest 		

Use Directions for Peanuts

Target Diseases	Product Rate (fl oz/ac)	Application Directions
Soilborne Diseases - mid-late season Rhizoctonia Peg and Pod Rot <i>(Rhizoctonia solani)</i> Stem Rot/White Mold <i>(Sclerotium rolfsii)</i>	8.0	Foliar Application: apply at approximately 60 and 90 days after planting or earlier if disease conditions develop. These applications will provide protection against the soilborne diseases and will also provide control of the foliar diseases listed for a 14 day period after each spray. An adjuvant may be added at recommended rates.
Foliar Diseases Early Leaf Spot <i>(Cercospora arachidicola)</i> Late Leaf Spot <i>(Cercosporidium personatum)</i> Leaf Scorch/ Pepper Spot <i>(Leptosphaerulina crassiasca)</i> Rust <i>(Puccinia arachidis)</i> Web Blotch <i>(Phoma arachidicola)</i>	5.0 – 7.0	Apply product beginning 35 to 40 days after planting or at the first appearance of disease. A lower rate may be applied for control of foliar diseases on a 14 day interval. Additional applications of other fungicides on a leaf spot application schedule will be required to provide season-long disease control of the leaf spot diseases. For multiple applications refer to the guidelines under Resistance Management.
Suppression of Soil Borne Diseases: White Mold / Southern Stem Rot <i>(Sclerotium rolfsii)</i> Rhizoctonia Peg and Pod Rot <i>(Rhizoctonia solani)</i>	5.0 – 7.0	Foliar Application: apply at approximately 60 and 90 days after planting or earlier if disease conditions develop. These applications will provide protection against the soilborne diseases and will also provide control of the foliar diseases listed for a 14 day period after each spray. An adjuvant may be added at recommended rates.
Restrictions: <ul style="list-style-type: none"> • Do not apply more than 32 fl oz/ac of product per acre per year • Do not apply more than 0.455 lb a.i. of flutriafol per acre per year • Do not apply more than 0.8 lb a.i. of azoxystrobin per acre per year • Do not apply more than 4 applications of product per year.. • No single application should exceed 8 fl.oz/A. • Restricted Entry Interval (REI): The REI is 12 hours • Preharvest Interval: Do not apply within 14 days of harvest 		

Use Directions for Soybeans

Target Diseases	Product Rate (fl oz/ac)	Application Directions
<p>Aerial Blight (<i>Rhizoctonia solani</i>) Alternaria Leaf Spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum truncatum</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora Blight and Purple Seed Stain (<i>Cercospora kikuchii</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Pod and Stem Blight (<i>Diaporthe phaseolorum</i>) Powdery Mildew (<i>Microsphaera diffusa</i>) Target Spot (<i>Corynespora cassicola</i>) Rust (<i>Phakopsora pachyrhizi</i>)</p> <p>Suppression: White Mold / Stem Rot (<i>Sclerotinia sclerotiorum</i>) Southern Blight (<i>Sclerotium rolfsii</i>)</p>	<p>5.0 – 8.0</p>	<p>For Cercospora Leaf Blight begin applications at first flower (R1)</p> <p>For Rust apply beginning at first visible flower (R1) or when conditions are favorable for disease development.</p> <p>For Aerial Blight use 8.0 fl oz/ac.</p> <p>For all other diseases apply preventatively or when conditions are favorable for disease development. Use the higher rate or additional application when environmental conditions are conducive for disease development.</p> <p>For multiple applications refer to the guidelines under Resistance Management.</p> <p>An adjuvant may be added at recommended rates. Use of a crop oil concentrate or non-ionic surfactant with the lower use rate is recommended.</p> <p>Minimum retreatment interval: For soybean rust, the minimum retreatment interval is 21 days, maximum retreatment interval is 35 days. All other diseases: the minimum retreatment interval is 14 days, maximum retreatment interval is 35 days</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> • Do not feed forage or hay to animals or permit animals to graze • Do not apply more than 15.5 fl oz/ac of product per acre per year • Only one application of product at 8 fl oz/ac may be made to any one field during a single growing season. • Do not apply more than 3 applications per year. • Do not apply more than 0.23 lb a.i. flutriafol per acre per year. • Do not apply more than 1.5 lb a.i. of azoxystrobin per acre per year. • Restricted Entry Interval (REI): The REI is 12 hours • Preharvest Interval: Do not apply within 21 days of harvest 		

Use Directions for Sugar Beets

Target Diseases	Product Rate (fl oz/ac)	Application Directions
Foliar Diseases Alternaria Leaf Spot <i>(Alternaria alternata)</i> Cercospora Leaf Spot <i>(Cercospora beticola)</i> PowderyMildew <i>(Erysiphe betae)</i>	5.0 – 7.0	Apply preventatively or when conditions are favorable for disease development and continue throughout the season every 14 days. For multiple applications refer to the guidelines under Resistance Management. An adjuvant may be added at recommended rates.
Restrictions: <ul style="list-style-type: none"> • Do not apply more than 2 applications of product per year. • No single application may exceed 7 fl.oz/A. • Do not apply more than 16 fl oz/ac of product per acre per year • Do not apply more than 0.23 lb a.i./ac of flutriafol per acre per year • Do not apply more than 2.0 lb a.i. of azoxystrobin per acre per year • Restricted Entry Interval (REI): The REI is 12 hours • Preharvest Interval: Do not harvest within 21 days of harvest. 		

WARRANTY DISCLAIMER

Cheminova warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CHEMINOVA MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Cheminova or the Seller. All such risks shall be assumed by Buyer and User. Buyer and User agree to hold Cheminova and the Seller harmless for any claims related to such factors.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to one of the following, at Cheminova's election:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, Cheminova shall not be liable for consequential, incidental, or special damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Cheminova or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

TOPGUARD is a registered trademark of Cheminova.