

67959-4

9/27/2007

1/29



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

SEP 27 2007

Mr. Miriam Eichelberger  
Office Manager/Registrations  
10260 Westheimer Road, Suite 230  
Houston, Texas 77042

Re: Update label in the State of California to Match the one registered in Washington  
EPA Reg. No.: 67959-4  
Date of Submission: August 7, 2007

Dear Mr. Eichelberger:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated August 7, 2007, for the product Trilin Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,  
*Theresa Leane for*  
Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number Trilin Herbicide	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tri Corporation	PM#	
5. Name and Address of Applicant (Include ZIP Code) 10260 Westheimer Rd., Suite 230 Houston, Texas 77042  <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. 67959-4  Product Name Trilin Herbicide	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

We are updating our label in the State of California to match the one registered with EPA in Washington. In the meantime, we have added the following statement in our label to assure ourselves that our customers in the bordering States to California do not enter our product into California.

This notification is consistent with the provision of PR Notice 98-10 and EPA regulations.

Section - III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> Text <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Miriam Eichelberger	Title Office Manager/Registrations	Telephone No. (Include Area Code) 713-785-0053
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received <b>(Stamped)</b>
2. Signature 	3. Title Office Manager/Registrations	
4. Typed Name Miriam Eichelberger	5. Date 8/6/2007	

C  
Notif  
3/29

# TRI CORPORATION

August 7, 2007

Ms. Joanne I. Miller  
Production Manager (23)  
Herbicide Branch  
Registration Division (7505C)  
US Environmental Protection Agency  
1801 S. Bell Street  
Arlington, VA 22202

Re: Trilin Herbicide  
EPA Registration No. 67959-4

Dear Ms. Miller,

We are in the process of updating our label for Trilin Herbicide in the State of California to match the one we have registered with EPA in Washington. Because this process will take a few months we have added a line in our labels reading: "For Sale and Use in States other than California". This way, we assure ourselves that our customers in the bordering States do not enter our product into California. Nothing else has been changed in the label.

This sentence will be taken out as soon as we obtain the approval of our label in California.

Enclosed is a specimen of how our label reads for your files. I would appreciate if you can send us your acceptance with a copy of this letter.

If you have any questions, do not hesitate in calling me.

Thank you!

Sincerely yours,

  
Miriam Eichelberger  
Office Manager/Registrations

Enclosures

# TriCorp Trilin<sup>®</sup>

HERBICIDE

## Herbicide for the Preemergence Control of Annual Grasses and Broadleaf Weeds

<b>ACTIVE INGREDIENT</b>	Trifluralin (2,6-dinitro-N,N-dipropyl-4-(trifluoromethyl)benzenamine).....	42.8%
<b>INERT INGREDIENTS</b>	.....	57.2%
<b>TOTAL</b>	.....	100.0%

Contains 4 Pounds Trifluralin Per Gallon

Contains Petroleum Distillate

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### FIRST AID

<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOTLINE NUMBER:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call 504-439-3140	
<b>NOTE TO PHYSICIAN:</b> This product may pose an aspiration pneumonia hazard.	
See Label for Additional Precautions and Directions for Use.	

TriCorp  
10260 Westheimer Suite 230  
Houston, Texas 77042

EPA REG. NO. 67959-4

### For Sale and Use In States Other Than California

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

#### CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or viton  $\geq$  14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and

#### INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

#### CONTROLLING DROPLET SIZE

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

#### BOOM LENGTH

For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

#### APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

#### SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

#### WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Applications should be avoided below 2 mph due to variable wind direction and high inversion potential.

**NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

#### TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

#### TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

#### SENSITIVE AREAS

**Field Cultivator:** Field cultivators are defined as 3 to 4 rows spaced at intervals of 7 inch soil is left unturned. Set to cut 3 to 4 inches more. Do not use chisel points.

**Combination Seedbed Conditioner:** Set to operate at a speed of at least 5 mph. The tillage devices combined and used as a 3 rows of field cultivator C- or S- shaped spacing of 6 to 9 inches (staggered) followed by a spike tooth or flexline harrow reel or basket.

**Rolling Cultivator:** Set to cut 2 to 4 inches of 6 to 8 mph. Rolling cultivators are for medium textured soils. The rolling cultivator soils when used in sugarcane.

**Bed Conditioner (Do-All):** Set to cut 2 to speed of 4 to 6 mph. The Do-All is effective medium textured soils only. Only one incorporation. Two passes with the planted culture.

**Mulch Treader (other similar disc-till Treader)** to cut 3 to 4 inches deep and open P.T.O. Driven Equipment (tillers, etc.) incorporation is required. Adjust to incorporate 3 inches of the seedbed using rotors spaced soil. P.T.O. equipment should not be open 4 mph.

Other equipment, including the flexible Melroe, is also recommended but only for is specified in this label. Use other equipment

#### CULTIVATION AFTER

Soil may be shallow cultivated without re- Trilin. Poor weed control may result if treated soil since this may bring untreated

#### CROP RECOMMENDATIONS

All recommendations are given as the broadcast For band applications, decrease the amount of surface treated per acre. Apply when the soil can be worked and is suitable for application. See specific crop recommendations under "APPLICATION" heading where specific recommendations are given. Use the lower rate for coarser soil matter. For soils containing 10% or more clay

#### MIXING AND APPLICATION

##### Trilin In Water

Thoroughly clean sprayer prior to use. Fill clean water and start agitation. Add product sufficient agitation and finish filling the tank

##### Trilin Tank Mixes in Water

For all tank mixes, continuous agitation is recommended. Generally provide the best agitation in sprayer avoid stirring or splashing air into the mixture end of the fill pipe below the surface of the allow the mixture to siphon back into the tank

##### Mixing Order

Fill the tank  $\frac{1}{2}$  to  $\frac{1}{3}$  full with clean water and powders and dry flowables and agitate until add aqueous suspensions (flowables); then ingredients and finish filling the tank. Provide filling and application. If spraying and agitating the tank is empty the materials may settle before continuing the spray application.

20/5

- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine and estuarine fish and aquatic invertebrates including shrimp and oysters. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, or viton ≥ 14 mils

while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

## GENERAL INFORMATION

Trilin is a preemergence herbicide which is incorporated into the soil to provide control of grasses and broadleaf weeds. Trilin controls weeds as they germinate. Trilin will not control established weeds.

## GRASSES AND BROADLEAF WEEDS CONTROLLED BY TRILIN

### Grasses

Annual Bluegrass	<i>Poa annua</i>
Barnyardgrass (Watergrass)	<i>Echinochloa</i> sp.
Brachiaria (Signalgrass)	<i>Brachiaria</i> sp.
Bromegrass	<i>Bromus tectorum</i>
(Cheatgrass)	
(Downy Brome)	
Cheat (chess)	<i>Bromus secalinus</i>
Crabgrass	<i>Digitaria</i> spp.
(Large Crabgrass)	
(Smooth Crabgrass)	
Foxtail	<i>Setaria</i> spp.
(Bottlegrass)	
(Bristlegrass)	
(Giant Foxtail)	
(Green Foxtail)	
(Foxtail Millet)	
(Pigeongrass)	
(Robust Foxtail)	
(Yellow Foxtail)	
Itchgrass (Raoulgrass)	<i>Rottboellia exaltata</i>
(See sugarcane for special instructions)	
Johnsongrass (from seed)	<i>Sorghum halepense</i>
(Rhizome - see special instructions for control in cotton, soybeans, fruit and nut crops, vineyards and cottonwood trees grown for pulpwood)	
Junglerice	<i>Echinochloa colonum</i>
Panicum	
Fall Panicum	<i>Panicum dichotomiflorum</i>
(Spreading Panicgrass - see special instructions in cotton and soybeans)	
Guineagrass	<i>Panicum maximum</i>
(See sugarcane for special instructions)	
Texas Panicum	<i>Panicum texanum</i>
(See special instructions in cotton and soybeans)	
(Buffalograss)	
(Coloradograss)	
Red Rice	<i>Oryza sativa</i>
(See suppression or partial control directions under soybeans)	
Sandbur (Burggrass)	<i>Cenchrus incertus</i>
Sprangletop	<i>Leptochloa filiformis</i>
Stinkgrass (Lovegrass)	<i>Eragrostis ciliaransensis</i>
Wild Can (Shattercane)	<i>Sorghum bicolor</i>

ing... and... filling' and application. If spraying and ag the tank is empty the materials may settle before continuing the spray application, r the bottom of the tank. A sparge agitator purpose. It may be more difficult to resus is to suspend it originally.

Read and carefully follow all label instruct the tank. To help assure good dispersion mixture by premixing dry and flowable to the slurry through a 20 or 35 mesh wet s finer than 50 mesh line screens in the tan If material builds up on the walls of the : soapy water between fillings. Rinse and c After completion of spraying, thoroughly cl The importance of accurate calibration an as the spray volume decreases. Check calibration and uniform application. To in: apply Trilin when the wind can cause driftir control may result if Trilin is applied to soi to prolonged periods of flooding.

### Ground Application

Using a low pressure herbicide sprayer uniformly, apply Trilin in 5 to 40 gallons of (broadcast spray).

### Aerial Application

For aerial spraying apply Trilin in 5 to 10 g per acre. Pump pressure, nozzle arrangen be adjusted to provide a uniform applicatic proper application spray widths use swath

### Fall Application (Areas receiving more th

See specific crop for recommendations. applications for all crops for which there instructions and for which Trilin is recor application. Trilin should not be applied in t and direct-seeded tomatoes. Apply an between October 15 and December 31. Le winter. On bedded ground, beds should height prior to planting, moving some trea Where soil is left over winter, care should b soil during spring bedding operations. destroy established weeds. Destroy established in furrows due to uncovering o Trilin should not be applied in the fall to soi prolonged periods of flooding or where rice

## GENERAL PREC AND RESTRIC

### PRECAUTIONS

Observe all precaution and limitations on i in tank mixes and overlays. Under nom applied according to directions, Trilin will n injury or soil residue may result from over a or crop injury may result from uneven incorporation of Trilin. Seeding disease, excessive moisture, high salt concentration seedlings and increase the possibility of de conditions, delayed crop development or r

### Plant back Restrictions

In Arizona, Colorado, Idaho, Montana, N Utah, Washington and Wyoming after a sp beets, red beets or spinach should not be p months after fall application. Soil should be prior to planting sugar beets to prevent the a spring application, sorghum (milo), proso be planted for 14 months after applicatio application of Trilin to avoid crop injury. If

- Chemical-resistant gloves, such as barrier laminate, or viton ≥ 14 mils
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter or allow others to enter until sprays have dried.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Avoid freezing. Store above 40°F. If frozen, poor weed control may result. Do not store near heat or flame.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## GENERAL INSTRUCTIONS AND INFORMATION CHEMIGATION

### General Instructions

Apply **Trilin** only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move or drip irrigation systems. Do not apply **Trilin** through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop may result from non-uniform distribution of treated water:

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label prescribed safety device for public water systems is in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### Specific Instructions for Public Water Systems

Public water system means a system for the distribution to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer, (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically

*Eragrostis amabilis*

Stinkgrass (Lovegrass)

Wild Cane (Shattercane)

(See soybean - Trilin Alone for special instructions)

Woolly Cupgrass

*Eragrostis cilianensis*

*Sorghum bicolor*

*Eriochloa villosa*

### Broadleaf Weeds

Carpetweed

Chickweed

Field Bindweed

(See fruit and nut crops and vineyards for special instructions)

Florida Pusley

(Florida Purslane)

(Mexican Clover)

(Pusley)

Goosefoot

Henbit (Fall application only)

Knotweed

Kochia

(Fireweed)

(Mexican Fireweed)

Lambsquarters

Pigweed

(Carlelessweed)

(Prostrate Pigweed)

(Redfoot)

(Rough Pigweed)

(Spiny Pigweed)

Puncturevine (Western U.S. only)

(Caltrop)

(Goathead)

Purslane

Russian Thistle (Tumbleweed)

Stinging Nettle (Nettle)

*Mollugo verticillata*

*Stellaria media*

*Convolvulus arvensis*

(See fruit and nut crops and vineyards for special instructions)

*Richardia scabra*

*Chenopodium hybridum*

*Lamium amplexicaule*

*Polygonum aviculare*

*Kochia scoparia*

*Chenopodium album*

*Amaranthus spp.*

*Tribulus terrestris*

*Portulaca oleracea*

*Salsola kali*

*Urtica dioica*

Long-term and continued use of trifluralin has resulted in the selection of tolerant populations in certain species of weeds. This situation is limited to a few weeds and is generally geographically specific. Weed species known to have some trifluralin tolerant population are goosegrass, green foxtail (pigeongrass) and Palmer amaranthus (Palmer pigweed). Trilin is not recommended for the control of goosegrass, tolerant green foxtail or Palmer amaranthus. Consult State Agricultural Extension Service or Experiment Station weed specialist for specific recommendations for local weed problems.

## SOIL PREPARATION

Soil surface should be smooth enough to enable the efficient use of sprayer and incorporation equipment to insure a uniform application and incorporation of Trilin. Interference can be caused by ground cover such as crop residue or existing weeds. Soil should be tilled prior to the application of Trilin to allow uniform incorporation into the top 2 to 3 inches of soil. Soil moisture should be such that any large clods will be broken up during incorporation process.

### Crop Residues or Existing Weeds

Ground cover such as crop residues of existing weeds can interfere with the incorporation of Trilin into the soil. A manageable level of such ground cover will allow the Trilin to be uniformly incorporated into the top 2 to 3 inches of soil. If the level of the ground cover is such that this cannot be done you must till the soil prior to the application of Trilin.

### Roughness

The soil surface should be smooth enough so that you can operate a sprayer and incorporation equipment efficiently and at speeds which insure a uniform application and incorporation of Trilin.

a spring application, sorghum (milo), or be planted for 14 months after application of Trilin to avoid crop injury not plant any of these crops for 18 months after a fall application of Trilin.

In those areas of Kansas, Nebraska, Dakota and Texas where at least 20 inches was used to produce the crop, sorghum 12 months after an application of Trilin.

Do not plant sorghum, proso millet application of Trilin if less than 20 inches produce the crop. Cool wet weather or growth may increase the possibility of it.

In all other areas receiving greater than moldboard plow before planting sugar be Trilin was made the previous season. Al in the section on control of rhizome j programs.

For vegetables other than those listed planted within 5 months following the ap

## FERTILIZER USE

### LIQUID FERTILIZERS

Trilin may be mixed with most liquid fertilizer with solutions and suspension-type grass control equal to the same rates of recommendations regarding rates and directions, special instructions, and labeling and applications relating to liquid state regulations are the responsibility of selling the fertilizer and chemical mixture.

### Compatibility Testing Tank Mix Partners

A tank mixture of Trilin alone or with aqueous suspensions, flowables liquid properly with some liquid fertilizer material before full-scale mixing to determine needed and which agent does the best surfactants designed for use with liquid following test to select the correct agent.

1. Measure one pint of intended spray water.
2. Check the pH of the liquid and adjust if needed.
3. Add in the given order the intended ingredients in addition.

(a) Surfactants (spreaders), acidifiers, activators; add one teaspoon for each gallon.

(b) Dry ingredients (wetttable powders); add one tablespoon for each pound/100 gallons.

(c) Emulsifiable concentrates; add one teaspoon for each gallon.

(d) Flowables; add one teaspoon for each gallon.

(e) Soluble ingredients; add one tablespoon for each gallon.

(f) Spreader-stickers; one teaspoon for each gallon.

4. The final mixture should be uniform a coagulation occurring. If incompatibility a compatibility agent added first. four ounces/100 gallons. If this does not work, use higher concentrations and other compatibility agents.

5. Allow the mixture to stand undisturbed. If occurs shake and observe the resulting mixture. If redisperses do not attempt to spray the mixture.

(a) More compatibility agents

(b) Different formulations of the active ingredient or emulsifiable concentrates to flow emulsifiable concentrates).

### Mixing Instructions

When mixing Trilin in liquid fertilizers cover prevent the Trilin from drying to the soil.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

#### Specific Instructions for Sprinkler Irrigation Systems

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Good agitation is required in the injection tank.

In moving systems, apply specified dosage of Trilin as a continuous injection. In non-moving systems inject Trilin for 15 to 20 minutes at end of cycle. Use the least amount of water possible for uniform coverage.

Mix the amount of Trilin needed for acreage to be treated into the quantity of water determined during prior calibration (refer to Mixing and Application Directions). For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.

Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all Trilin is flushed from system.

## AERIAL DRIFT REDUCTION ADVISORY INFORMATION

Avoid spray drift at the application site is the responsibility of the applicator. The interactions of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.

the incorporation of Trilin into the soil. A manageable level of such ground cover will allow the Trilin to be uniformly incorporated into the top 2 to 3 inches of soil. If the level of the ground cover is such that this cannot be done you must till the soil prior to the application of Trilin.

#### Roughness

The soil surface should be smooth enough so that you can operate a sprayer and incorporation equipment efficiently and at speeds which insure a uniform application and incorporation of Trilin.

#### General Soil Conditions

To assure incorporation of Trilin, soil moisture conditions should be such that any large clods can be broken up during the incorporation process.

## SOIL TEXTURE GUIDE

The amount of chemical applied will vary with the soil texture and organic matter. A fine textured soil will require more Trilin per acre than a coarse soil. Where rates are based on coarse, medium or fine textured soils, it is understood that soil textural classes are generally categorized as follows:

Coarse	Medium	Fine
Sand	Loam	Sandy Clay
Loamy Sand	Silt	Clay Loam
Sandy Loam	Silt Loam	Silty Clay, Clay
	Silty Clay Loam*	
	Sandy Clay Loam*	

\*Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are mostly sand or silt, they are usually classified as medium textured soils. If they are mostly clay, they are usually classified as fine textured soils.

Do not exceed recommended rates.

## INCORPORATION DIRECTIONS

#### General Directions

Erratic weed control and/or crop injury may result if Trilin is not incorporated into the top 2 to 3 inches of the final seedbed.

#### Before Planting

Initial incorporation of Trilin must take place within 24 hours after application. A second incorporation prior to planting should be done by running the equipment in a different direction from the first. Incorporate Trilin uniformly into the top 2 to 3 inches of the final seedbed.

#### After Planting

For directions concerning incorporation after planting check label for crop or crops of interest for specific instructions.

#### 72 Hour Incorporation Delay - Arizona and California Only

When Trilin is applied as a preplant incorporated treatment, the first incorporation pass must be accomplished within 24 hours after application. In Arizona and California the incorporation delay has been extended from 24 to 72 hours when applied to dry soils. When Trilin is applied to warm soil or if wind velocity is 10 mph or higher, variable weed control may result from delaying the first incorporation beyond 24 hours. Where two incorporation passes are required, the second incorporation may occur anytime prior to planting.

#### Bedded Culture

Trilin needs to be incorporated into the top 2 to 3 inches of the final seedbed for effective weed control.

#### Application Prior To Bedding

Apply Trilin prior to bedding and incorporate with recommended equipment. The bedding operation serves as the second incorporation. Avoid removal of untreated soil from the seedbed before or during the planting operation. This would expose untreated soil allowing weeds to germinate in the drill row.

#### Application After Bedding

Knock off beds to planting height before application and incorporate Trilin with recommended equipment that will conform to the bed shape. Do not

- (b) Different formulations of the active ingredient or emulsifiable concentrates to flow emulsifiable concentrates).

#### Mixing Instructions

When mixing Trilin in liquid fertilizers do not prevent the Trilin from rising to the surface. Use a compatibility agent to make the Trilin emulsifiable concentrates with dry flowal suspensions, flowables liquids or solu compatibility agent is especially important the fertilizer as an oil and the emulsion is combine with the formulation or susceptible to redispersion. A compatibility agent concentrates to form non-oiling mixtures. Compatibility agents can be used at rate of liquid fertilizer and should be mixed with the liquid concentrate. Follow the direct label.

The following is a list of some phosphates to be used with liquid fertilizers. They use agents in tank mixtures in plain water.

- \*Amoco Spray Mate (Amoco Oil Company)
- \*Compat (Farm Chemicals Incorporated)
- \*Kem-Link (Universal Coop., Minneapolis)
- \*Rigo Compatibility Agent (Rigo Company)
- \*Sponto 168D (Witco Chemicals Company)
- \*T-Mulz 734-2 (Thompson-Hayward Chemicals)
- \*Unite (Hopkins Ag Chemicals, Madison)
- \*Not for use in California

#### Application

Use a properly calibrated applicator mixture and apply material uniformly to

#### Incorporation

Follow normal Trilin incorporation procedure

## DRY BULK FE

Trilin may be used for impregnation. Application of dry bulk fertilizers impregnated with Trilin should be equal to the seed. Follow all Trilin label recommendations for approved crops, incorporation, special minimum of 200 pounds per acre of dry Trilin. The recommended rates should be impregnation of any commonly used fertilizer and straight limestone. Trilin materials. Blends containing mixture impregnated.

#### Impregnation

A closed drum belt ribbon or other blender may be used. Trilin should be

#### Rates

Specific crop recommendations for Trilin should be followed. Check the rate table below for Trilin to be impregnated into a ton of dry bulk fertilizer which will be applied per acre

The pesticide injection pipeline must also contain a closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Good agitation is required in the injection tank.

In moving systems, apply specified dosage of Trilin as a continuous injection. In non-moving systems inject Trilin for 15 to 20 minutes at end of cycle. Use the least amount of water possible for uniform coverage.

Mix the amount of Trilin needed for acreage to be treated into the quantity of water determined during prior calibration (refer to Mixing and Application Directions). For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.

Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all Trilin is flushed from system.

## AERIAL DRIFT REDUCTION ADVISORY INFORMATION

Avoid spray drift at the application site is the responsibility of the applicator. The interactions of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Only clay loam soils be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are mostly sand or silt, they are usually classified as medium textured soils. If they are mostly clay, they are usually classified as fine textured soils.

Do not exceed recommended rates.

## INCORPORATION DIRECTIONS

### General Directions

Erratic weed control and/or crop injury may result if Trilin is not incorporated into the top 2 to 3 inches of the final seedbed.

### Before Planting

Initial incorporation of Trilin must take place within 24 hours after application. A second incorporation prior to planting should be done by running the equipment in a different direction from the first. Incorporate Trilin uniformly into the top 2 to 3 inches of the final seedbed.

### After Planting

For directions concerning incorporation after planting check label for crop or crops of interest for specific instructions.

### 72 Hour Incorporation Delay - Arizona and California Only

When Trilin is applied as a preplant incorporated treatment, the first incorporation pass must be accomplished within 24 hours after application. In Arizona and California the incorporation delay has been extended from 24 to 72 hours when applied to dry soils. When Trilin is applied to warm soil or if wind velocity is 10 mph or higher, variable weed control may result from delaying the first incorporation beyond 24 hours. Where two incorporation passes are required, the second incorporation may occur anytime prior to planting.

### Bedded Culture

Trilin needs to be incorporated into the top 2 to 3 inches of the final seedbed for effective weed control.

### Application Prior To Bedding

Apply Trilin prior to bedding and incorporate with recommended equipment. The bedding operation serves as the second incorporation. Avoid removal of untreated soil from the seedbed before or during the planting operation. This would expose untreated soil allowing weeds to germinate in the drill row.

### Application After Bedding

Knock off beds to planting height before application and incorporate Trilin with recommended equipment that will conform to the bed shape. Do not leave untreated soil exposed.

### Recommended Equipment

Two incorporation passes are necessary unless specifically stated otherwise on this label. The second incorporation should not be deeper than the first.

**Disc:** Set disc to cut 4 to 6 inches deep and operate at 4 to 6 mph. A 4 to 6 inch cut should result in Trilin being incorporated into the top 2 to 3 inches of the soil.

Sponto 168D (Witco Chemicals Comp  
\*T-Mulz 734-2 (Thompson-Hayward Ct  
Unite (Hopkins Ag Chemicals, Madiso  
\*Not for use in California

### Application

Use a properly calibrated applicator mixture and apply material uniformly to

### Incorporation

Follow normal Trilin incorporation proc

## DRY BULK FE

Trilin may be used for impregnation Application of dry bulk fertilizers imp weed and grass control equal to the s Follow all Trilin label recommend approved crops, incorporation, specic minimum of 200 pounds per acre of dr the recommended rates should be impregnation of any commonly used nitrate and straight limestone. Trilin materials. Blends containing mixtu impregnated.

### Impregnation

A closed drum belt ribbon or other blender may be used. Trilin should be

### Rates

Specific crop recommendations for t followed. Check the rate table below be impregnated into a ton of dry bu fertilizer which will be applied per acr



**Field Cultivator:** Field cultivators are defined as implements with sweeps of 3 to 4 rows spaced at intervals of 7 inches or less staggered so that no soil is left unturned. Set to cut 3 to 4 inches deep and operate at 5 mph or more. Do not use chisel points.

**Combination Seedbed Conditioner:** Set to cut 3 to 4 inches deep and operate at a speed of at least 5 mph. These are defined as three or more tillage devices combined and used as a single tool. For example, 2 to 3 rows of field cultivator C- or S- shaped shanks with an effective sweep spacing of 6 to 9 inches (staggered so that no soil is left unturned) followed by a spike tooth or flexline harrow followed by a ground driven reel or basket.

**Rolling Cultivator:** Set to cut 2 to 4 inches deep and operate at a speed of 6 to 8 mph. Rolling cultivators are effective for use on coarse and medium textured soils. The rolling cultivator may be used on fine textured soils when used in sugarcane.

**Bed Conditioner (Do-All):** Set to cut 2 to 4 inches deep and operate at a speed of 4 to 6 mph. The Do-All is effective when used on coarse or medium textured soils only. Only one incorporation pass is necessary in bedded culture. Two passes with the Do-All are necessary in flat planted culture.

**Mulch Treader (other similar disc-type implements):** Set Mulch Treader to cut 3 to 4 inches deep and operate at 5 to 8 mph.

**P.T.O. Driven Equipment (tillers, cultivators, hoes):** Only one incorporation is required. Adjust to incorporate Trilin into the top 2 to 3 inches of the seedbed using rotors spaced to give a clean sweep of the soil. P.T.O. equipment should not be operated at a speed greater than 4 mph.

Other equipment, including the flexible tine-tooth harrow (Flexline, Melroe), is also recommended but only for the special usages for which it is specified in this label. Use other equipment only as specified herein.

## CULTIVATION AFTER PLANTING

Soil may be shallow cultivated without reducing the weed control activity of Trilin. Poor weed control may result if cultivation is deeper than the treated soil since this may bring untreated soil to the surface.

## CROP RECOMMENDATIONS

All recommendations are given as the broadcast rates of Trilin per acre. For band applications, decrease the amount of Trilin in proportion to the amount of surface treated per acre. Apply Trilin any time after January 1 when the soil can be worked and is suitable for good incorporation. For fall application, see specific crop recommendations or "FALL APPLICATION" heading where specific crop recommendations are not given. Use the lower rate for coarser soils or soils with lower organic matter. For soils containing 10% or more organic matter do not use Trilin.

## MIXING AND APPLICATION DIRECTIONS

### Trilin In Water

Thoroughly clean sprayer prior to use. Fill the sprayer 1/3 to 1/2 full with clean water and start agitation. Add proper amount of Trilin providing sufficient agitation and finish filling the tank.

### Trilin Tank Mixes in Water

For all tank mixes, continuous agitation is required (Sparger pipe agitators generally provide the best agitation in spray tanks). To prevent foaming avoid stirring or splashing air into the mixture during filling by placing the end of the fill pipe below the surface of the water in the spray tank. Do not allow the mixture to siphon back into the water source.

### Mixing Order

Fill the tank 1/3 to 1/2 full with clean water and start agitation. Add wettable powders and dry flowables and agitate until completely dispersed. Then add aqueous suspensions (flowables); then add Trilin. Lastly, add soluble ingredients and finish filling the tank. Provide continuous agitation during filling and application. If spraying and agitation must be stopped before the tank is empty the materials may settle to the bottom. If this happens, before continuing the spray application, resuspend all of the material in

## Rate Chart for Impregnating Fertilizer With Trilin Added to a Ton of Fertilizer

CHART QUANTITIES LISTED ARE QUARTS OF TRILIN PER TON OF FERTILIZER

Fertilizer Rate Per Acre	TRILIN Rate Per Acre				
	1 pt.	1 1/2 pts.	2 pts.	3 pts.	4 pts.
200 lbs.	5	7.5	10	15	20
250 lbs.	4	6	8	12	16
300 lbs.	3.33	5	6.67	10	13.33
350 lbs.	2.75	4.25	5.75	8.5	11.5
400 lbs.	2.5	3.75	5	7.5	10
450 lbs.	2.25	3.33	4.5	6.67	9

For rates other than those listed above use the following formula to calculate the amount of Trilin to be impregnated on a ton of dry bulk fertilizer

$$\text{Pts. Trilin/Acre} \times \frac{1000 \text{ lbs. Fertilizer/Acre}}{1000} = \frac{\text{Qts. Trilin/Ton of Fertilizer}}{1000}$$

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

### Application

Use a properly calibrated applicator to spread the fertilizer/chemical mixture. The material should be applied uniformly to the soil surface.

### Incorporation

Follow Trilin incorporation procedures.

## APPLICATION INSTRUCTIONS ALFALFA

### Preplant

Trilin may be applied as a preplant incorporated for preemergence control of labeled weeds in direct seeded alfalfa. Apply and incorporate prior to planting according to product label directions.

### Broadcast Application Rates

Soil Texture	Trilin (pt/Acre)
Coarse	1.0
Medium	1.0-1.5*
Fine	1.5

\*Use lower range in areas receiving less than 20 inches of rainfall and irrigation.

Precautions: Some crop stand reduction and stunting may occur with use of Trilin, however, reduced weed competition will allow establishment of a quality stand.

### Mechanically Incorporated

Use a broadcast rate of 1.5 pints per acre on coarse soils and 2 pints on medium and fine soils (in areas receiving less than 20 inches average annual rainfall). Damage to the established alfalfa may be caused if the proper incorporation equipment is not used for thorough soil mixing. Recommended soil preparation application and incorporation instructions should be followed.

### Chemigation or Water Incorporated

Trilin may be surface applied for annual grass control in established alfalfa by chemigation, or ground or aerial broadcast application equipment. Refer to "Chemigation" section of this label for use directions for chemigation.

### Surface Application Activated by Rainfall or Irrigation

Broadcast surface applications of Trilin to established alfalfa may be

## COLE CROP

### Broccoli, Brussels Sprout, Cabbage

#### Transplant

Apply and incorporate Trilin prior to transplant

#### Broadcast Rates Per Acre

	Coarse
Trilin (pts.)	
Areas receiving less than 20" average annual rainfall	1
Areas receiving greater than 20" average annual rainfall	1
For soils with 2-5% organic matter use 1.5 and 2 pints on fine soils. On soils with 5-10% on all soils.	

#### Direct-Seeded

Use Trilin before planting at a broadcast rate and medium soils and 1.5 pints on fine soils matter. Direct-seeded cole crops have ext recommended rates of Trilin.

**Warning:** Stunting or reduced stands may c

## COTTON

Apply Trilin before planting, immediately aft

#### Preemergence

#### Broadcast Rates Per Acre

	Coarse
Trilin (pts.)	
Areas receiving less than 20" average annual rainfall	1
Areas receiving greater than 20" average annual rainfall	1
Use 1.5 pints per acre on coarse and med fine soils with 2-5% organic matter and 2 5-10% organic matter.	

#### Post Plant

Do not disturb the seed when incorporating

#### Layby

Trilin can be applied and incorporated any than 90 days before harvest. Direct layt between the rows and beneath emerged cc as for a preemergence application.

#### Fall Application

Trilin may be applied and incorporated any December 31. The ground may be left fl: Where soil is left flat, take care not to turn up furrows. On bedded ground, knock down b planting, moving some treated soil into the established weeds during preparation o destroy weeds which may have become es uncovering of untreated soil during bedding to wet soil or soils which are subject to prol

#### Application Over Standing or Shredded

Trilin may be broadcast applied, after S standing cotton stalks or after shredc equipment. Trilin must be incorporated application using a disc operated at 4 to 6 n may be delayed until spring prior to bed herbicide or bringing untreated soil to the not be chiseled, ripped or deep plowed folk

Broadcast Rates Per Acre - Fall Applic

the tank is empty the materials may settle to the bottom. If this happens, before continuing the spray application, resuspend all of the material in the bottom of the tank. A sparge agitator is particularly useful for this purpose. It may be more difficult to resuspend the settled material than it is to suspend it originally.

Read and carefully follow all label instructions for each material added to the tank. To help assure good dispersion in the tank water, make a slurry mixture by premixing dry and flowable formulations with water and pour the slurry through a 20 or 35 mesh wet screen in the top of the tank. No finer than 50 mesh line screens in the tank should be used.

If material builds up on the walls of the spray tank, wash the tank with soapy water between fillings. Rinse and continue the spraying operation. After completion of spraying, thoroughly clean the tank lines and screens. The importance of accurate calibration and uniform application increases as the spray volume decreases. Check sprayer daily to insure proper calibration and uniform application. To insure uniform application do not apply Trilin when the wind can cause drifting of spray particles. Poor weed control may result if Trilin is applied to soils which are wet or are subject to prolonged periods of flooding.

#### Ground Application

Using a low pressure herbicide sprayer which will apply the spray uniformly, apply Trilin in 5 to 40 gallons of water per acre (broadcast spray).

#### Aerial Application

For aerial spraying apply Trilin in 5 to 10 gallons of water or liquid fertilizer per acre. Pump pressure, nozzle arrangements and height should be adjusted to provide a uniform application to soil surface. To assure proper application spray widths use swath markers or flagmen.

#### Fall Application (Areas receiving more than 20" average annual rainfall)

See specific crop for recommendations. Use the rates listed for spray applications for all crops for which there are no specific fall application instructions and for which Trilin is recommended as a preemergence application. Trilin should not be applied in the fall for sugarbeets, potatoes and direct-seeded tomatoes. Apply and incorporate Trilin any time between October 15 and December 31. Leave ground flat or bedded over winter. On bedded ground, beds should be knocked down to desired height prior to planting, moving some treated soil from beds into furrows. Where soil is left over winter, care should be taken not to turn up untreated soil during spring bedding operations. During seedbed preparation destroy established weeds. Destroy weeds which have become established in furrows due to uncovering of untreated soil during bedding. Trilin should not be applied in the fall to soils which are wet, are subject to prolonged periods of flooding or where rice was grown the previous year.

## GENERAL PRECAUTIONS AND RESTRICTIONS

#### PRECAUTIONS

Observe all precaution and limitations on the label of each product used in tank mixes and overlays. Under normal growing conditions and if applied according to directions, Trilin will not harm the treated crop. Crop injury or soil residue may result from over application. Erratic weed control or crop injury may result from uneven application or improper soil incorporation of Trilin. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage from Trilin. Under these conditions, delayed crop development or reduced yields may result.

#### Plant back Restrictions

In Arizona, Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming after a spring application of Trilin, sugar beets, red beets or spinach should not be planted for 12 months or for 14 months after fall application. Soil should be plowed to a depth of 12 inches prior to planting sugar beets to prevent the possibility of crop injury. After a spring application, sorghum (milo), proso millet, corn or oats should not be planted for 14 months after application or for 16 months after a fall application of Trilin to avoid crop injury. If land has not been irrigated do not plant any of these crops for 18 months after a spring application or 20

#### Surface Application Activated by Rainfall or Irrigation

Broadcast surface applications of Trilin to established alfalfa may be activated by rainfall, sprinkler, flood or furrow irrigation. Rainfall or a single overhead sprinkler irrigation of 0.5 acre inch or more is required to activate Trilin. If activated by furrow irrigation, care should be taken to thoroughly wet beds between furrows. If rainfall or irrigation has not occurred within 3 days after application, Trilin may be mechanically incorporated. If mechanically incorporated, use equipment that will insure thorough soil mixing with minimum damage to the established alfalfa.

#### Application Timing

Applications to established alfalfa for annual grass control can be made during dormancy or semi-dormancy, or during the growing season immediately after a cutting. Because Trilin does not control established weeds, application must be made prior to the expected time of weed germination. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather. To control these weeds, apply Trilin immediately after a cutting between August 1 and October 1; but prior to weed germination. When fall applied, Trilin controls bromegrass and cheat in addition to other labeled weeds that germinate after application.

Apply Trilin at a broadcast rate of 4 pints per acre on all soil textures by chemigation or surface applied and incorporated by rainfall or irrigation.

#### RESTRICTIONS

Do not harvest forage within 21 days after application and do not harvest hay within 20 days after application. Do not apply more than 8 pints (4 lbs. a.i./A) per year. Apply no more than 4 pints of Trilin during any growing season. In the growing season following application of 4 pints of Trilin to alfalfa, plant only those crops for which Trilin is registered as a preplant treatment or crop injury does not occur.

## ASPARAGUS Established

Follow recommended soil preparation, application and incorporation procedures for Trilin. Trilin can be applied to established asparagus as a single or as a split application. In the winter or early spring, apply to asparagus after ferns are removed but before spear emergence or apply after harvest in the late spring or early summer before turning begins. Trilin will suppress volunteer seedling asparagus and field bindweed used at the following recommended rates and application schedules.

#### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Split Application	1	1.5	2
OR			
Single Application	2	3	4

In any single calendar year, the maximum Trilin to apply is 2 pints per acre on coarse soils, 3 pints on medium soils and 4 pints on fine soils.

## BEAN Dry

Apply Trilin before planting using the following rates:

#### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. For soils with 5-10% organic matter use 2 pints on all soils.

#### TRILIN WITH EPTAM TANK MIX

A tank mix of Trilin and Eptam will effectively control all the following weeds in addition to those weeds listed for Trilin.

Henbit (spring applications) Oat Wild

not be chiseled, ripped or deep plowed follow

#### Broadcast Rates Per Acre - Fall Application

In Alabama, Arkansas, Northern Florida, Georgia, Southwestern Missouri, South Carolina, South Carolina, Tennessee and Texas, apply broadcast rate of 2 pints per acre on all soil. In Arizona, California and Nevada, a broadcast should be used on coarse soil, 2 pints on fine soil.

For other states where cotton may be grown rate of 1 pint per acre on coarse soils, 1.5 pints on fine soils. For coarse soils with 1.5 pints. For soils with 5-10% organic matter

#### Incorporation of Trilin With Bedding Implements

Bedding implements (listers and hipplers) may be used for weed control in cotton. Because provide thorough soil mixing under all conditions follow use directions to optimize weed control from single pass incorporation with bedding compared to conventional double pass incorporation rate recommended for the soil texture to be

#### Soil Preparation

Crop Residues or Existing Weeds: Ground or existing weeds, can interfere with uniform manageable level of such ground cover will into the top 2 to 3 inches of soil. Ground excessive, should be reduced by appropriate application of Trilin.

#### General Soil Conditions

The soil surface should be smooth enough to and efficient incorporation of Trilin. Apply sufficient to allow the breakup of large clods incorporation process.

#### Use Directions for Bedding Equipment

A lister or disk bedder may be used to implement according to the manufacturer's produce beds of the desired height. A ripper should be mounted on the bedder in a position but ahead of the bedder tool to help distribute bed. The use of bed tillage equipment such driven rod weeders or bed conditioners after provide additional soil mixing. Avoid deep untreated soil to the surface resulting in to conditions, cultural practices, bed tillage affect the distribution of Trilin treated soil. V dependent upon how uniformly Trilin treated surface at the time of planting.

If Trilin treated soil is moved, exposing untreated planting, a band application of Trilin at application may be required to restore uniform

#### PRECAUTION

Do not incorporate with bedding equipment if soil mixing.

#### Trilin for Weed Control in Conservation Tillage

Trilin may be applied and incorporated before to crop emergence, or at layby for weed control cotton. Single or multiple application may be application rates are not exceeded and recommended followed.

#### Strip Planting into Small Grain Cover Crops

Fall planted cover crops may be utilized protect developing crop seedlings from winter cotton, the cover crop may be treated with Trilin continued growth and development and p seedlings for water and soil nutrients. The

be planted for 18 months after application or for 10 months after a fall application of Trilin to avoid crop injury. If land has not been irrigated do not plant any of these crops for 18 months after a spring application or 20 months after a fall application of Trilin.

In those areas of Kansas, Nebraska, North Dakota, Oklahoma, South Dakota and Texas where at least 20 inches of rain and/or irrigation (total) was used to produce the crop, sorghum or oats should not be planted for 12 months after an application of Trilin.

Do not plant sorghum, proso millet or oats for 18 months after an application of Trilin if less than 20 inches of total water was used to produce the crop. Cool wet weather conditions during the early stage of growth may increase the possibility of injury to sorghum.

In all other areas receiving greater than 20 inches rainfall per year, moldboard plow before planting sugar beets where a spring application of Trilin was made the previous season. Also note planting restrictions listed in the section on control of rhizome johnsongrass and other high rate programs.

For vegetables other than those listed on this label, crops should not be planted within 5 months following the application of Trilin.

## FERTILIZER USE DIRECTIONS

### LIQUID FERTILIZERS

Trilin may be mixed with most liquid fertilizer materials. A combination of Trilin with solutions and suspension-type fertilizers will provide weed and grass control equal to the same rates of Trilin applied to water. Trilin label recommendations regarding rates per acre, crops, incorporation directions, special instructions, and precautions should be followed. Labeling and applications relating to liquid fertilizer mixing for individual state regulations are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

### Compatibility Testing Tank Mix Partners

A tank mixture of Trilin alone or with dry flowables, wettable powders, aqueous suspensions, flowables liquids or solutions may not combine properly with some liquid fertilizer materials. Always test a small quantity before full-scale mixing to determine whether a compatibility agent is needed and which agent does the best job. Phosphate ester type surfactants designed for use with liquid fertilizers are suggested. Use the following test to select the correct agent for your mixture.

1. Measure one pint of intended spray water or fertilizer solution into a jar.
2. Check the pH of the liquid and adjust if necessary.
3. Add in the given order the intended ingredients shaking well after each addition.
  - (a) Surfactants (spreaders), acidifiers, compatibility agents and activators; add one teaspoon for each pint/100 gallons.
  - (b) Dry ingredients (wetable powders, or dry flowables); add one tablespoon for each pound/100 gallons.
  - (c) Emulsifiable concentrates; add one teaspoon for each pint/100 gallons.
  - (d) Flowables; add one teaspoon for each pint/100 gallons.
  - (e) Soluble ingredients; add one tablespoon for each pound/100 gallons.
  - (f) Spreader-stickers; one teaspoon for each pint/100 gallons.
4. The final mixture should be uniform and smooth with no evidence of coagulation occurring. If incompatibility is evident, begin test again with a compatibility agent added first. Six drops is equivalent to four ounces/100 gallons. If this does not smooth the mixture try higher concentrations and other compatibility agents.

5. Allow the mixture to stand undisturbed thirty minutes. If separation occurs shake and observe the resulting mixture. If the mixture does not redispense do not attempt to spray the mixture. You may try:

- (a) More compatibility agents
- (b) Different formulations of the active ingredients (switch from wettable or emulsifiable concentrates to flowable or from wettable powder to emulsifiable concentrates).

### Mixing Instructions

When mixing Trilin in liquid fertilizers continuous agitation is required to prevent the Trilin from rising to the surface as an oily layer. Use a compatibility agent to make the Trilin emulsify properly. When tank mixing

weeds in addition to those weeds listed for Trilin.

Henbit (spring applications)	Oat Wild
Nightshade, Black	Ragweed, Common
Nightshade, Hairy	Smartweed, Pennsylvania
Nutsedge	Velvetleaf (Buttonweed)

Follow recommended soil preparation and incorporation procedures for Trilin. Apply this tank mix from 2 days prior to planting up to planting. Incorporate immediately after application.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2
PLUS			
Eptam 7E* (pts.)	2.5-3.5	2.5-3.5	2.5-3.5

\*To control annual grasses use Eptam 7E at a rate of 2.5 pints per acre. To control nutsedge and additional broadleaf weeds use 3.5 pints.

On soils with 2-5% organic matter use Trilin at 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. For soils with 5-10% organic matter use 2 pints of Trilin on all soils.

### PRECAUTION

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use this tank mix on some blackeyed peas (beans), lima beans, soybeans, mung beans and other flatpodded beans except Romano. Do not use foliage from treated plants for feed, forage or for grazing.

### Fall application in Idaho, Oregon and Washington

Apply and incorporate Trilin between October 15 and December 31. Use a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils and 1.5 pints on fine soils. Destroy established weeds during seedbed preparation.

## BEAN

### Guar and Mungbean

Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

## BEAN

### Snap and Lima

Apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils.

## BEAN

### Lentil

Apply and incorporate Trilin before planting. Plant no deeper than 1.5 inches when temperature and moisture in seedbed promote fast germination and emergence.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Spring Application	1.0	1.0	1.5
Fall Application	1.25	1.25	1.75

For soils with 2-5% organic matter use 1.5 pints per acre.

### PRECAUTION

Lentil tolerance to trifluralin is marginal. Injury may occur under conditions of plant stress such as cold weather, low fertility and disease or insect damage. Spring application may increase the probability of stand reductions.

protect developing crop seedlings from wind erosion, the cover crop may be treated with a continued growth and development and provide seedlings for water and soil nutrients. The soil continues to control wind erosion and provide crop until it is well established.

In strip planting, cotton is seeded into comp in the cover crop. Competition-free bands in unseeded drill rows when seeding the cover contact herbicide to prepare competition-free

### Application Before or After Planting Bands

Apply Trilin using low pressure ground equipment (weed free zone) or as a broadcast incorporation may occur before planting or emergence. If applied after planting, set not disturb the planted seed.

### Incorporation

Equipment should be adapted to the width of zone. Use equipment that will uniformly mix Trilin zone. Weed control, resulting from single incorporation equipment that does not resurface treated soil may be reduced compared to incorporation. Implements used to incorporate be operated so that they do not disturb the

### Broadcast Rates Per Acre

Use 1.0 to 2.0 pints per acre on all soils. Use range when additional sequential applications. Use the higher rate in the rate range when present, and where dense weed populations. For band treatments, reduce the application spacing and band width treated. For example where the row spacing is 36 inches would be broadcast rate per acre (12 inches divided by

### Layby Applications

Layby applications may be made in establish stage of growth up to layby, but not less than Trilin uniformly to the soil surface using drop broadcast rate of 1.0 pint per acre for coarse, and 2.0 pints per acre for fine textured soils. So a sweep-type cultivator or properly adjust cultivation equipment at speeds sufficient to provide exercise care to avoid mechanical injury to application rate may not exceed the layby application texture.

### Chemigation

Trilin may be applied through properly equipped control in conservation tillage cotton. Apply the 0.5 to 1.0 acre in the sprinkler irrigation. Apply planting, but prior to weed or crop germination established weeds. Soil incorporation is not chemigation. Soil treated with Trilin may be shallow weed control activity. See "Chemigation" section. Apply at a broadcast rate of 1.0-2.0 pints per acre rate in the rate range when additional sequential applications are present, where dense weed populations additional sequential applications of Trilin a Applications section).

### Sequential Applications

Trilin may be applied one or more times per season using the rates and methods of a season weed control. The maximum dosage application cannot exceed the rates shown. The maximum cumulative application rate for the same growing season cannot exceed 4 pint

Crop Rotation

1/2



ers are defined as implements with sweeps  
ls of 7 inches or less staggered so that no  
to 4 inches deep and operate at 5 mph or

**tioners:** Set to cut 3 to 4 inches deep and  
mph. These are defined as three or more  
used as a single tool. For example, 2 to  
3- shaped shanks with an effective sweep  
gged so that no soil is left untuned)  
xtine harrow followed by a ground driven

to 4 inches deep and operate at a speed  
y are effective for use on coarse and  
ng cultivator may be used on fine textured

to cut 2 to 4 inches deep and operate at a  
-All is effective when used on coarse or  
ly one incorporation pass is necessary in  
with the Do-All are necessary in flat

**ar disc-type implements):** Set Mulch  
ep and operate at 5 to 8 mph.

**tillers, cultivators, hoes):** Only one  
st to incorporate Trilin into the top 2 to  
ctors spaced to give a clean sweep of the  
not be operated at a speed greater than

he flexible tine-tooth harrow (Flextine,  
ut only for the special usages for which it  
ner equipment only as specified herein.

## I AFTER PLANTING

without reducing the weed control activity  
y result if cultivation is deeper than the  
untreated soil to the surface.

## OMMENDATIONS

as the broadcast rates of Trilin per acre.  
e the amount of Trilin in proportion to the  
acre. Apply Trilin any time after January 1  
nd is suitable for good incorporation. For  
rop recommendations or "FALL  
specific crop recommendations are not  
coarser soils or soils with lower organic  
or more organic matter do not use Trilin.

## LICATION DIRECTIONS

to use. Fill the sprayer 1/3 to 1/2 full with  
Add proper amount of Trilin providing  
ng the tank.

gitation is required (Sparger pipe agitators  
tion in spray tanks). To prevent foaming  
to the mixture during filling by placing the  
face of the water in the spray tank. Do not  
into the water source.

## Rate Chart for Impregnating Fertilizer With Trilin Added to a Ton of Fertilizer

CHART QUANTITIES LISTED ARE QUARTS OF TRILIN  
PER TON OF FERTILIZER

Fertilizer Rate Per Acre	TRILIN Rate Per Acre				
	1 pt.	1½ pts.	2 pts.	3 pts.	4 pts.
200 lbs.	5	7.5	10	15	20
250 lbs.	4	6	8	12	16
300 lbs.	3.33	5	6.67	10	13.33
350 lbs.	2.75	4.25	5.75	8.5	11.5
400 lbs.	2.5	3.75	5	7.5	10
450 lbs.	2.25	3.33	4.5	6.67	9

For rates other than those listed above use the following formula to  
calculate the amount of Trilin to be impregnated on a ton of dry bulk  
fertilizer.

$$\begin{array}{rcl} \text{Pts.} & & \text{Qts.} \\ \text{Trilin/Acre} & \times & 1000 \text{ lbs.} \\ & & \text{Fertilizer/Acre} \end{array} = \begin{array}{r} \text{Trilin/Ton of} \\ \text{Fertilizer} \end{array}$$

All individual state regulations relating to dry bulk fertilizer blending,  
registration, labeling and application are the responsibility of the individual  
and/or company selling the fertilizer and chemical mixture.

### Application

Use a properly calibrated applicator to spread the fertilizer/chemical  
mixture. The material should be applied uniformly to the soil surface.

### Incorporation

Follow Trilin incorporation procedures.

## APPLICATION INSTRUCTIONS ALFALFA

### Preplant

Trilin may be applied as a preplant incorporated for preemergence control  
of labeled weeds in direct seeded alfalfa. Apply and incorporate prior to  
planting according to product label directions.

### Broadcast Application Rates

Soil Texture	Trilin (pt/Acre)
Coarse	1.0
Medium	1.0-1.5*
Fine	1.5

\*Use lower range in areas receiving less than 20 inches of rainfall and  
irrigation.

Precautions: Some crop stand reduction and stunting may occur with use  
of Trilin, however, reduced weed competition will allow establishment of a  
quality stand.

### Mechanically Incorporated

Use a broadcast rate of 1.5 pints per acre on coarse soils and 2 pints on  
medium and fine soils (in areas receiving less than 20 inches average  
annual rainfall). Damage to the established alfalfa may be caused if the  
proper incorporation equipment is not used for thorough soil mixing.  
Recommended soil preparation application and incorporation instructions  
should be followed.

## COLE CROPS

### Broccoli, Brussels Sprout, Cabbage & Cauliflower

#### Transplant

Apply and incorporate Trilin prior to transplanting only.

#### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium  
and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints  
on all soils.

#### Direct-Seeded

Use Trilin before planting at a broadcast rate of 1 pint per acre on coarse  
and medium soils and 1.5 pints on fine soils and soils with 2-5% organic  
matter. Direct-seeded cole crops have exhibited marginal tolerance to  
recommended rates of Trilin.

**Warning:** Stunting or reduced stands may occur.

## COTTON

Apply Trilin before planting, immediately after planting, or at layby.

#### Preemergence

#### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

Use 1.5 pints per acre on coarse and medium textured soils, 2 pints on  
fine soils with 2-5% organic matter and 2 to 2.5 pints on all soils with  
5-10% organic matter.

#### Post Plant

Do not disturb the seed when incorporating Trilin postplant.

#### Layby

Trilin can be applied and incorporated any time up to layby, but not less  
than 90 days before harvest. Direct layby applications onto the soil  
between the rows and beneath emerged cotton plants at the same rates as  
as for a preemergence application.

#### Fall Application

Trilin may be applied and incorporated any time between October 15 and  
December 31. The ground may be left flat or bedded-up over winter.  
Where soil is left flat, take care not to turn up untreated soil from beds into  
furrows. On bedded ground, knock down beds to desired heights before  
planting, moving some treated soil into the furrow from the beds. Destroy  
established weeds during preparation of seedbed. Before planting,  
destroy weeds which may have become established in furrows during the  
uncovering of untreated soil during bedding. Trilin should not be applied  
to wet soil or soils which are subject to prolonged periods of flooding.

Application Over Standing or Shredded Cotton Stalks

13/29

as the broadcast rates of Trilin per acre. e the amount of Trilin in proportion to the acre. Apply Trilin any time after January 1 and is suitable for good incorporation. For crop recommendations or "FALL" specific crop recommendations are not coarser soils or soils with lower organic or more organic matter do not use Trilin.

## APPLICATION DIRECTIONS

to use. Fill the sprayer 1/3 to 1/2 full with water. Add proper amount of Trilin providing enough water to fill the tank.

Agitation is required (Sparger pipe agitators or agitators in spray tanks). To prevent foaming in the mixture during filling by placing the face of the water in the spray tank. Do not stir into the water source.

When water and start agitation. Add wettable powder and agitate until completely dispersed. Then add Trilin. Lastly, add soluble material. Provide continuous agitation during application and agitation must be stopped before material may settle to the bottom. If this happens, stop application, resuspend all of the material in the water. Agitation is particularly useful for this material to resuspend the settled material than it

Follow instructions for each material added to the tank. To insure proper dispersion in the tank water, make a slurry of wettable formulations with water and pour through wet screen in the top of the tank. No stirring in the tank should be used.

After use of the spray tank, wash the tank with water and continue the spraying operation. Thoroughly clean the tank lines and screens. Proper operation and uniform application increases efficiency. Check sprayer daily to insure proper operation. To insure uniform application do not use drifting of spray particles. Poor weed control is related to soils which are wet or are subject

Use the sprayer which will apply the spray evenly. Allow 10 to 15 gallons of water per acre.

Use 5 to 10 gallons of water or liquid fertilizer per acre. The nozzle height and height should be adjusted to the soil surface. To assure uniform application use swath markers or flagmen.

For areas with more than 20" average annual rainfall, use the rates listed for spring application. Use the rates listed for spring application where there are no specific fall application rates. It is recommended as a preemergence application in the fall for sugarbeets, potatoes and other crops. Apply and incorporate Trilin any time after January 31. Leave ground flat or bedded over. Do not be knocked down to desired height. Do not be taken not to turn up untreated soil from beds into furrows. Do not be taken not to turn up untreated soil from beds into furrows. Destroy weeds which have become established in untreated soil during bedding.

Trilin may be applied as a preplant incorporated for preemergence control of labeled weeds in direct seeded alfalfa. Apply and incorporate prior to planting according to product label directions.

### Broadcast Application Rates

Soil Texture	Trilin (pt/Acre)
Coarse	1.0
Medium	1.0-1.5*
Fine	1.5

\*Use lower range in areas receiving less than 20 inches of rainfall and irrigation.

Precautions: Some crop stand reduction and stunting may occur with use of Trilin, however, reduced weed competition will allow establishment of a quality stand.

### Mechanically Incorporated

Use a broadcast rate of 1.5 pints per acre on coarse soils and 2 pints on medium and fine soils (in areas receiving less than 20 inches average annual rainfall). Damage to the established alfalfa may be caused if the proper incorporation equipment is not used for thorough soil mixing. Recommended soil preparation application and incorporation instructions should be followed.

### Chemigation or Water Incorporated

Trilin may be surface applied for annual grass control in established alfalfa by chemigation, or ground or aerial broadcast application equipment. Refer to "Chemigation" section of this label for use directions for chemigation.

### Surface Application Activated by Rainfall or Irrigation

Broadcast surface applications of Trilin to established alfalfa may be activated by rainfall, sprinkler, flood or furrow irrigation. Rainfall or a single overhead sprinkler irrigation of 0.5 acre inch or more is required to activate Trilin. If activated by furrow irrigation, care should be taken to thoroughly wet beds between furrows: If rainfall or irrigation has not occurred within 3 days after application, Trilin may be mechanically incorporated. If mechanically incorporated, use equipment that will insure thorough soil mixing with minimum damage to the established alfalfa.

### Application Timing

Applications to established alfalfa for annual grass control can be made during dormancy or semi-dormancy, or during the growing season immediately after a cutting. Because Trilin does not control established weeds, application must be made prior to the expected time of weed germination. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather. To control these weeds, apply Trilin immediately after a cutting between August 1 and October 1, but prior to weed germination. When fall applied, Trilin controls bromegrass and cheat in addition to other labeled weeds that germinate after application.

Apply Trilin at a broadcast rate of 4 pints per acre on all soil textures by chemigation or surface applied and incorporated by rainfall or irrigation.

### RESTRICTIONS

Do not harvest forage within 21 days after application and do not harvest hay within 20 days after application. Do not apply more than 8 pints (4 lbs. a.i./A) per year. Apply no more than 4 pints of Trilin during any growing season. In the growing season following application of 4 pints of Trilin to alfalfa, plant only those crops for which Trilin is registered as a preplant treatment or crop injury does not occur.

## ASPARAGUS Established

Follow recommended soil preparation, application and incorporation procedures for Trilin. Trilin can be applied to established asparagus as a single or as a split application. In the winter or early spring, apply to asparagus after ferns are removed but before spear emergence or apply after harvest in the late spring or early summer before turning begins. Trilin will suppress volunteer seedling asparagus and field bindweed used at the following recommended rates and application schedules.

### Broadcast Rates Per Acre

Coarse	Medium	Fine
--------	--------	------

Use 1.5 pints per acre on coarse and medium textured soils, 2 pints on fine soils with 2-5% organic matter and 2 to 2.5 pints on all soils with 5-10% organic matter.

### Post Plant

Do not disturb the seed when incorporating Trilin postplant.

### Layby

Trilin can be applied and incorporated any time up to layby, but not less than 90 days before harvest. Direct layby applications onto the soil between the rows and beneath emerged cotton plants at the same rates as for a preemergence application.

### Fall Application

Trilin may be applied and incorporated any time between October 15 and December 31. The ground may be left flat or bedded-up over winter. Where soil is left flat, take care not to turn up untreated soil from beds into furrows. On bedded ground, knock down beds to desired heights before planting, moving some treated soil into the furrow from the beds. Destroy established weeds during preparation of seedbed. Before planting, destroy weeds which may have become established in furrows during the uncovering of untreated soil during bedding. Trilin should not be applied to wet soil or soils which are subject to prolonged periods of flooding.

### Application Over Standing or Shredded Cotton Stalks

Trilin may be broadcast applied, after September 1, over the top of standing cotton stalks or after shredding using ground or aerial equipment. Trilin must be incorporated once within 24 hours after application using a disc operated at 4 to 6 mph. The second incorporation may be delayed until spring prior to bedding. To avoid dilution of the herbicide or bringing untreated soil to the surface, the treatment should not be chiseled, ripped or deep plowed following incorporation.

### Broadcast Rates Per Acre - Fall Application Only

In Alabama, Arkansas, Northern Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee and Texas, apply and incorporate Trilin at a broadcast rate of 2 pints per acre on all soil types.

In Arizona, California and Nevada, a broadcast rate of 1.5 pints per acre should be used on coarse soil, 2 pints on medium soil and 2.5 pints on fine soil.

For other states where cotton may be grown, apply Trilin at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. For coarse soils with 2-5% organic matter use 1.5 pints. For soils with 5-10% organic matter use 2 pints.

### Incorporation of Trilin With Bedding Implements for Weed Control in Cotton

Bedding implements (listers and hippers) may be used to soil incorporate Trilin for weed control in cotton. Because bedding implements do not provide thorough soil mixing under all conditions, it is important to closely follow use directions to optimize weed control. Weed control resulting from single pass incorporation with bedding equipment will be reduced compared to conventional double pass incorporation. Use the application rate recommended for the soil texture to be treated.

### Soil Preparation

Crop Residues or Existing Weeds: Ground cover, such as crop residues or existing weeds, can interfere with uniform soil incorporation of Trilin. A manageable level of such ground cover will allow uniform incorporation into the top 2 to 3 inches of soil. Ground cover of crop residues, if excessive, should be reduced by appropriate soil tillage prior to the application of Trilin.

### General Soil Conditions

The soil surface should be smooth enough to allow for uniform application and efficient incorporation of Trilin. Apply Trilin when soil moisture is sufficient to allow the breakup of large clods and uniform mixing during the incorporation process.

### Use Directions for Bedding Equipment

A lister or disk bedder may be used to incorporate Trilin. Operate the implement according to the manufacturers use directions in order to produce beds of the desired height. A ripper shank, sweep or chisel shank should be mounted on the bedder in a position behind the spray nozzle.

fresh wet screen in the top of the tank. No  
in the tank should be used.

ills of the spray tank, wash the tank with  
inse and continue the spraying operation.  
oroughly clean the tank lines and screens.  
ibration and uniform application increases  
s. Check sprayer daily to insure proper  
tion. To insure uniform application do not  
ause drifting of spray particles. Poor weed  
plied to soils which are wet or are subject  
J.

ide sprayer which will apply the spray  
gallons of water liquid fertilizer per acre

n 5 to 10 gallons of water or liquid fertilizer  
le arrangements, ed and height should  
n application to oil surface. To assure  
use swath markers or flagmen.

ng more than 20" average annual rainfall)  
ndations. Use the rates listed for spring  
hich there are no specific fall application  
lin is recommended as a preemergence  
plied in the fall for sugarbeets, potatoes  
Apply and incorporate Trilin any time  
mber 31. Leave ground flat or bedded over  
eds should be knocked down to desired  
ome treated soil from beds into furrows.  
re should be taken not to turn up untreated  
perations. During seedbed preparation  
Destroy weeds which have become  
ncovering of untreated soil during bedding.  
re fall to soils which are wet, are subject to  
r where rice was grown the previous year.

## PRECAUTIONS RESTRICTIONS

ications on the label of each product used  
Under normal growing conditions and if  
Trilin will not harm the treated crop. Crop  
from over application. Erratic weed control  
om uneven application or improper soil  
ng disease, cold weather, deep planting,  
oncentration or drought may weaken crop  
ssibility of damage from Trilin. Under these  
ppment or reduced yields may result.

Montana, Nevada, New Mexico, Oregon,  
ng after a spring application of Trilin, sugar  
ould not be planted for 12 months or for 14  
oil should be plowed to a depth of 12 inches  
o prevent the possibility of crop injury. After  
(milo), proso millet, corn or oats should not  
er application or for 16 months after a fall  
op injury. If land has not been irrigated do  
r 18 months after a spring application or 20  
of Trilin.

braska, North Dakota, Oklahoma, South  
ast 20 inches of rain and/or irrigation (total)  
sorghum or oats should not be planted for  
of Trilin.

o millet or oats for 18 months after an  
ian 20 inches of total water was used to  
eather conditions during the early stage of

### Application Timing

Applications to established alfalfa for annual grass control can be made during dormancy or semi-dormancy, or during the growing season immediately after a cutting. Because Trilin does not control established weeds, application must be made prior to the expected time of weed germination. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather. To control these weeds, apply Trilin immediately after a cutting between August 1 and October 1, but prior to weed germination. When fall applied, Trilin controls bromegrass and cheat in addition to other labeled weeds that germinate after application.

Apply Trilin at a broadcast rate of 4 pints per acre on all soil textures by chemigation or surface applied and incorporated by rainfall or irrigation.

### RESTRICTIONS

Do not harvest forage within 21 days after application and do not harvest hay within 20 days after application. Do not apply more than 8 pints (4 lbs. a.i./A) per year. Apply no more than 4 pints of Trilin during any growing season. In the growing season following application of 4 pints of Trilin to alfalfa, plant only those crops for which Trilin is registered as a preplant treatment or crop injury does not occur.

## ASPARAGUS Established

Follow recommended soil preparation, application and incorporation procedures for Trilin. Trilin can be applied to established asparagus as a single or as a split application. In the winter or early spring, apply to asparagus after ferns are removed but before spear emergence or apply after harvest in the late spring or early summer before turning begins. Trilin will suppress volunteer seedling asparagus and field bindweed used at the following recommended rates and application schedules.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Split Application	1	1.5	2
OR			
Single Application	2	3	4

In any single calendar year, the maximum Trilin to apply is 2 pints per acre on coarse soils, 3 pints on medium soils and 4 pints on fine soils.

## BEAN Dry

Apply Trilin before planting using the following rates:

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. For soils with 5-10% organic matter use 2 pints on all soils.

### TRILIN WITH EPTAM TANK MIX

A tank mix of Trilin and Eptam will effectively control all the following weeds in addition to those weeds listed for Trilin.

Henbit (spring applications)	Oat Wild
Nightshade, Black	Ragweed, Common
Nightshade, Hairy	Smartweed, Pennsylvania
Nutsedge	Velvetleaf (Buttonweed)

Follow recommended soil preparation and incorporation procedures for Trilin. Apply this tank mix from 2 days prior to planting up to planting. Incorporate immediately after application.

### Broadcast Rates Per Acre

Coarse	Medium	Fine
--------	--------	------

fine soil.

For other states where cotton may be grown, apply Trilin at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. For coarse soils with 2-5% organic matter use 1.5 pints. For soils with 5-10% organic matter use 2 pints.

### Incorporation of Trilin With Bedding Implements for Weed Control in Cotton

Bedding implements (listers and hippers) may be used to soil incorporate Trilin for weed control in cotton. Because bedding implements do not provide thorough soil mixing under all conditions, it is important to closely follow use directions to optimize weed control. Weed control resulting from single pass incorporation with bedding equipment will be reduced compared to conventional double pass incorporation. Use the application rate recommended for the soil texture to be treated.

### Soil Preparation

Crop Residues or Existing Weeds: Ground cover, such as crop residues or existing weeds, can interfere with uniform soil incorporation of Trilin. A manageable level of such ground cover will allow uniform incorporation into the top 2 to 3 inches of soil. Ground cover or crop residues, if excessive, should be reduced by appropriate soil tillage prior to the application of Trilin.

### General Soil Conditions

The soil surface should be smooth enough to allow for uniform application and efficient incorporation of Trilin. Apply Trilin when soil moisture is sufficient to allow the breakup of large clods and uniform mixing during the incorporation process.

### Use Directions for Bedding Equipment

A lister or disk bedder may be used to incorporate Trilin. Operate the implement according to the manufacturers use directions in order to produce beds of the desired height. A ripper shank, sweep or chisel shank should be mounted on the bedder in a position behind the spray nozzles but ahead of the bedder tool to help distribute Trilin in the center of the bed. The use of bed tillage equipment such as rolling cultivators, P.T.O. driven rod weeders or bed conditioners after the bedding operation will provide additional soil mixing. Avoid deep tillage which might bring untreated soil to the surface resulting in loss of weed control. Weather conditions, cultural practices, bed tillage and planting procedures can affect the distribution of Trilin treated soil. Weed control obtained will be dependent upon how uniformly Trilin treated soil is distributed over the soil surface at the time of planting.

If Trilin treated soil is moved, exposing untreated soil, during bed tillage or planting, a band application of Trilin at planting or a postemergence application may be required to restore uniform weed control.

### PRECAUTION

Do not incorporate with bedding equipment if the soil is too wet for uniform soil mixing.

### Trilin for Weed Control in Conservation Tillage Cotton

Trilin may be applied and incorporated before planting, after planting prior to crop emergence, or at layby for weed control in conservation tillage cotton. Single or multiple application may be made so long as maximum application rates are not exceeded and rotational crop restrictions are followed.

### Strip Planting into Small Grain Cover Crops

Fall planted cover crops may be utilized to control wind erosion and protect developing crop seedlings from wind damage. Prior to planting cotton, the cover crop may be treated with a contact herbicide to prevent continued growth and development and prevent competition with crop seedlings for water and soil nutrients. The standing cover crop (now dead) continues to control wind erosion and provide protection to the developing crop until it is well established.

In strip planting, cotton is seeded into competition free bands established in the cover crop. Competition-free bands may be established by leaving unseeded drill rows when seeding the cover crop, or by tillage of use of a contact herbicide to prepare competition-free bands prior to planting.

### Application Before or After Planting (Within Competition-Free

20 inches of rain and/or irrigation (total) sorghum or oats should not be planted for Trilin.

millet or oats for 18 months after an n 20 inches of total water was used to other conditions during the early stage of maturity of injury to sorghum.

reater than 20 inches rainfall per year, sugar beets where a spring application of Trilin. Also note planting restrictions listed for johnsongrass and other high rate

e listed on this label, crops should not be planted until the application of Trilin.

## USE DIRECTIONS

liquid fertilizer materials. A combination of Trilin-type fertilizers will provide weed and Trilin rates of Trilin applied to water. Trilin label rates per acre, crops, incorporation and precautions should be followed. Refer to liquid fertilizer mixing for individual Trilin of the individual and/or company label for mixture.

### Lix Partners

or with dry flowables, wettable powders, Trilin liquids or solutions may not combine with other materials. Always test a small quantity to determine whether a compatibility agent is needed. Phosphate ester type Trilin liquid fertilizers are suggested. Use the Trilin agent for your mixture.

l spray water or fertilizer solution into a jar. lid adjust if necessary.

ended ingredients shaking well after each

l, acidifies, compatibility agents and Trilin for each pint/100 gallons.

ble powders or dry flowables); add Trilin pound/100 gallons.

add one teaspoon Trilin each pint/100 gallons.

one tablespoon for each pound/100 gallons.

aspoon for each pint/100 gallons. uniform and smooth with no evidence of incompatibility is evident. begin test again with Trilin first. Six drops is equivalent to Trilin does not smooth the mixture try higher Trilin compatibility agents.

undisturbed thirty minutes. If separation e resulting mixture. If the mixture does not spray the mixture. You may try:

ne active ingredients (switch from wettable Trilin to flowable or from wettable powder to Trilin).

Trilin continuous agitation is required to Trilin to the surface as an oily layer. Use a Trilin emulsify properly. When tank mixing dry flowables, wettable powders, aqueous Trilin or solutions in liquid fertilizer using a Trilin y important. If Trilin rises to the surface of Trilin emulsion is not properly formed, the oil may or suspension to form oil curds which is

### Nutsedge

### Velvetleaf (Buttonweed)

Follow recommended soil preparation and incorporation procedures for Trilin. Apply this tank mix from 2 days prior to planting up to planting. Incorporate immediately after application.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2
PLUS			
Eptam 7E* (pts.)	2.5-3.5	2.5-3.5	2.5-3.5

\*To control annual grasses use Eptam 7E at a rate of 2.5 pints per acre. To control nutsedge and additional broadleaf weeds use 3.5 pints.

On soils with 2-5% organic matter use Trilin at 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. For soils with 5-10% organic matter use 2 pints of Trilin on all soils.

### PRECAUTION

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use this tank mix on some blackeyed peas (beans), lima beans, soybeans, mung beans and other flatpodded beans except Romano. Do not use foliage from treated plants for feed, forage or for grazing.

### Fall application in Idaho, Oregon and Washington

Apply and incorporate Trilin between October 15 and December 31. Use a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils and 1.5 pints on fine soils. Destroy established weeds during seedbed preparation.

## BEAN

### Guar and Mungbean

Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

## BEAN

### Snap and Lima

Apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils.

## BEAN

### Lentil

Apply and incorporate Trilin before planting. Plant no deeper than 1.5 inches when temperature and moisture in seedbed promote fast germination and emergence.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Spring Application	1.0	1.0	1.5
Fall Application	1.25	1.25	1.75

For soils with 2-5% organic matter use 1.5 pints per acre.

### PRECAUTION

Lentil tolerance to trifluralin is marginal. Injury may occur under conditions of plant-stress such as cold weather, low fertility and disease or insect damage. Spring application may increase the probability of stand reductions.

## CARROT

Apply and incorporate Trilin prior to planting.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			

In strip planting, cotton is seeded into competition free bands established in the cover crop. Competition-free bands may be established by leaving unseeded drill rows when seeding the cover crop, or by tillage of use of a contact herbicide to prepare competition-free bands prior to planting.

### Application Before or After Planting (Within Competition-Free Bands)

Apply Trilin using low pressure ground equipment as a band (within the weed free zone) or as a broadcast treatment. Application and incorporation may occur before planting or after planting prior to crop emergence. If applied after planting, set incorporation equipment so as to not disturb the planted seed.

### Incorporation

Equipment should be adapted to the width of the competition-free band. Use equipment that will uniformly mix Trilin into the weed germination zone. Weed control resulting from single pass incorporation or with incorporation equipment that does not result in thorough mixing of Trilin treated soil may be reduced compared to conventional double pass incorporation. Implements used to incorporate Trilin after planting should be operated so that they do not disturb the planted seed or growing crop.

### Broadcast Rates Per Acre

Use 1.0 to 2.0 pints per acre on all soils. Use the lower rate in the rate range when additional sequential applications of Trilin are anticipated. Use the higher rate in the rate range where high crop residues are present, and where dense weed populations are anticipated.

For band treatments, reduce the application rate in proportion to the row spacing and band width treated. For example, treating a 12-inch band where the row spacing is 36 inches would require 1/3 of the recommended broadcast rate per acre (12 inches divided by 36 inches = 1/3).

### Layby Applications

Layby applications may be made in established cotton from the 4 true leaf stage of growth up to layby, but not less than 90 days before harvest. Apply Trilin uniformly to the soil surface using drop nozzles if necessary. Apply at a broadcast rate of 1.0 pint per acre for coarse, 1.5 pints per acre for medium and 2.0 pints per acre for fine textured soils. Soil incorporate using one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Operate cultivation equipment at speeds sufficient to provide vigorous soil mixing and exercise care to avoid mechanical injury to the crop. Cumulative layby application rate may not exceed the layby application rate given for each soil texture.

### Chemigation

Trilin may be applied through properly equipped chemical systems for weed control in conservation tillage cotton. Apply the recommended rate of Trilin in 0.5 to 1.0 acre in the sprinkler irrigation. Apply Trilin before planting or after planting, but prior to weed or crop germination. Trilin does not control established weeds. Soil incorporation is not required when applied by chemigation. Soil treated with Trilin may be shallow-cultivated without reducing weed control activity. See "Chemigation" section for detailed use directions. Apply at a broadcast rate of 1.0-2.0 pints per acre for all soils. Use the lower rate in the rate range when additional sequential applications of Trilin are anticipated. Use the higher rate in the rate range when high crop residue levels are present, where dense weed populations are anticipated, or where no additional subsequential applications of Trilin are to be made (See Sequential Applications section).

### Sequential Applications

Trilin may be applied one or more times sequentially during the growing season using the rates and methods of application described for full season weed control. The maximum dosage that can be used for a single application cannot exceed the rates shown for each application method. The maximum cumulative application rate that may be applied within the same growing season cannot exceed 4 pints per acre.

### Crop Rotation

See the Plantback Restrictions section for specific rotational crop restrictions. When the cumulative application rate exceeds the application rate of 1.5 pints per acre for coarse, 1.5 pints per acre for medium and 2.0 pints per acre for fine textured soils, plant only those crops for which Trilin can be applied as a preplant incorporated treatment in the season.



or suspension to form oil curds which is compatibility agent is helpful in causing liquid oiling mixtures with liquid fertilizers. Used at rates as low as 1.5 to 2 pints per ton mixed well with the fertilizer before adding the directions on the compatibility agent

phosphate ester type surfactants designed . They usually do not work as compatibility water.

il Company, Chicago, IL)  
 orporated, Aberdeen, NC)  
 inneapolis, MN)  
 o Company, Buckner, KY)  
 ls Company, Chicago, IL)  
 ward Chemical Co.)  
 Madison, WI)

applicator to spread the fertilizer/pesticide omly to the soil surface.

on procedures.

## DK FERTILIZERS

gnation or coating of dry bulk fertilizers. ers impregnated with Trilin has provided o the same rates of Trilin applied in water. imendations regarding rates per acre, special instructions, and precautions . A re of dry fertilizer impregnated with Trilin at uld be applied. Trilin can be used for used fertilizer, except coated ammonium ). Trilin will not be absorbed by these mixtures of these materials can be

other commonly used dry bulk fertilizer ould be applied uniformly to the fertilizer.

s for the rate of Trilin per acre should be low to determine the amount of Trilin to dry bulk fertilizer based on the amount of er acre.

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

## CASTOR BEAN

Apply and incorporate Trilin before or immediately after planting. If applied and incorporated after planting, set equipment so as to not disturb the seed.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

## CELERY

Apply Trilin as a soil incorporated treatment. Trilin may be applied to direct-seeded or transplant celery before planting, at planting or immediately after planting.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.25-1.5	1.5-2.0

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

Use the lower rate in the range in areas receiving less than 20 inches total annual rainfall and/or irrigation.

## CHICORY

Trilin may be applied as a preplant incorporated treatment to chicory grown either as a root crop or leafy vegetable.

**Cichorium intybus** Considered to be a root crop, may yield the following:

Chicory - the dried and processed root used as a coffee substitute.

Radicchio - green leaves harvested from field grown plantings.

Belgian Endive - white leaves grown in the dark growth from field grown rootstalks.

**Cichorium endiva** Considered to be a leafy vegetable, may yield the following:

Escarole - curly green leaves from field grown plantings.

Endive - very curly green leaves from field grown planting.

Apply Trilin as a soil incorporated treatment in spring or early summer prior to planting.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1.0	1.5	2.0

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils and 2.0 pints on fine soils. On soils with 5-10% organic matter use 2.0 pints on all soil textures.

2.0 pints per acre for fine textured soils, plant only those crops for which Trilin can be applied as a preplant incorporated treatment in the season following the Trilin application or crop injury may result.

Small grain cover crops that will not be grazed or harvested and are intended for prevention of wind erosion in conservation tillage cotton may be planted in the fall following spring applications of up to 2 pints per acre Trilin. Injury in the form of reduced stands or delayed emergence and development may result when small grains are planted under these conditions.

## SPECIAL USE DIRECTIONS FOR COTTON

### Fall Panicum

Apply and incorporate Trilin broadcast at the rate of 2 pints per acre on coarse and medium soils.

### Texas Panicum

Apply and incorporate at the labeled use rate for the soil type. Uniformly incorporate to a depth of at least 3 inches. Adequate soil moisture is essential for herbicide activation. Any deviation from these recommendations can result in reduced control.

### Rhizome Johnsongrass

Rhizome johnsongrass can be controlled in all cotton producing states except Arizona by using a double rate program applied for two consecutive years.

### Soil Preparation

For satisfactory results proper soil preparation is essential. To bring rhizomes to the top of the soil use a chisel plow or similar implement. Follow with a disc twice before application to cut rhizomes into small 2 to 3 inch pieces. Any emerged johnsongrass should be destroyed.

### Application

Choose one of the following programs which best meets your cultural practices:

For spring application apply Trilin prior to planting in the spring for 2 consecutive years. A broadcast rate of 2 pints per acre should be used on all soils.

### OR

For fall application use Trilin between October 15 and December 31 for 2 consecutive years at the same rates as a spring application.

### Incorporation

For good rhizome johnsongrass control deep incorporation is necessary. Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two passes are necessary with the second pass in a different direction from the first.

### Cultivation

To remove johnsongrass plants which have escaped control, timely cultivations during the crop season are necessary to obtain effective control. Effective control cannot be obtained with only 1 year of double rate Trilin use.

### Crop Rotation

Plant only rice and those crops for which Trilin can be applied as a preplant treatment following a double rate treatment season or injury may result

### Pigweed and Seedling Johnsongrass Control

In Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, South Carolina, Tennessee and Southern Virginia, apply Trilin preplant at a broadcast rate of 1 to 1.5 pints per acre on coarse soils, 1.5 to 2 pints on medium soils and 2 pints on fine soils.

17/29

### Additional Weed Grass Control

In the Texas Gulf Coast counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, apply Trilin at a broadcast rate of 1.5 pints per acre on coarse soils, 2 pints on medium soils and 3 pints on fine soils two weeks prior to planting.

### **PRECAUTION**

When using higher usage rates, plant cotton after early season adverse weather conditions to avoid additional stress to the cotton plants due to cool, wet weather early in the growth cycle, which could cause reduced stands, delayed maturity and reduced yields.

### **TANK MIXES AND OVERLAYS**

Observe all precautions and limitations on the label of each product used in tank mixes and overlays.

### TRILIN WITH COTTON-PRO – Arizona, New Mexico and West Texas

### TRILIN WITH CAPAROL 4L – Arizona, New Mexico and the upper and lower El Paso Valley, Texas

Trilin with Cotton-Pro or Caparol 4L will control certain grasses and broadleaf weeds listed for Trilin alone plus the following weeds.

Annual Morningglory	Prickly Sida (Teaweed)
Groundcherry (Annual)	Ragweed
Malva	Smartweed
Mustard	Wild Oat

Shallow germinating seedlings of cocklebur and coffeeweed will also be controlled.

### **Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.25-1.5	2
PLUS			
Cotton-Pro (pts.)	3.125	4	4
OR			
Caparol 4L (pts.)	2.4-3.2*	4	4

Do not use on sands and loamy sands. Use proportionally less for band application. \*Use less than 3.2 pints per acre only in Arizona.

### **Mixing Directions**

Add Cotton-Pro or Caparol 4L to a partially filled tank of water. Add Trilin and fill tank. During the filling and spraying operation, agitate continuously. If bypass agitation is used, minimize foaming by hanging the bypass line stop at the bottom of the tank. Avoid leaving the spray mixture in the tank without constant agitation.

### **Crop Rotations**

After a spring application of Trilin plus Cotton-Pro or Caparol 4L, cabbage, okra, onions and peas may be planted in the fall. Winter barley, winter rye and winter wheat, if plowed down and not used for food or feed, can be planted in the fall also. Refer to the Cotton-Pro or Caparol 4L label for directions, and precautions.

### **PRECAUTION**

Do not use Trilin with Cotton-Pro or Caparol 4L tank mix in the cut areas of newly leveled fields, in areas of excess salt or where flooding over the beds is likely to happen. Do not plant cotton in tractor wheel depressions. These conditions may cause crop injury. On mulch planted cotton, water back only after cotton seedlings are well-established. Do not feed foliage from treated plants to livestock or graze treated areas.

### TRILIN WITH METURON 4L, METURON 80DF, COTORAN 4L OR COTORAN DF (Except in Arizona)

Follow recommended soil preparation and incorporation procedures for Trilin. A tank mix of Trilin with Meturon 4L, Meturon 80DF, Cotoran 4L or Cotoran DF effectively controls all the annual grasses and broadleaf weeds listed for Trilin alone plus these additional weeds:

Buttonweed	Morningglory	Sesbania
Cocklebur	Prickly Sida (Teaweed)	Sicklepod

### **Incorporation Directions**

Incorporate one time within 24 hours after application. The second incorporation should be performed in the spring prior to seeding flax. The incorporation operations should result in a thorough mixing of Trilin with soil. Otherwise, erratic weed control may result.

### **Incorporation Equipment**

Follow recommended soil incorporation procedures for Trilin. Incorporation operations or any other tillages performed in the spring prior to seeding should be relatively shallow so as to maintain a firm seedbed, and the seedbed should be packaged just prior to seeding. Seeding should be done with a press drill or hoe drill. Seed into a moist seedbed no more than 1.5 inches deep. Do not delay the first incorporation more than 24 hours after application.

## **FRUIT AND NUT CROPS AND VINEYARDS**

### **For areas receiving more than 20" average annual rainfall**

On new plantings of citrus, pecan trees and vineyards apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. On soils with 2-5% organic matter use 1.5 pints on fine soils and on soils with 5-10% organic matter use 2 pints. Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures for non-bearing established plantings of citrus and pecan trees and bearing plantings of grapefruit, lemon, orange, pecan, tangelo and tangerine trees.

### **For areas receiving less than 20" average annual rainfall**

On new plantings of almond, apricot, citrus, nectarine, peach, pecan and walnut trees apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils, 1.5 pints on fine soils, on soils with 2-5% organic matter use 1.5 to 2 pints and 2 pints on soils with 5-10% organic matter.

### **New Plantings of Vineyards**

Apply and incorporate Trilin prior to planting at a broadcast rate of 1 to 1.5 pints per acre on coarse soils, 1.5 to 3 pints on medium soils and 3 to 4 pints on fine soils or soils with 2-10% organic matter. Do not use more than 2 pints per acre on heat-treated grape rootings.

### **Postplant Application**

For postplant applications on bearing or non-bearing established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, plum, prune, tangelo, tangerine and walnut trees. Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures. Do not apply to vineyards within 60 days of harvest. In established plantings, use Trilin as a directed spray to the soil. Use incorporation methods not injurious to the trees or vines.

### **Rhizome Johnsongrass Control**

For areas receiving less than 20 inches average annual rainfall. Control rhizome johnsongrass with postplant applications in bearing and non-bearing established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees by applying Trilin for 2 consecutive years.

### **Soil Preparation**

Soil should be worked thoroughly to bring the rhizomes near the surface.

### **Application**

Use Trilin at a broadcast rate of 2 quarts per acre on all soil textures each year for 2 consecutive years. Do not apply to vineyards within 60 days of harvest.

### **Incorporation**

Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary using a different direction for the second.

### **Cultivation**

As some johnsongrass plants will escape, cultivation is necessary to obtain effective control. Effective control cannot be obtained with only

## **POTATC**

### **(All States Except**

Apply and incorporate Trilin after plar immediately following dragoff or after emerged.

### **Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	
Areas receiving less than 20" average annual rainfall	1
Areas receiving greater than 20" average annual rainfall	1

For soils with 2-5% organic matter use 1.5 soils, 2 pints on soils with 5-10% organic r

Incorporation equipment should be set to furrow with a layer of treated soil. If the her bed, potato emergence may be retarded. Do not completely cover the foliage with tr and incorporated after potato plants l completely cover foliage at subsequent incorporation machinery does not dar elongating sprouts.

### **Split Application in Idaho, Oregon and I**

Apply and incorporate 0.75 pint of Trilin 0.75 pint after planting when potato plant soils, except do not apply to soils containi Follow incorporation directions listed abov planting.

### TRILIN WITH EPTAM TANK MIX In Ka North Dakota, Oklahoma, South Dakota

This tank mix will effectively control the those weeds controlled by Trilin.

Henbit (spring applications)	O
Nightshade Black	R
Nightshade Harry	S
Nutsedge	V

Follow the recommendations for soil p procedures for Trilin. Trilin with Eptam t planting, but before crop emergence. I normally dragged off, this tank mix should l to or immediately following drag off.

### **Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	
Areas receiving less than 20" average annual rainfall	1
Areas receiving greater than 20" average annual rainfall	1
PLUS	
Eptam 7E	1.75-7**

\*\*For nutsedge control use the higher rate On soils with 2-5% organic matter use 1.5 medium soils and on soil with 5-10% org soils.

### **PRECAUTION**

Follow direction and precautions on th Observe all precautions and limitations on in tank mixes and overlays. Do not use feed, forage or for grazing.

Groundcherry, Wright  
Jimsonweed

Ragweed  
Ryegrass

Smartweed  
Tumbleweed

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Meturon 4L (pts.)	2	3.125	4
OR			
Meturon 80DF (lbs.)	1.25	1.9	2.5
OR			
Cotoran 4L (pts.)	2	3.125	4
OR			
Cotoran DF (lbs.)	1.2	2	2.4

**Mixing Directions**

Add Meturon or Cotoran to a partially filled tank of water. Add the Trilin after Meturon or Cotoran is thoroughly mixed and continue filling. Agitate continuously throughout the filling and application operations. Do not leave spray mixture in tank without constant agitation. If bypass agitation is used, minimize foaming by hanging the bypass line stop at the bottom of the tank. Apply in 15 to 40 gallons of water per acre.

**West Texas Only**

Do not use tank of Trilin plus Meturon or Cotoran on sandy, loamy sand or fine sandy loam soils. Do not use on cotton planted in furrows.

**Arkansas, Louisiana and Mississippi Only**

On sandy loam soils low in organic matter use 1.5 pints Meturon 4L or Cotoran 4L or 1 pound Meturon 80DF or Cotoran DF in tank mix with Trilin.

**New Mexico**

Do not plant treated land with crops other than cotton until 1 year after the last application. Do not use on sandy or coarse textured soils of less than 1% organic matter.

**PRECAUTION**

Crop injury may result if treated land is planted to anything but cotton within six months of the application of Trilin plus Meturon or Cotoran. Do not feed foliage from treated cotton plants or gin trash to livestock. Do not mix Trilin plus Meturon or Cotoran with liquid fertilizer.

**TRILIN PREPLANT FOLLOWED BY METURON 4L, METURON 80DF, COTORAN 4L OR COTORAN DF OVERLAY**

Apply and incorporate Trilin as recommended for the specific soil texture. Apply Meturon 4L or Cotoran 4L at 2 to 4 pints per acre or Meturon 80DF or Cotoran DF at 1.2 to 2.4 pounds per acre as a preemergence surface treatment. Use the lower rate on light silt and sandy soil low in organic matter.

**PRECAUTION**

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

**TRILIN PREPLANT FOLLOWED BY DIREX 4L OR KARMEX DF OVERLAY (East of the Mississippi River plus Arkansas, Southeastern Missouri, Louisiana and Eastern Texas)**

Incorporate Trilin prior to planting. Follow with a preemergence application of Direx 4L or Karmex DF. This will effectively control all the weeds controlled by Trilin alone plus these additional weeds.

Annual Ground, Cherry	Pennycress	Velvetgrass
Annual Morningglory	Ragweed	Wild Lettuce
Dogfennel	Shepherdspurse	Wild Mustard

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Direx 4L (pts.)	0.5	1	1.75

1 year of thin use.

**PRECAUTION**

Crop injury may result if the 2 quart rate is used on new plantings. Orchard or vineyards should not be interplanted with other crops. If the treated vineyards and orchards are diverted to other crop uses, plant only those crops for which Trilin has been registered as a preplant treatment for the next cropping season.

**Bindweed control In California**

Use Trilin for the control of field bindweed in vineyards and for almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees.

Use Trilin at a broadcast rate of 4 pints per acre on all soil textures. Trilin must be applied in the spring with a specially designed spray blade which applies a thin concentrated layer at a soil depth of 4 to 6 inches. This layer of Trilin prevents bindweed shoots from emerging.

**Land Preparation**

All weeds and grasses should be destroyed with soil tillage prior to applying Trilin. This tillage is necessary to prevent trash from interfering with the operation of the spray blade.

**Equipment**

A spray blade capable of running 4 to 6 inches below the surface of the soil should be used. The spray blade should be equipped with nozzles located under the blade and directed so that the Trilin spray will be trapped under the soil which is flowing over the blade as it is pulled through the soil. A sufficient number of nozzles should be used with spacing that will uniformly apply the Trilin underground in a thin horizontal layer.

**Application**

Use Trilin in 40 to 80 gallons of water per acre. Operate the spray blade at a depth of 4 to 6 inches.

**PRECAUTION**

After rainfall or irrigation some soils may crack as they dry. Bindweed may emerge if the cracks extend through the Trilin layer. Prevent or eliminate cracks by shallow discing or other tillage. Avoid deep tillage which disturbs the subsurface layer. Cultivation or tillage also aids the control of germinating seeds.

**IRRIGATION WATER RINGS  
NON-BEARING CITRUS TREES**

Apply to non-bearing citrus trees through irrigation water rings to provide preemergence weed control. Mix at a rate of 12 fluid ounces of Trilin per 500 gallons of water. Agitate until uniformly dispersed in tank.

Apply 10 gallons of the mixture per four foot diameter water ring per tree. Trilin should be applied at the second or third watering and should not be applied in combination with any other pesticide.

**GREENS**

**Turnip greens grown for processing,  
Collards, Kale and Mustard Greens**

Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

**HOPS**

Apply and incorporate Trilin while the crop is dormant using a broadcast rate of 1 pint per acre on coarse soils, 1.25 pints on medium soils and 1.5 pints on fine soils. Use incorporation equipment that will insure thorough soil mixing with minimum damage to the crop.

**KENAF**

**Ground Application**

Use a low pressure herbicide sprayer which will apply the spray

**Oregon and Washington**

Trilin with Eptam may be applied prior to or after planting at a rate of 0.75 pint of Trilin per acre and 3.5 pints of Eptam per acre on all soil textures. Incorporate immediately.

**PRECAUTION**

Do not use this tank mix both before and after planting. Observe all precautions and limit product used in tank mixes and overlays. Do not use on plants for feed, forage or for grazing.

**RADISH**

Apply and incorporate as a preplant soil treatment.

**Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	1

**RAPES EED (CANOLA)**

(Except In Alaska)

Follow recommended procedures for soil preparation. Trilin may be applied in the fall or early spring. Use incorporation equipment to incorporate to depth specified on label.

**Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	1

**SAFFLOW**

Recommended soil preparation application for Trilin should be followed. Use Trilin between October 15 and December 31.

**Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	1

Areas receiving less than 20" average annual rainfall

Areas receiving greater than 20" average annual rainfall

On coarse and medium soils use 1.5 pints of Trilin with 2-5% organic matter. Use 2 to 2.5 pints on fine soils with 2-5% organic matter.

**For Fall Application in Arizona, Idaho, Utah, Washington and Wyoming**

Apply and incorporate Trilin any time between October 15 and December 31. Ground may be left flat or bedded-up. Weeds should be knocked down to desired height. Care should be taken during spring bedding up untreated soil. Destroy established weeds before planting. Destroy weeds that become established during spring bedding up. Apply and incorporate Trilin at a rate of 1.5 pints per acre on coarse soils, 2 pints on medium soils, and 2.5 pints on fine soils. Trilin should not be applied to soils that are subject to prolonged periods of flooding.

**SOUTHERN**

Apply and incorporate Trilin prior to planting.

**Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	1

Areas receiving less than 20" average annual rainfall

10/16

OR Karmex DF (lb.) 0.33 0.67 1

**PRECAUTION**

Observe all precautions and limitations on the label of each product used in tank mixes and overlays. Direx 4L or Karmex DF should not be used on soils with less than 1% organic matter as crop injury may result. Do not feed foliage from treated plants to livestock or graze treated areas.

**COTTON: USE RESTRICTION**

Do not apply more than 2.0 pounds a.i./A (4 pints Trilin Herbicide) per application and do not apply more than 4 pints per crop year (either fall application through layby application or preplant plus post plant through layby).

**TREES GROWN FOR PULPWOOD  
Aspen, Cottonwood and Poplar**

**New Plantings**

Apply and incorporate Trilin before planting.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1.0	1.25-1.5	1.5-2.0

For soils with 2-5% organic matter use 1.5 to 2 pints. On soils with 5-10% organic matter use 2.0 pints. Use lower rate in areas receiving less than 20 inches total rainfall and irrigation.

**Established Plantings**

Apply as a directed spray to the soil and use incorporation methods not injurious to the crop.

**Broadcast Rates Per Acre**

**All Soil Texture**

	2.0-4.0
Trilin (pts.)	2.0-4.0

Application rate within the rate range may be adjusted according to weed pressure.

**Johnsongrass Suppression**

Proper soil preparation before application is necessary for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the soil surface. Then work the soil twice using a tandem disc to cut rhizomes into small (2 to 3 inch) pieces and to destroy emerging johnsongrass.

**Broadcast Rates Per Acre**

**All Soil**

	4.0
Trilin (pts.)	4.0

**Incorporation:** Incorporate twice with a tandem disc set to cut 4 to 6 inches deep and operated at 4 to 6 mph.

**Cultivation:** Some johnsongrass plants will escape. Timely cultivation with tillage implements or spot spraying with effective postemergence herbicides will improve the level of johnsongrass control.

**CUCURBITS**

**Cantaloupe, Cucumber and Watermelon**

**Postplant Emerged in Western United States including Texas**

Apply Trilin as a directed spray to the soil between the rows and beneath plants which are in the 3 to 4 true leaf stage.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints

Using a low-pressure broadcast sprayer which can apply and spray uniformly, apply Trilin in 5 to 40 gallons of water (broadcast spray) per acre.

**Aerial Application**

For aerial spraying apply Trilin in 5 to 10 gallons of water. Pump pressure, nozzle arrangements, speed and height should be adjusted to provide a uniform application to the soil surface. To assure proper application spray widths use swath markers or flagmen.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2

**Incorporation Directions**

**General Directions**

Erratic weed control and/or crop injury may result if directions are not followed. The soil surface should be free of existing weeds and excessive trash or clods.

**Before Planting**

Incorporation of Trilin must take place within 24 hours after application and may be accomplished in one pass with a sweep type or a rolling cultivator. Follow recommended soil incorporation procedures for Trilin.

**PRECAUTION**

Without reducing the weed control activity of Trilin, soil treated with Trilin may be shallow cultivated, rotary hoed or handhoed. Do not cultivate deeper than (1 to 2 inches) the Trilin treated layer of soil as this may bring untreated soil to the surface and poor weed control may result. Kenaf treated with Trilin must not be used as a food, forage or feed.

**LUPINE**

Apply and incorporate before planting.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints.

**MUSTARD**

Grown for seed or processing for food in Minnesota, Montana and North Dakota. Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

**NUT CROPS**

See FRUIT AND NUT CROPS AND VINEYARDS for instructions.

**OKRA**

Apply and incorporate Trilin before or immediately after planting. If applied and incorporated after planting, set equipment so as to not disturb the seed.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

20" average annual rainfall

Areas receiving greater than 20" average annual rainfall

For soils with 2-5% organic matter use 1.5 and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

**SOYBEAN**

**Preemergence**

Follow recommended soil preparation, and procedures for Trilin.

**Broadcast Rates Per Acre**

	Coarse
Trilin (pts.)	1

Use 1.5 pints per acre on coarse and medium soils with 2-5% organic matter. Use 2 pints on fine soils with 5-10% organic matter.

**Fall Application**

Trilin may be applied and incorporated in October and December 31. The ground may be left bare. Where soil is left flat, care should be taken to bed down to desired heights before planting, and to bed down furrows from the beds. Established weed preparation of seedbed. Before planting, weeds should become established in furrows due to the bedding. Trilin should not be applied subject to prolonged periods of flooding of the previous year.

Apply and incorporate Trilin at a broadcast rate of 1.5 pints on coarse and medium soils and 2.5 pints on fine soils in northern Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, and Texas.

For other states where soybeans may be grown, use 1 pint per acre on coarse soils, 1.5 pints on medium soils, and 2 pints on fine soils. For coarse soils use 1.5 pints. For soils with 5-10% organic matter use 2 pints.

**SPECIAL USE DIRECTIONS FOR**

**Fall Panicum**

Apply and incorporate Trilin broadcast at 1.5 pints per acre on coarse and medium soils.

**Crop Rotation**

Plant only rice and those crops for which preplant treatment following a double rate result.

**Texas Panicum**

Apply and incorporate at the labeled use rate and incorporate to a depth of at least 3 inches. Recommendations can result in reduced control of weeds.

**Pigweed and Seedling Johnsongrass Control**

In Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Tennessee and southern Virginia, apply Trilin at 1 to 1.5 pints per acre on coarse soils, and 2 pints on fine soils. **Exception:** Use 1.5 pints on Louisiana.

**Additional Weed and Grass Control**

In the Texas Gulf Coast counties of Brazoria, Galveston, Harris, Jackson, Jefferson, Victoria, Waller and Wharton, apply Trilin at 1.5 pints per acre on coarse and medium soils, and 2 pints on fine soils.

# FIELD CORN, GRAIN SORGHUM AND CORN FODDER, FORAGE AND SILAGE

Apply Trilin to field corn or grain sorghum (8 inches or taller) as an over-the-top or directed spray to effectively control weeds listed for Trilin. Trilin applied as an over-the-top spray or as a directed spray in field corn and grain sorghum will control shattercane in addition to those other weeds listed on the label for Trilin.

## Soil Preparation

Cultivate before a Trilin application to insure loose, friable soil, to remove established weeds and to cover the base of plants with soil.

## Application Directions

Trilin should be applied and incorporated at the recommended rates for the soil texture when the crop is well established (8 inches or taller). Trilin may be applied either as an over-the-top spray or as a directed spray. Drop nozzles should be used if foliage prevents uniform coverage of soil surface. Soil incorporation may be accomplished with only one pass of a sweep-type cultivator or a properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at 6 to 8 mph. Set the middle sweeps so as to avoid exposing untreated soil. Adjust the incorporation tools to prevent crop injury.

## Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2

## Corn Only

Apply 1 to 1.5 pints per acre in Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia to control fall panicum and Texas panicum.

## TRILIN WITH ATRAZINE TANK MIX

Trilin can be tank mixed with atrazine for additional weed control in field corn and grain sorghum.

## Broadcast Rates Per Acre

	Coarse*	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2

## PLUS

AAtrex 4L (pts.) 2.4 4.75 6  
When using AAtrex 4L use the rates listed above. For other atrazine formulations, use equivalent rates. When using AAtrex NineO 1 pint of AAtrex 4L equals 0.55 pound of NineO. One pint of AAtrex 4L equals 0.62 pound of Atrazine 80W.

\*Do not use the above tank mix on coarse soils for grain sorghum.

Apply and incorporate the Trilin/Atrazine tank mix as directed on the Trilin label for field corn and grain sorghum.

## PRECAUTION

Do not apply Trilin to sweet corn or corn grown for seed. Do not apply to corn or sorghum as a preplant or preemergence treatment or crop injury may occur. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

# FLAX

## Fall Application

Trilin may be applied and incorporated in the fall for weed control in spring seeded flax. Ground cover from existing weeds or previous crop should be at a manageable level so that there is no interference with incorporation.

Apply at the broadcast rates per acre of 1 pint on coarse soil, 1.5 pints on medium soil and 2 pints for fine soils.

PCA

# Dry and English

## Trilin Alone

Apply and incorporate Trilin prior to planting at a rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils.

## Fall application in Idaho, Oregon and Washington

Apply and incorporate Trilin any time between October 15 and December 31 using a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils and 1.5 pints on fine soils. Destroy established weeds during seedbed preparation. Do not apply Trilin in the fall to soils which are wet or are subject to prolonged periods of flooding.

## TRILIN WITH FAR-GO TANK MIX in Idaho, Oregon and Washington

Trilin plus Far-Go controls wild oat in addition to other annual grasses and broadleaf weeds controlled by Trilin.

## Application Rates

Broadcast 0.75 pint of Trilin per acre on coarse and medium soils, 1 pint on fine soils. Use 1.25 quarts of Far-Go per acre for all soil textures.

## Incorporation Directions

Apply and incorporate up to 3 weeks prior to planting. Follow recommended incorporation procedures for Trilin.

## PRECAUTION

Do not apply to lentils. Leaf crinkling and delayed maturity of peas may occur particularly on clay points in the northwest; but this is usually more than offset by a reduction of wild oat. Do not use foliage from treated plants for feed or grazing. Observe all precautions and limitations on the label of each product used in tank mixes and overlays.

# PEANUT

## Spanish Peanut, Florunner and Florigiant in New Mexico, Oklahoma and Texas

Apply and incorporate Trilin prior to planting, at planting or immediately after planting using a broadcast rate of 1 pint per acre on coarse soils. When incorporating after planting take care not to disturb the seed.

# PEPPER

## Transplant

Apply and incorporate Trilin prior to transplanting only.

## Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium, 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

up to two weeks before planting.

## Charcoal Soils in Arkansas, Louisiana

Freshly cleared land sometimes contain charcoal from burning debris. This charcoal tendency to bind Trilin and reduce its effectiveness. If such conditions exist, higher rates of Trilin are necessary. Injury can occur if increased rates are used where charcoal matter is not present in the soil. In the burr present, consequently, poor weed control. Use increased rate of Trilin.

Use Trilin broadcast at the rate of 1.5 to 2.5 pints on coarse soils and 3 pints on fine soils. Soil preparation, application and incorporation.

## Red Rice in Arkansas, Louisiana, Mississippi

Partial suppression or control of red rice can be achieved by the application of Trilin at the following recommended rate.

## Broadcast Rates Per Acre

Application	Coarse	Medium	Fine
Year 1	2	3	4
Year 2	1	1.5	2

If high organic matter and/or charcoal are present, use the following rates for the second year as follows for Arkansas, Louisiana, Mississippi.

## Broadcast Rates Per Acre

### Coarse

Trilin (pts.)	1
---------------	---

## Crop Rotation

Use a two year program for red rice control. In the first year plant soybeans. The second year plant rice. Trilin can be used preplant using the type and charcoal level. Do not plant rice until the third year.

## Rhizome Johnsongrass in Eastern United States

Rhizome johnsongrass can be acceptably controlled by Trilin for two consecutive years as follows.

## Soil Preparation

For satisfactory results proper soil preparation is essential. For rhizomes to the top of the soil use a chisel plow. Follow twice with a disc prior to application. Use 3 inch pieces and to destroy any emerged plants.

## Application

Choose one of the following programs and follow the practices.

For spring application use Trilin prior to planting. A broadcast rate of 1.5 pints on coarse soils, 3 pints on medium soils, 3 pints on coarse soils with 2-5% organic matter and 5-10% organic matter.

## OR

For fall application use Trilin between October and November for two consecutive years using the same rate.

## OR

A split application of Trilin may be used for two consecutive years using the following rates.

20/12

## Control

ties of Brazoria, Calhoun, Chambers, Fortson, Jefferson, Liberty, Matagorda, Orange, apply Trilin at a broadcast rate of 1.5 pints on medium soils and 3 pints on fine soils

As plant cotton after early season adverse additional stress to the cotton plants due to growth cycle, which could cause reduced reduced yields.

Simulations on the label of each product used

### Arizona, New Mexico and West Texas Arizona, New Mexico and the upper and

Caparol 4L will control certain grasses and alone plus the following weeds.

Prickly Sida (Teaweed)

Ragweed

Smartweed

Wild Oat

Species of cocklebur and coffee-weed will also be

Coarse	Medium	Fine
1	1.25-1.5	2
3.125	4	4
2.4-3.2*	4	4

my sands. Use proportionally less for band 2 pints per acre only in Arizona.

to a partially filled tank of water. Add Trilin filling and spraying operation, agitate on is used, minimize foaming by hanging the of the tank. Avoid leaving the spray mixture agitation.

Trilin plus Cotton-Pro or Caparol 4L, cabbage, planted in the fall. Winter barley, winter rye low and not used for food or feed, can be r to the Cotton-Pro or Caparol 4L label for

Pro or Caparol 4L tank mix in the cut areas as of excess salt or where flooding over the ot plant cotton in tractor wheel depressions. crop injury. On mulch planted cotton, water gs are well-established. Do not feed foliage k or graze treated areas.

### IL METURON 80DF, COTORAN 4L OR (Arizona)

eparation and incorporation procedures for Meturon 4L, Meturon 80DF, Cotoran 4L or rils all the annual grasses and broadleaf plus these additional weeds:

## Incorporation Directions

Incorporate one time within 24 hours after application. The second incorporation should be performed in the spring prior to seeding flax. The incorporation operations should result in a thorough mixing of Trilin with soil. Otherwise, erratic weed control may result.

## Incorporation Equipment

Follow recommended soil incorporation procedures for Trilin. Incorporation operations or any other tillages performed in the spring prior to seeding should be relatively shallow so as to maintain a firm seedbed, and the seedbed should be packaged just prior to seeding. Seeding should be done with a press drill or hoe drill. Seed into a moist seedbed no more than 1.5 inches deep. Do not delay the first incorporation more than 24 hours after application.

## FRUIT AND NUT CROPS AND VINEYARDS

### For areas receiving more than 20" average annual rainfall

On new plantings of citrus, pecan trees and vineyards apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. On soils with 2-5% organic matter use 1.5 pints on fine soils and on soils with 5-10% organic matter use 2 pints. Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures for non-bearing established plantings of citrus and pecan trees and bearing plantings of grapefruit, lemon, orange, pecan, tangelo and tangerine trees.

### For areas receiving less than 20" average annual rainfall

On new plantings of almond, apricot, citrus, nectarine, peach, pecan and walnut trees apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils, 1.5 pints on fine soils, on soils with 2-5% organic matter use 1.5 to 2 pints and 2 pints on soils with 5-10% organic matter.

### New Plantings of Vineyards

Apply and incorporate Trilin prior to planting at a broadcast rate of 1 to 1.5 pints per acre on coarse soils, 1.5 to 3 pints on medium soils and 3 to 4 pints on fine soils or soils with 2-10% organic matter. Do not use more than 2 pints per acre on heat-treated grape rootings.

### Postplant Application

For postplant applications on bearing or non-bearing established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, plum, prune, tangelo, tangerine and walnut trees. Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures. Do not apply to vineyards within 60 days of harvest. In established plantings, use Trilin as a directed spray to the soil. Use incorporation methods not injurious to the trees or vines.

### Rhizome Johnsongrass Control

For areas receiving less than 20 inches average annual rainfall. Control rhizome johnsongrass with postplant applications in bearing and non-bearing established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees by applying Trilin for 2 consecutive years.

### Soil Preparation

Soil should be worked thoroughly to bring the rhizomes near the surface.

### Application

Use Trilin at a broadcast rate of 2 quarts per acre on all soil textures each year for 2 consecutive years. Do not apply to vineyards within 60 days of harvest.

### Incorporation

Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary using a different direction for the second.

### Cultivation

As some johnsongrass plants will escape, cultivation is necessary to

## POTATO

### (All States Except Maine)

Apply and incorporate Trilin after planting prior to emergence or immediately following dragoff or after the potato plants have fully emerged.

### Broadcast Rates Per Acre

Trilin (pts.)	Coarse	Medium	Fine
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils, 2 pints on soils with 5-10% organic matter.

Incorporation equipment should be set to uniformly cover the bed and furrow with a layer of treated soil. If the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. Do not completely cover the foliage with treated soil when Trilin is applied and incorporated after potato plants have fully emerged. Do not completely cover foliage at subsequent cultivations. Be careful that incorporation machinery does not damage potato seed pieces or elongating sprouts.

### Split Application in Idaho, Oregon and Washington

Apply and incorporate 0.75 pint of Trilin per acre before planting and 0.75 pint after planting when potato plants have fully emerged from all soils, except do not apply to soils containing a 2% or more organic matter. Follow incorporation directions listed above for application to potato after planting.

### TRILIN WITH EPTAM TANK MIX in Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota and Texas

This tank mix will effectively control the following weeds in addition to those weeds controlled by Trilin.

Henbit (spring applications)	Oat, Wild
Nightshade Black	Ragweed Common
Nightshade Harry	Smartweed Pennsylvania
Nutsedge	Velvetleaf Buttonweed

Follow the recommendations for soil preparation and incorporation procedures for Trilin. Trilin with Eptam tank mix may be applied after planting but before crop emergence. In areas where potatoes are normally dragged off, this tank mix should be applied and incorporated up to or immediately following drag off.

### Broadcast Rates Per Acre

Trilin (pts.)	Coarse	Medium	Fine
Areas receiving less than 20" average annual rainfall	1	1-1.5	1-1.5
Areas receiving greater than 20" average annual rainfall PLUS	1	1-1.5	1-2
Eptam 7E	1.75-7**	1.75-7	1.75-7

\*\*For nutsedge control use the higher rate of Eptam 7E.

On soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and on soil with 5-10% organic matter use 2 pints on all soils.

### PRECAUTION

Follow direction and precautions on the Eptam label before using. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use foliage from treated plants for food forage or for grazing.

in or grass treated areas.

**L. METURON 80DF, COTORAN 4L OR izona)**

eparation and incorporation procedures for Meturon 4L, Meturon 80DF, Cotoran 4L or ols: all the annual grasses and broadleaf lus these additional weeds:

Morningglory	Sesbania
Tricky Sida (Teaweed)	Sicklepod
Tagweed	Smartweed
Triggrass	Tumbleweed

Coarse	Medium	Fine
1	1.5	2
2	3.125	4
1.25	1.9	2.5
2	3.125	4
1.2	2	2.4

partially filled tank of water. Add the Trilin oroughly mixed and continue filling. Agitate filling and application operations. Do not ithout constant agitation. If bypass agitation hanging the bypass line stop at the bottom gallons of water per acre.

plus Meturon or Cotoran on sandy, loamy Do not use on cotton planted in furrows.

**Mississippi Only.**

organic matter use 1.5 pints Meturon 4L or iron 80DF or Cotoran DF in tank mix with

crops other than cotton until 1 year after the sandy or coarse textured soils of less than

ated land is planted to anything but cotton ation of Trilin plus Meturon or Cotoran. Do otton plants or gin trash to livestock. Do not oran with liquid fertilizer.

**VED BY METURON 4L, METURON 80DF, N DF OVERLAY**

s recommended for the specific soil texture, 4L at 2 to 4 pints per acre or Meturon 80DF ounds per acre as a preemergence surface on light silt and sandy soil low in organic

mitations on the labels of each product used

**VED BY DIREX 4L OR KARMEX DF Mississippi River plus Arkansas, lsiana and Eastern Texas)**

ing. Follow with a preemergence application This will effectively control all the weeds these additional weeds.

Panngrass Velvetgrass

harvest.

**Incorporation**

Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary using a different direction for the second.

**Cultivation**

As some johnsongrass plants will escape, cultivation is necessary to obtain effective control. Effective control cannot be obtained with only 1 year of Trilin use.

**PRECAUTION**

Crop injury may result if the 2 quart rate is used on new plantings. Orchard or vineyards should not be interplanted with other crops. If the treated vineyards and orchards are diverted to other crop uses, plant only those crops for which Trilin has been registered as a preplant treatment for the next cropping season.

**Bindweed control In California**

Use Trilin for the control of field bindweed in vineyards and for almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees.

Use Trilin at a broadcast rate of 4 pints per acre on all soil textures. Trilin must be applied in the spring with a specially designed spray blade which applies a thin concentrated layer at a soil depth of 4 to 6 inches. This layer of Trilin prevents bindweed shoots from emerging.

**Land Preparation**

All weeds and grasses should be destroyed with soil tillage prior to applying Trilin. This tillage is necessary to prevent trash from interfering with the operation of the spray blade.

**Equipment**

A spray blade capable of running 4 to 6 inches below the surface of the soil should be used. The spray blade should be equipped with nozzles located under the blade and directed so that the Trilin spray will be trapped under the soil which is flowing over a blade as it is pulled through the soil. A sufficient number of nozzles should be used with spacing that will uniformly apply the Trilin underground in a thin horizontal layer.

**Application**

Use Trilin in 40 to 80 gallons of water per acre. Operate the spray blade at a depth of 4 to 6 inches.

**PRECAUTION**

After rainfall or irrigation some soils may crack as they dry. Bindweed may emerge if the cracks extend through the Trilin layer. Prevent or eliminate cracks by shallow disking or other tillage. Avoid deep tillage which disturbs the subsurface layer. Cultivation or tillage also aids the control of germinating seeds.

**IRRIGATION WATER RINGS NON-BEARING CITRUS TREES**

Apply to non-bearing citrus trees through irrigation water rings to provide preemergence weed control. Mix at a rate of 12 fluid ounces of Trilin per 500 gallons of water. Agitate until uniformly dispersed in tank.

Apply 10 gallons of the mixture per four foot diameter water ring per tree. Trilin should be applied at the second or third watering and should not be applied in combination with any other pesticide.

**GREENS**

**Turnip greens grown for processing, Collards, Kale and Mustard Greens**

Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

**HOPS**

On soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and on soil with 5-10% organic matter use 2 pints on all soils.

**PRECAUTION**

Follow direction and precautions on the Eptam label before using. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use foliage from treated plants for feed, forage or for grazing.

**TRILIN WITH EPTAM APPLICATION BEFORE PLANTING in Idaho, Oregon and Washington**

Trilin with Eptam may be applied prior to planting using a broadcast rate of 0.75 pint of Trilin per acre and 3.5 pints of Eptam 7E per acre on all soil textures. Incorporate immediately.

**PRECAUTION**

Do not use this tank mix both before and after planting in the same season. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use foliage from treated plants for feed, forage or for grazing.

**RADISH**

Apply and incorporate as a preplant soil treatment.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	1.5

**RAPESÉD (CANOLA), CRAMBE**

(Except In Alaska)

Follow recommended procedures for soil preparation and application of Trilin. Trilin may be applied in the fall or early spring prior to seeding. Set incorporation equipment to incorporate to a depth of 3 to 4 inches with equipment specified in this label.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2

**SAFFLOWER**

Recommended soil preparation application and incorporation procedures for Trilin should be followed. Use Trilin before planting in the spring or between October 15 and December 31.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

On coarse and medium soils use 1.5 pints per acre and 2 pints on fine soils with 2-5% organic matter. Use 2 to 2.5 pints on all soils with 5-10% organic matter.

**For Fall Application in Arizona, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming**

Apply and incorporate Trilin any time between October 15 and December 31. Ground may be left flat or bedded-up over winter. On bedded ground, beds should be knocked down to desired height before planting, moving some treated soil from tops into furrows where soil is left flat over winter. Care should be taken during spring bedding operations to prevent turning up untreated soil. Destroy established weeds during seedbed preparation. Before planting destroy weeds that become established in furrows due to uncovering of untreated soil. Apply and incorporate Trilin at a broadcast rate of 1.5 pints per acre on coarse soils, 2 pints on medium and 2.5 pints on fine soils. Trilin should not be applied in the fall to soils which are wet

Hagweed Wild Lettuce  
Shepherdspurse Wild Mustard

Coarse	Medium	Fine
1	1.5	2
0.5	1	1.75
0.33	0.67	1

mitations on the label of each product used  
rex 4L or Karmex DF should not be used on  
ic-matter as crop injury may result. Do not  
ts to livestock or graze treated areas.

**IN**  
pounds a.i./A (4 pints Trilin Herbicide) per  
more than 4 pints per crop year (either fall  
lication or preplant plus post plant through

## WN FOR PULPWOOD tonwood and Poplar

efore planting.

Coarse	Medium	Fine
1.0	1.25-1.5	1.5-2.0

matter use 1.5 to 2 pints. On soils with  
0 pints. Use lower rate in areas receiving  
fall and irrigation.

the soil and use incorporation methods not

### All Soil Texture 2.0-4.0

range may be adjusted according to weed

e application is necessary for satisfactory  
similar implement to bring rhizomes to the  
oil twice using a tandem disc to cut rhizomes  
and to destroy emerging johnsongrass.

### All Soil 4.0

twice with a tandem disc set to cut 4 to  
at 4 to 6 mph.

grass plants will escape. Timely cultivation  
not spraying with effective postemergence  
vel of johnsongrass control.

## UCURBITS cumber and Watermelon

ern United States including Texas

y to the soil between the rows and beneath  
true leaf stage.

Coarse	Medium	Fine
--------	--------	------

Apply and incorporate Trilin while the crop is dormant using a broadcast  
rate of 1 pint per acre on coarse soils, 1.25 pints on medium soils and  
1.5 pints on fine soils. Use incorporation equipment that will insure  
thorough soil mixing with minimum damage to the crop.

## KENAF

### Ground Application

Using a low-pressure herbicide sprayer which will apply the spray  
uniformly, apply Trilin in 5 to 40 gallons of water (broadcast spray) per  
acre.

### Aerial Application

For aerial spraying apply Trilin in 5 to 10 gallons of water. Pump pressure,  
nozzle arrangements, speed and height should be adjusted to provide a  
uniform application to the soil surface. To assure proper application spray  
widths use swath markers or flagmen.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2

### Incorporation Directions

#### General Directions

Erratic weed control and/or crop injury may result if directions are not  
followed. The soil surface should be free of existing weeds and excessive  
trash or clods.

#### Before Planting

Incorporation of Trilin must take place within 24 hours after application  
and may be accomplished in one pass with a sweep type or a rolling  
cultivator. Follow recommended soil incorporation procedures for Trilin.

### PRECAUTION

Without reducing the weed control activity of Trilin, soil treated with Trilin  
may be shallow cultivated, rotary hoed or handhoed. Do not cultivate  
deeper than (1 to 2 inches) the Trilin treated layer of soil as this may bring  
untreated soil to the surface and poor weed control may result. Kenaf  
treated with Trilin must not be used as a food, forage or feed.

## LUPINE

Apply and incorporate before planting.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils  
and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints.

## MUSTARD

Grown for seed or processing for food in Minnesota, Montana and North  
Dakota. Apply and incorporate Trilin prior to planting at 1 pint per acre on  
coarse soils and 1.5 pints on medium and fine soils.

## NUT CROPS

See FRUIT AND NUT CROPS AND VINEYARDS for instructions.

## OKRA

Apply and incorporate Trilin before or immediately after planting. If applied  
and incorporated after planting, set equipment so as to not disturb the  
seed.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			

## SOUTHERN PEA

Apply and incorporate Trilin prior to planting only.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium  
and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints  
on all soils.

## SOYBEANS

### Preemergence

Follow recommended soil preparation, application, and incorporation  
procedures for Trilin.

### Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2

Use 1.5 pints per acre on coarse and medium textured soils and 2 pints  
on fine soils with 2-5% organic matter. Use 2 to 2.5 pints on all soils with  
5-10% organic matter.

### Fall Application

Trilin may be applied and incorporated any time between October 15  
and December 31. The ground may be left flat or bedded-up over winter.  
Where soil is left flat, care should be taken not to turn up untreated soil  
from beds into furrows. On bedded grounds, beds should be knocked  
down to desired heights before planting, moving some treated soil into the  
furrow from the beds. Established weeds should be destroyed during  
preparation of seedbed. Before planting, destroy weeds which may have  
become established in furrows due to the uncovering of untreated soil  
during bedding. Trilin should not be applied to wet soil or soils which are  
subject to prolonged periods of flooding of ground where rice was grown  
the previous year.

Apply and incorporate Trilin at a broadcast rate of 2 pints per acre on  
coarse and medium soils and 2.5 pints on fine soils in Alabama, Arkansas,  
northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri  
bootheel, North Carolina, Oklahoma, South Carolina, Tennessee and  
Texas.

For other states where soybeans may be grown, use Trilin at a broadcast  
rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and  
2 pints on fine soils. For coarse soils with 2-5% organic matter use  
1.5 pints. For soils with 5-10% organic matter use 2 to 2.5 pints.

## SPECIAL USE DIRECTIONS FOR SOYBEANS

### Fall Panicum

Apply and incorporate Trilin broadcast at the rate of 2 pints per acre on  
coarse and medium soils.

### Crop Rotation

Plant only rice and those crops for which Trilin can be applied as a  
preplant treatment following a double rate treatment season or injury may  
result.

### Texas Panicum

Apply and incorporate at the labeled use rate for the soil type. Uniformly  
incorporate to a depth of at least 3 inches. Adequate soil moisture is  
essential for herbicide activation. Any deviation from these  
recommendations can result in reduced control.

### Pigweed and Seedling Johnsongrass Control

In Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi,



1 1.25-1.5 1.5  
 1 1.5 2  
 matter use 1.5 pints on coarse and medium soils with 5-10% organic matter use 2 pints

**GRAIN SORGHUM AND FORAGE AND SILAGE**

ain sorghum (8 inches or taller) as an over-actively control weeds listed for Trilin. Trilin may or as a directed spray in field corn and rattercane in addition to those other weeds

ation to insure loose, friable soil, to remove er the base of plants with soil.

incorporated at the recommended rates for is well established (8 inches or taller). Trilin over-the-top spray or as a directed spray. If foliage prevents uniform coverage of soil ay be accomplished with only one pass of a perly adjusted rolling cultivator. The sweep- to 5 sweeps per row, middle and be operated sweeps so as to avoid exposing untreated tools to prevent crop injury.

Coarse	Medium	Fine
0.75-1	1-1.5	1.5-2

Alabama, Florida, Georgia, North Carolina, control fall panicum and Texas panicum.

**NK MIX**  
 atrazine for additional weed control in field

Coarse*	Medium	Fine
0.75-1	1-1.5	1.5-2

2.4 4.75 6  
 the rates listed above. For other atrazine rates. When using AAtrex NineO 1 pint of id of NineO. One pint of AAtrex 4L equals

nix on coarse soils for grain sorghum.  
 lin/Atrazine tank mix as directed on the Trilin sorghum.

orn or corn grown for seed. Do not apply to rt and preemergence treatment or crop injury utions and limitations on the labels of each nd overlays.

**FLAX**

orporated in the fall for weed control in spring om existing weeds or previous crop should l so that there is no interference with

per acre of 1 pint on coarse soil, 1.5 pints on

Areas receiving greater than 1 1.5 2  
 20" average annual rainfall  
 For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

**PEA  
 Dry and English**

**Trilin Alone**  
 Apply and incorporate Trilin prior to planting at a rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils.

**Fall application in Idaho, Oregon and Washington**  
 Apply and incorporate Trilin any time between October 15 and December 31 using a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils and 1.5 pints on fine soils. Destroy established weeds during seedbed preparation. Do not apply Trilin in the fall to soils which are wet or are subject to prolonged periods of flooding.

**TRILIN WITH FAR-GO TANK MIX in Idaho, Oregon and Washington**  
 Trilin plus Far-Go controls wild oat in addition to other annual grasses and broadleaf weeds controlled by Trilin.

**Application Rates**  
 Broadcast 0.75 pint of Trilin per acre on coarse and medium soils, 1 pint on fine soils. Use 1.25 quarts of Far-Go per acre for all soil textures.

**Incorporation Directions**  
 Apply and incorporate up to 3 weeks prior to planting. Follow recommended incorporation procedures for Trilin.

**PRECAUTION**  
 Do not apply to lentils. Leaf crinkling and delayed maturity of peas may occur particularly on clay points in the northwest; but this is usually more than offset by a reduction of wild oat. Do not use foliage from treated plants for feed or grazing. Observe all precautions and limitations on the label of each product used in tank mixes and overlays.

**PEANUT  
 Spanish Peanut, Florunner and Florigiant  
 in New Mexico, Oklahoma and Texas**

Apply and incorporate Trilin prior to planting, at planting or immediately after planting using a broadcast rate of 1 pint per acre on coarse soils. When incorporating after planting take care not to disturb the seed.

**PEPPER  
 Transplant**

Apply and incorporate Trilin prior to transplanting only.  
**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2
For soils with 2-5% organic matter use 1.5 pints on coarse and medium, 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.			

of 1 to 1.5 pints per acre on coarse soils, 1.5 to 2 pints on medium soils and 2 pints on fine soils. **Exception:** Use 3 pints per acre on fine soils in Louisiana.

**Additional Weed and Grass Control**  
 In the Texas Gulf Coast counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, apply Trilin at a broadcast rate of 1.5 pints per acre on coarse soils, 2 pints on medium soils and 3 pints on fine soils, up to two weeks before planting.

**Charcoal Soils in Arkansas, Louisiana and Mississippi**  
 Freshly cleared land sometimes contains 5-10% organic matter and charcoal from burning debris. This charcoal and organic matter has a tendency to bind Trilin and reduce its weed control activity. If these conditions exist, higher rates of Trilin are needed for weed control. Crop injury can occur if increased rates are used and the charcoal or organic matter is not present in the soil. In the burn row a high level of charcoal is present, consequently, poor weed control may result even with an increased rate of Trilin.

Use Trilin broadcast at the rate of 1.5 to 2.5 pints per acre on coarse soils, 2.5 pints on medium soils and 3 pints on fine soils. Follow recommended soil preparation, application and incorporation procedures for Trilin:

**Red Rice in Arkansas, Louisiana, Mississippi and Texas only**  
 Partial suppression or control of red rice can be obtained when Trilin is applied at the following recommended rates.

**Broadcast Rates Per Acre**

Application	Coarse	Medium	Fine	Coarse with 2-5% organic matter	Coarse with 5-10% organic matter
Year 1	2	3	4	3	4
Year 2	1	1.5	2	1.5	2-2.5

If high organic matter and/or charcoal are present in the soil use Trilin the second year as follows for Arkansas, Louisiana and Mississippi:

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5-2.5	2.5-3

**Crop Rotation**  
 Use a two year program for red rice control in soybeans. Use rates listed for 1 year and plant soybeans. The second year plant only those crops for which Trilin can be used preplant using the normal rates listed for your soil type and charcoal level. Do not plant rice the second year. Rice may be planted the third year.

**Rhizome Johnsongrass in Eastern United States and Texas**  
 Rhizome johnsongrass can be acceptably controlled using a double rate program for two consecutive years as follows.

**Soil Preparation**  
 For satisfactory results proper soil preparation is essential. To bring rhizomes to the top of the soil use a chisel plow or similar implement. Follow twice with a disc prior to application to cut rhizomes into small 2 to 3 inch pieces and to destroy any emerged johnsongrass.

**Application**  
 Choose one of the following programs which best meets your cultural practices.

For spring application use Trilin prior to planting in the spring for two consecutive years. A broadcast rate of 2 pints per acre should be used on coarse soils, 3 pints on medium soils and 4 pints on fine soils. Use 3 pints on coarse soils with 2-5% organic matter and 4 pints on soils with 5-10% organic matter.

**OR**  
 For fall application use Trilin between October 15 and December 31 for two consecutive years using the same rates as for spring application.

**OR**  
 A split application of Trilin may be used in spring and fall for

**POTATO**

**es Except Maine)**

after planting prior to emergence or before or after the potato plants have fully emerged.

Coarse	Medium	Fine
1	1.25-1.5	1.5
1	1.5	2

After use 1.5 pints on coarse and medium soils with 2-5% organic matter.

Trilin should be set to uniformly cover the bed and not be concentrated over the top. If the herbicide is concentrated over the top, retarded and stem brittleness can occur. Do not apply Trilin to treated soil when Trilin is applied to plants have fully emerged. Do not use subsequent cultivations. Be careful that Trilin does not damage potato seed pieces or tubers.

**Oregon and Washington**

Use 1 pint of Trilin per acre before planting and 1.5 pints per acre after emergence. Potato plants have fully emerged from all soils containing a 2% or more organic matter. Listed above for application to potato after emergence.

**MIX in Kansas, Minnesota, Nebraska, North Dakota and Texas**

Control the following weeds in addition to those listed above:

- Oat, Wild
- Ragweed Common
- Smartweed Pennsylvania
- Velvetleaf Buttonweed

For soil preparation and incorporation in Eptam tank mix may be applied after emergence. In areas where potatoes are planted, Trilin should be applied and incorporated up to 2 weeks before planting.

Coarse	Medium	Fine
1	1-1.5	1-1.5
1	1-1.5	1-2

Use 75-7\*\* or higher rate of Eptam 7E.

Use 1.5 pints per acre on coarse and 2 pints on all soils with 2-10% organic matter.

Read the Eptam label before using. Observe the labels of each product used to remove foliage from treated plants for disposal.

**Broadcast Rates Per Acre**

Application	Coarse	Medium	Fine	Coarse with 2-5% organic matter	Coarse with 5-10% organic matter
Trilin (pts.)					
Spring & Fall	1	1	2	1.5	2

**Incorporation**

For good rhizome johnsongrass control deep incorporation is necessary. Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two passes are necessary with the second pass in a different direction from the first.

**Cultivation**

To control johnsongrass plants which have escaped Trilin, timely cultivations during the crop season may be necessary to obtain control. Control cannot be obtained with only 1 year of double rate Trilin use.

**Crop Rotation**

Plant only rice and those crops for which Trilin can be applied as a preplant treatment following a double rate treatment season or injury may result.

**Wild Cane (Shattercane)**

Follow recommended soil preparation and application procedures for Trilin.

Germination of wild cane (shattercane) occurs throughout the growing season and from deeper in the soil than most other weed seeds. Effective control can be obtained by using the following increased rates of Trilin.

A broadcast rate of 1 pint per acre on coarse soils, 2 pints on medium soils and 2.5 pints on fine soils.

**Incorporation**

For good wild cane control deep incorporation is necessary. Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two passes are necessary with the second pass in a different direction from the first.

**Cultivation**

Cultivation during the crop season will also contribute to control.

**PRECAUTION**

When using higher rates, soybeans should be planted after early season adverse weather conditions. This is to avoid additional stress to the plants due to cool, wet weather early in the growth period. Use of high rates and adverse growing conditions can cause reduced stands, delayed maturity and reduced yields.

**TANK MIXES AND OVERLAYS**

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

**TRILIN WITH SENCOR OR LEXONE**

Trilin with Sencor or Lexone will control certain grasses and broadleaf weeds listed for Trilin alone and the following weeds:

- Jimsonweed
- Mallow, Venice
- (Flower-of-an-hour)
- Mustard, Wild
- Prickly Sida
- Ragweed, Common
- Sesbania, Hemp
- Smartweed, Pennsylvania
- Velvetleaf

Cocklebur, morningglory and giant ragweed (horseweed) control may be erratic. Timely cultivation may improve control. An overlay of Sencor or Lexone may be preferred to the tank mix where cocklebur is a serious problem.

Trilin tank mixed with Sencor or Lexone may be applied from 2 weeks prior to planting up to planting.

**Broadcast Rates Per Acre**

Coarse	Medium	Fine
--------	--------	------

**Seeding Directions**

Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone of soil into which Trilin has been incorporated.

**PRECAUTION**

Crop injury such as delayed emergence and development may occur when wheat is planted in direct contact with treated soil.

**Fallow soil application in Washington and Oregon**

To control cheatgrass and certain annual grasses and broadleaf weeds, apply Trilin and shallowly incorporate into fallow soil up to four months ahead of planting. As long as the seed is placed below the zone of soil treated with Trilin the growth development or yield will not be adversely affected. Use deep or semi-deep furrow drills. Use at a broadcast rate of 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. Trilin can be applied any time from May to September prior to the fall planting of winter wheat.

**Incorporation**

Incorporate Trilin using a flexible tine-tooth harrow (Flextine or Melroe) set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Two passes over the field in different directions are necessary for thorough incorporation. Incorporate one time within 24 hours after application followed by a second incorporation prior to seeding. When a flexible tine harrow has been used to apply Trilin do not till the soil with a disc.

**PRECAUTION**

Deep furrow or semi-deep furrow drills only should be used. Seed should be placed below the zone of soil into which Trilin has been incorporated. Injury to the crop or delay in emergence or development may occur if wheat is planted directly into the zone of soil treated with Trilin.

**WHEAT (SPRING), DURUM AND BARLEY**

**Preplant Barley**

Apply and incorporate Trilin before planting in the spring or in the fall after September 1, at the rate of 1 pint per acre on coarse soils, 1.5 pints per acre on medium soils and 1.5 pints on fine soils. Use 1.5 pints of Trilin per acre on coarse and medium soils with 2-5% organic matter. Incorporate Trilin into the soil with a flexible tine-tooth harrow, (Flextine or Melroe) set to cut 1 to 2 inches deep and operate at 3 to 6 mph. Within 24 hours after application, incorporate one time followed by a second incorporation in a different direction from the first, prior to planting. After the Trilin has been incorporated with a flexible tine harrow do not till the soil with a disc. Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone of soil into which Trilin has been incorporated.

**PRECAUTION**

Crop injury such as delayed emergence and development may occur when barley is planted in direct contact with treated soil. DO NOT MAKE MORE THAN ONE TRILIN APPLICATION PER SEASON.

**Postemergence Wheat (Spring), Durum and Barley**

**Trilin Alone**

To control foxtail (pigeongrass) Trilin is recommended as a postplant incorporated treatment. Use Trilin at a broadcast rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils. Seedbed should be well-tilled and seed planted 2 to 3 inches deep. Use Trilin after seeding but prior to crop emergence. Use flextine or diamond harrow operated two times in different directions to incorporate. Incorporate by operating equipment at a speed of at least 5 mph and set at up to 1.5 inches deep. Apply and incorporate the first time in the same operation if possible. Both incorporations must be done with 24 hours.

**TRILIN WITH FAR-GO TANK MIX**

This tank mix will effectively control all the weeds controlled by Trilin alone plus these additional weeds:

- Foxtail
  - Pigeongrass
  - Wild Oat
- Apply Trilin with Far-Go as a postplant incorporated treatment.

3/1/81

**CAUTION BEFORE PLANTING in Idaho.**

and prior to planting using a broadcast rate of 3.5 pints of Eptam 7E per acre on all soil types.

before and after planting in the same areas and limitations on the labels of each product overlays. Do not use foliage from treated plants for feed or forage.

**RADISH**

plant soil treatment.

Coarse	Medium	Fine
1	1.5	1.5

**CANOLA), CRAMBE**

(pret in Alaska)

Use for soil preparation and application of the fall or early spring prior to seeding. Set furrows to a depth of 3 to 4 inches with a disk.

Coarse	Medium	Fine
1	1.5	2

**FLOWER**

Use application and incorporation procedures listed for Trilin before planting in the spring or after March 31.

Coarse	Medium	Fine
1	1.25-1.5	1.5
1	1.5	2

Use 1.5 pints per acre and 2 pints on fine soils. Use 2 to 2.5 pints on all soils with 5-10% organic matter.

**Idaho, Montana, Nevada, Oregon, Washington**

Apply between October 15 and December 15. On bedded ground, apply to desired height before planting, moving furrows where soil is left flat over winter. Avoid bedding operations to prevent turning wind-blown weeds during seedbed preparation. Do not become established in furrows due to soil applied and incorporate Trilin at a broadcast rate on coarse soils, 2 pints on medium and 2.5 pints on fine soils applied in the fall to soils which are wet or prone to flooding.

**HERN PEA**

Use prior to planting only.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Sencor 4L (pt.)	0.5	0.75	1
OR			
Lexone DF or Sencor DF (lb.)	0.33	0.5	0.67

Do not use Sencor or Lexone on coarse soils with less than 1% organic matter.

**PRECAUTION**

Do not plant any crop other than soybeans within 4 months after treatment. Crop injury, herbicide residue or erratic weed control may result from over application, uneven application or improper soil incorporation. Cold weather, deep planting, seedling disease, excessive moisture, soil pH over 7.5, high salt concentration or drought are additional stress factors. Any of these may weaken crop seedlings and possible damage from the tank mix is increased. These factors may also delay crop development or reduce yields when Sencor or Lexone is applied. Do not use foliage from treated plants for feed or forage.

**TRILIN PREPLANT FOLLOWED BY SENCOR OR LEXONE AS AN OVERLAY**

After Trilin has been applied as a preplant incorporated herbicide make a single application of Sencor or Lexone as either a broadcast or band spray either during planting or after planting before the soybeans emerge. Crop injury may result if Sencor or Lexone is sprayed over the top of emerged soybeans.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Sencor 4L (pts.)	0.75-1	0.75-1.5	1-1.75
OR			
Sencor DF (lbs.)	0.5-0.67	0.5-1	0.67-1.17
OR			
Lexone DF (lb.)	0.5	0.5-0.67	0.67

Lexone or Sencor should not be applied to sands or soils with less than 1/2% organic matter or to coarse soils (sandy loam and loamy sand) containing less than 2% organic matter.

**PRECAUTION**

Tracy, Semmes, Altona, Vansoy or Coker 102 soybean varieties are sensitive and crop injury may result if Lexone or Sencor is used. Before a Sencor or Lexone application, seeds must be planted at least 1.5 inches but not more than 2 inches below the soil surface. Only one application per season should be used at these rates. Do not plant areas treated with Sencor or Lexone to any crop other than soybeans within 4 months after treatment. Soybean injury may occur if Lexone or Sencor is used on soils having a calcareous surface, a pH of 7.5 or higher, or if they are used in conjunction with soil-applied organic phosphate pesticides. Do not use foliage from treated plants for feed or forage. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

**SUGAR BEET**

Use Trilin when plants are between 2 to 6 inches tall as a broadcast over-the-top spray at a rate of 1 pint per acre on coarse soils and 1.25 to 1.5 pints on medium and fine soils. Use the higher rate for medium and fine soils in areas receiving greater than 20" average annual rainfall. Set incorporation machinery to throw treated soil toward the plants in the row. Care should be taken that incorporation machinery does not damage the sugarbeet taproot.

**PRECAUTION**

To reduce the possibility of girdling, exposed beet roots should be covered

Apply Trilin with Far-Go as a postplant incorporated treatment. Plant 2 to 3 inches deep in a well-tilled seedbed. Trilin with Far-Go should be applied after seeding but prior to crop emergence. Use flexline or diamond harrows to incorporate. Make two passes each in different directions at speeds of at least 5 mph operating equipment 1 to 1.5 inches deep. Application and the first incorporation should be done in the same operation if possible. If not incorporate immediately after application.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pts.)			
Barley, Durum, Spring Wheat	1	1	1.5
PLUS			
Far-Go (pts.)			
Durum, Spring Wheat	2.5	2.5	2.5
Barley	2	2	2

**PRECAUTION**

Over application may result in crop injury. Observe all precautions and limitations on the label of each product used in tank mixes and overlays. DO NOT MAKE MORE THAN ONE TRILIN APPLICATION PER SEASON.

**ORNAMENTALS**

Apply and mechanically incorporate Trilin prior to planting new nursery stock liners, ornamentals, trees and woody shrubs and gladioli. Gladioli corms less than 1 inch in diameter may be injured by pre-plant application. Trilin may also be applied to these and other listed ornamentals (see below) after they are established. When mechanically incorporated after planting, the implement should be adjusted so that the treated soil is thrown toward and around the plants in the row.

**Broadcast Rates Per Acre**

	Coarse	Medium	Fine
Trilin (pints)	1.0	1.5	2.0

For the indicated ornamental groundcovers, apply 1 gallon/acre (3 oz/1000 sq ft) of Trilin in 5 to 40 gallons of water and incorporate within 24 hours with at least a 1/2 inch rain or its equivalent in sprinkler irrigation.

Woody	Trees	Ground Cover
Andromeda, Japanese	Almond	Aaronsbeard
Arborvitae, Americana	Apple, Crabapple	Bellflower, Adriatic
Azalea	Apricot	Bellflower, Poscharsky
Barberry, Japanese	Ash, White	Ceanothus
Barberry, Mento	Baldcypress	Cereopsis
Boxwood, Common	Birch, European White	Cotoneaster
Boxwood, Harlands	Black gum	Coyote Brush
Boxwood, Littleleaf	Cherry	Crown Vetch
Camellia, Japanese	Chestnut, Chinese	Daisy Trailing African
Camellia, Sasanqua	Cottonwood	Fern, Asparagus
Cherrylaurel, Americana	Dogwood, Flowering	Gazania
Cinquefoil	Dogwood, Kousa	Germander
Cleyera, Japanese	Douglasfir	Ice Plant, Largeleaf
Cotoneaster, Cranberry	Fir, Balsam	Ivy, Algerian
Cotoneaster, Zabel	Hemlock, Canada	Ivy, English
Deutzia	Honeylocust	Lily-of-the-Nile
Elaeagnus, Silverberry	Larch, Japanese	Lilyturf, Bigblue
Euonymus, Spreading	Locust, Black	Marigold
Euonymus, Winged	Maple, Norway	Myoporum
Euonymus, Wintercreeper	Maple, Red	Plumbago, Dwarf

2/27

Coarse	Medium	Fine
1	1.25-1.5	1.5
1	1.5	2

For coarse and medium soils use 1.5 pints on coarse and medium soils with 5-10% organic matter use 2 pints

## SOYBEANS

Preparation, application, and incorporation

Coarse	Medium	Fine
1	1.5	2

For coarse and medium textured soils use 2 pints on coarse soils with 2 to 2.5 pints on all soils with

Incorporated any time between October 15 and November 15 may be left flat or bedded-up over winter. Do not be taken not to turn up untreated soil on bedded grounds, beds should be knocked during planting, moving some treated soil into the bedded weeds should be destroyed during planting, destroy weeds which may have due to the uncovering of untreated soil it be applied to wet soil or soils which are flooding of ground where rice was grown

Use a broadcast rate of 2 pints per acre on 5 pints on fine soils in Alabama, Arkansas, Louisiana, Mississippi, southeastern Missouri, Oklahoma, South Carolina, Tennessee and

Soybeans may be grown, use Trilin at a broadcast rate of 2 pints on coarse soils, 1.5 pints on medium soils and 1 pint on fine soils with 2-5% organic matter use 2 to 2.5 pints.

## INSTRUCTIONS FOR SOYBEANS

Broadcast at the rate of 2 pints per acre on

Conditions for which Trilin can be applied as a double rate treatment season or injury may

Recommended use rate for the soil type. Uniformly to a depth of 3 inches. Adequate soil moisture is essential for activation. Any deviation from these conditions will reduce control.

### Preplant Grass Control

In Alabama, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, use Trilin preplant at a broadcast rate of 2 pints on coarse soils, 1.5 to 2 pints on medium soils and 1 pint on fine soils. **Application:** Use 3 pints per acre on fine soils in

### PRECAUTION

To reduce the possibility of girdling, exposed beet roots should be covered with soil before applying Trilin.

In Colorado, Idaho, Montana, Nebraska, Oregon, Texas, Utah, Washington and Wyoming use a tine-tooth harrow (Flexline or Melroe) for incorporation of Trilin for effective weed control in sugar beets. The tine-tooth harrow should be operated two times over the field, the second time in opposite direction at a speed of 3 to 6 mph. The harrow should be set to cut 1 or 2 inches deep. Care should be taken that the tine-tooth harrow does not damage the sugar beet taproot. All recommendations for application procedures and broadcast rates per acre for sugar beet should be followed.

## SUGARCANE Plant Cane

Apply and incorporate Trilin twice a year at a broadcast rate of 2 to 4 pints per acre for all soil textures. Make the first application of Trilin in the fall immediately after the seed pieces are planted and the second application of Trilin in the spring before or shortly after the cane emerges. Before the spring application rain-packed beds should be loosened 2 to 3 inches deep. Care should be taken so that the seed pieces or emerging shoots are not damaged by incorporation machinery.

### Hawaii (postplant) for control of most annual grasses including guineagrass

Surface apply Trilin after planting (for plant cane) or after harvesting (for ratoon cane) at a broadcast rate of 6 to 8 pints per acre for all soil textures. Apply Trilin prior to germination and emergence of grass weeds. For best results in plant cane, the soil surface should be smooth and finely tilled. Apply Trilin as soon as possible after tillage and planting, before germination and emergence of grass weeds. For optimum efficacy in ratoon cane, minimize surface residue from previous crop before applying. For best result, apply Trilin before anticipated rainfall in non-irrigated and furrow-irrigated sugarcane. Apply 0.5 inch or more irrigation in drip-irrigated or sprinkler-irrigated sugarcane as soon as possible after applying Trilin.

### Repeat Applications

Subsequent germination flushes of grass weeds may occur prior to the development of a full dense canopy of sugarcane. Once this occurs, additional grass weed establishments is strongly suppressed. One or two additional applications of Trilin can be applied to maintain weed control during the early crop development period. For repeat applications, direct the spray to the soil surface to minimize interception of the herbicide by the crop.

### PRECAUTION

Do not apply Trilin as a postplant surface applied treatment within 180 days of harvest.

### Applications in Louisiana or Texas up to 10 days before layby for plant cane or ratoon cane

Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures. This should be done in the spring from before or shortly after the cane emerges up to layby. Use Trilin after the beds have been shaved or false shaved. Rain-packed beds should be loosened 2 to 3 inches deep before application. Care should be taken that seed pieces or emerging shoots are not damaged by incorporation machinery. A rolling cultivator or bed chopper may be used to incorporate Trilin layby applications in sugarcane on all soil textures. Normal incorporation directions for the rolling cultivator should be followed. Bed chopper should be set to cut 3 to 4 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary.

### Itchgrass (Raouigrass) control in Louisiana

Apply and incorporate Trilin on either plant or ratoon cane at a broadcast rate of 4 pints per acre for all soil textures. Directions above for sugarcane layby application in Louisiana and Texas should be followed.

Eucalyptus, wintercreeper	maple, Red	Flamingo, Dwarf
Firethorn	Maple, Silver	Rockrose
Forsythia	Maple, Sugar	Rosemary
Guava, Pineapple	Oak, Pin	Rupturewort
Holly	Oak, Red	Snow-in-Summer
Honeysuckle	Oak, Scarlet	Speedwell
India hawthorn	Peach	St. Johnswort
Juniper	Pine, Austrian	Stoncrop (Sedum)
Laurel, Mountain	Pine, Easter White	Strawberry, Beach
Lilac, Common	Pine, Japanese Black	Thrift
Mockrange	Pine, Loblolly	Verbena
Pittosporum, Japanese	Pine, Red	Wirevine, Creeping
Privet	Pine, Scotch	Yarrow, Woolly
Redcedar, Eastern	Planetree, London	Zoysiagrass
Rhododendro	Plum	
Spiraea, Vanhoutte	Red, bud, Eastern	
Virburnum	Spruce, Colorado	
Weigela	Spruce, Norway	
Willow	Spruce, White	
Yew, Anglojap	Sweetgum	
Yew, Japanese	Sycamore	
Yewpine	Tuliptree	
	Walnut Black	

### Roses and Other Established Flowers

African Daisy	Marigold
Aster (perennial)	Marigold, Cape
Balsam	Morningglory
Black-eyed Susan	Nasturtium
Calendula	Petunia
Carnation	Phlox
Entaurea, Velvet	Pincushion flower
Chrysanthemum	Poppy, California
Coreopsis	Portulaca
Cornflower	Rose
Cosmos	Salvia
Dahlia	Shasta Daisy
Dianthus	Snapdragon
Dusty Miller	Snow-on-the-mountain
Floss Flower	Stock
Forget-me-not	Sunflower
Four O'Clock	Sweet Allosome
Gaillardia	Sweet Pea
Gladiolus	Sweet Sultan
Golden Glow	Sweet Williams
Impatiens	Vinca
Ixora	Yarrow
Lobelia	Zinnia
Lupine	

## UNDER PAVED SURFACES

General Use Instructions and Site Preparation: Trilin should be used only where the soil to be treated has been prepared according to good construction practices. If rhizomes, stolens, tubers or other vegetative plant parts are present in the site, they should be removed by scalping with a grader blade to a depth sufficient to insure their complete removal.

on, Jefferson, Liberty, Matagorda, Orange, apply Trilin at a broadcast rate of 1.5 pints on medium soils and 3 pints on fine soils, 3.

**Louisiana and Mississippi**

Soils contains 5-10% organic matter and This charcoal and organic matter has a reduce its weed control activity. If these Trilin are needed for weed control. Crop types are used and the charcoal or organic matter. In the burn row a high level of charcoal weed control may result even with an

of 1.5 to 2.5 pints per acre on coarse soils, 3 pints on fine soils. Follow recommended incorporation procedures for Trilin:

**Louisiana, Mississippi and Texas only**

of red rice can be obtained when Trilin is needed rates.

	Coarse with 2-5% organic matter	Coarse with 5-10% organic matter
Coarse	4	3
Fine	2	1.5
Medium	3	4
Fine	5	2-2.5

Charcoal are present in the soil use Trilin the rates, Louisiana and Mississippi:

Coarse	Medium	Fine
1	1.5-2.5	2.5-3

of rice control in soybeans. Use rates listed The second year plant only those crops for it using the normal rates listed for your soil t plant rice the second year. Rice may be

**Western United States and Texas**

is acceptably controlled using a double rate rates as follows.

For soil preparation is essential. To bring Trilin use a chisel plow or similar implement. application to cut rhizomes into small 2 to 3 inches emerged johnsongrass.

Programs which best meets your cultural

Trilin prior to planting in the spring for broadcast rate of 2 pints per acre should be used on medium soils and 4 pints on fine soils. On soils with 2-5% organic matter and 4 pints on soils

between October 15 and December 31 for the same rates as for spring application.

Trilin may be used in spring and fall for the following rates.

Use Trilin in the spring or in the fall between October 15 and December 31. Recommendations for soil preparation application and incorporation procedures for Trilin should be followed.

**Broadcast Rates Per Acre**

Trilin (pts.)	Coarse	Medium	Fine
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

On coarse and medium soils use 1.5 to 2 pints and 2 pints on fine soils with 2-5% organic matter. Use 2 pints on all soils with 5-10% organic matter.

**TOMATO**

For direct-seeded tomato use Trilin at blocking or thinning as a directed spray to the soil between rows and beneath the plants and incorporate. For transplant tomato, apply and incorporate Trilin preplant, postplant or at layby.

**Broadcast Rates Per Acre**

Trilin (pts.)	Coarse	Medium	Fine
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1-2
Areas receiving greater than 20" average annual rainfall	1	1.5	2

On soils with 2-5% organic matter use 1.5 pints on coarse and medium textured soils and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints.

**VINEYARDS**

See FRUIT AND NUT CROPS AND VINEYARDS for instructions.

**WHEAT (WINTER)**

Colorado, Idaho, Kansas, Montana, Nebraska, Oregon, Washington and Wyoming

Apply Trilin for preplant preemergence control of cheatgrass and other annual grasses and broadleaf weeds controlled by Trilin. If the seed is placed below the zone of soil treated with Trilin the growth development and yield of winter wheat will not be adversely affected.

Use Trilin any time during a period from 3 weeks up to immediately prior to planting. Apply Trilin at a broadcast rate of 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils.

**Incorporation**

Incorporate Trilin into the soil with a flexible tine-tooth harrow (Flexline or Melroe) set to cut 1 to 2 inches deep and operate at 3 to 6 mph. Within 24 hours after application incorporate one time followed by a second incorporation in a different direction from the first prior to planting. After the Trilin has been incorporated with a flexible tine harrow do not till the soil with a disc.

and do not apply Trilin to soil where application is to be applied directly on top of the treated soil. Paving should follow Trilin application as soon as possible.

**Large Areas:** Apply Trilin in sufficient water to insure thorough wetting of the soil surface or penetration of the spray solution through the base rock layer. A minimum of 150 gallons per acre is recommended. Apply uniformly with a ground sprayer. Using a ground sprayed add Trilin to clean water during filling of spray tank. Agitate before spraying.

**Small Areas:** For treating small areas, a hand sprayer or sprinkling can may be used. Before application determine the amount of water and Trilin necessary to uniformly cover the area to be treated. Shake or stir the spray solution prior to application.

	Ounces/1000 Sq. Ft.	Gallons/Acre
Trilin	9.0-12.0	3-4

**WARRANTY STATEMENT**

TRICORP and SELLER OFFER THIS PRODUCT AND THE BUYER AND USER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to take into account all variables and to eliminate all risks associated with its use. Except as may be otherwise provided in any controlling State Law, injury or damage may result due to conditions that are beyond the control of TRICORP or the Seller. TRICORP warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use when this product is used as directed under normal conditions. **TRICORP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.** In no case shall TRICORP or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. The exclusive remedy for any buyer of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling or application of this product, including claims based on contract, negligence, strict liability or legal theories or otherwise shall not exceed the purchase price paid for this product or, at the election of TRICORP, the replacement of the product.

Tri Corporation, Houston, Texas 770042

TCN 101905-US  
CPC 091201

Caparol is a registered trademark of Syngenta

Cotoran is a registered trademark of Makhteshim Agan of North America, Inc.

Cotton Pro is a registered trademark of Griffin Corporation

Direx 4L is a registered trademark of Griffin Corporation

Eptam is a registered trademark of Gowan Company

Far-Go is a registered trademark of Gowan Company

Karmex is a registered trademark of E.I. Dupont de Nemours and Company

Meturon is a registered trademark of Griffin Corporation

Sencor is a registered trademark of Bayer

Trilin is a registered trademark of TriCorp.

3/12