

67979-8

12/5/2008

1/12

DEC 05 2008

Ms. Demetra Vlachos  
Senior Regulatory Affairs Manager  
Syngenta Seeds, Inc.  
P.O. Box 12257  
Research Triangle Park, NC 27709-2257

Dear Ms. Vlachos:

Subject: Your May 19, 2008 Amendment Request to Amend the Insect Resistance Management Terms and Conditions for Bt11 x MIR604 Corn  
EPA Registration No. 67979-8

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the following comments.

**Insect Resistance Management:**

The listed IRM terms and conditions are being modified as follows.

- 1) Under 6a. Refuge Requirements, the following statement is added "These refuge requirements do not apply to seed propagation of inbred and hybrid corn seed corn up to a total of 20,000 acres per county and up to a combined U.S. total of 250,000 acres per PIP active ingredient per registrant per year." Additionally, the phrase ", and preferably 6" is deleted.
- 2) Under 6d. Insect Resistance Monitoring, the Cry1Ab section is removed and replaced by the following and the Cry1Ab section under 6e. Remedial Action Plans is being removed.

**Cry1Ab Insect Resistance Monitoring**

The Agency is imposing the following conditions for this product:

The registrant will monitor for resistance to its lepidopteran-resistant Bt corn. The monitoring program shall consist of two approaches: (1) focused population sampling and laboratory testing; and (2) investigation of reports of less-than expected control of labeled insects. Should field-

CONCURRENCES

SYMBOL	7511P	7511P						
SURNAME	Madh	Smith						
DATE	12/4/08	12/4/08						

relevant resistance be confirmed, an appropriate resistance management action plan will be implemented.

**(1) Focused Population Sampling**

The registrant shall annually sample and bioassay populations of the key target pests *Ostrinia nubilalis* (European corn borer; ECB), *Diatraea grandiosella* (Southwestern corn borer; SWCB), and *Helicoverpa zea* (corn earworm; CEW). Sampling for the target pests will be focused in areas identified as those with the highest risk of resistance development (e.g., where lepidopteran-active Bt hybrids are planted on a high proportion of the corn acres, and where the insect species are regarded as key pests of corn). Bioassay methods must be appropriate for the goal of detecting field-relevant shifts in population response to lepidopteran resistant Bt corn and/or changes in resistance allele frequency in response to the use of Bt corn and, as far as possible, should be consistent across sampling years to enable comparisons with historical data.

The number of populations to be collected shall reflect the regional importance of the insect species as a pest, and specific collection regions will be identified for each pest. For ECB, a minimum of 12 populations across the sampling region will be targeted for collection at each annual sampling. For SWCB, the target will be a minimum of six populations. For CEW, the target will be a minimum of 10 populations. Pest populations should be collected from multiple corn-growing states reflective of different geographies and agronomic conditions. To obtain sufficient sensitivity to detect resistance alleles before they become common enough to cause measurable field damage, each population collection shall attempt to target 400 insect genomes (egg masses, larvae, mated females, and/or mixed-sex adults), but a successful population collection will contain a minimum of 100 genomes. It is recognized that it may not be possible to collect the target number of insect populations or genomes due to factors such as natural fluctuations in pest density, environmental conditions, and area-wide pest suppression.

The sampling program and geographic range of collections may be modified as appropriate based on changes in pest importance and for the adoption levels of lepidopteran-resistant Bt corn. The Agency shall be consulted prior to the implementation of such modifications.

The registrant will report to the Agency before August 31 each year the results of the population sampling and bioassay monitoring program.

Any incidence of unusually low sensitivity to the Bt protein in bioassays shall be investigated as soon as possible to understand any field relevance of such a finding. Such investigations shall proceed in a stepwise manner until the field relevance can be either confirmed or refuted, and results of these shall be reported to the Agency annually before August 31. The investigative steps will include:

1. Re-test progeny of the collected population to determine whether the unusual bioassay response is reproducible and heritable. If it is not reproducible and heritable, no further action is required.

		CONCURRENCES						
SYMBOL	▶							
SURNAME	▶			2				
DATE	▶							

2. If the unusual response is reproducible and heritable, progeny of insects that survive the diagnostic concentration will be tested using methods that are representative of exposure to Bt corn hybrids under field conditions. If progeny do not survive to adulthood, any suspected resistance is not field relevant and no further action is required.

3. If insects survive steps 1 and 2, resistance is confirmed, and further steps will be taken to evaluate the resistance. These steps may include:

- determining the nature of the resistance (*i.e.*, recessive or dominant, and the level of functional dominance);
- estimating the resistance-allele frequency in the original population;
- determining whether the resistance-allele frequency is increasing by analyzing field collections in subsequent years sampled from the same site where the resistance allele(s) was originally collected;
- determining the geographic distribution of the resistance allele by analyzing field collections in subsequent years from sites surrounding the site where the resistance allele(s) was originally collected.

Should field-relevant resistance be confirmed, and the resistance appears to be increasing or spreading, the registrant will consult with the Agency to develop and implement a case-specific resistance management action plan.

**(2) Investigation of Reports of Unexpected Levels of Damage by the Target Pests:**

The registrant will follow up on grower, extension specialist or consultant reports of unexpected levels of damage by the lepidopteran pests listed on the pesticide label. The registrant will instruct its customers to contact them if such incidents occur. The registrant will investigate all legitimate reports submitted to the company or the company's representatives.

If reports of unexpected levels of damage lead to the suspicion of resistance in any of the key target pests (ECB, SWCB, and CEW), the registrant will implement the actions described below, based on the following definitions of *suspected resistance* and *confirmed resistance*.

CONCURRENCES

SYMBOL								
SURNAME			3					
DATE								

*Suspected resistance*

EPA defines *suspected resistance* to mean field reports of unexpected levels of insect feeding damage for which:

- the corn in question has been confirmed to be lepidopteran-active Bt corn;
- the seed used had the proper percentage of corn expressing Bt protein;
- the relevant plant tissues are expressing the expected level of Bt protein; and
- it has been ruled out that species not susceptible to the protein could be responsible for the damage, that no climatic or cultural reasons could be responsible for the damage, and that there could be no other reasonable causes for the damage.

The Agency does not interpret *suspected resistance* to mean grower reports of possible control failures or suspicious results from annual insect monitoring assays, nor does the Agency intend that extensive field studies and testing be undertaken to confirm scientifically the presence of insects resistant to Bt corn in commercial production fields before responsive measures are undertaken.

If resistance is *suspected*, the registrant will instruct growers to do the following:

- Use alternative control measures in the Bt corn fields in the affected region to control the target pest during the immediate growing season.
- Destroy Bt corn crop residues in the affected region within one month after harvest with a technique appropriate for local production practices to minimize the possibility of resistant insects over-wintering and contributing to the next season's target pest population.

Additionally, if possible, and prior to the application of alternative control measures or destruction of crop residue, the registrant will collect samples of the insect population in the affected fields for laboratory rearing and testing. Such rearing and testing shall be conducted as expeditiously as practical.

*Confirmed resistance*

EPA defines *confirmed resistance* to mean, in the case of field reports of unexpected levels of damage from the key target pests, that all the following criteria are met:

CONCURRENCES

SYMBOL							
SURNAME			4				
DATE							

- There is >30% insect survival and commensurate insect feeding in a bioassay, initiated with neonate larvae, that uses methods that are representative of exposure to Bt corn hybrids under field conditions (ECB and SWCB only).
- In standardized laboratory bioassays using diagnostic concentrations of the Bt protein suited to the target pest in question, the pest exhibits resistance that has a genetic basis and the level of survivorship indicates that there may be a resistance allele frequency of  $\geq 0.1$  in the sampled population.
- In standardized laboratory bioassays, the  $LC_{50}$  exceeds the upper limit of the 95% confidence interval of the  $LC_{50}$  for susceptible populations surveyed both in the original baselines developed for this pest species and in previous years of field monitoring.

**(3) Response to Confirmed Resistance in a Key Target Pest as the Cause of Unexpected Levels of Damage in the Field**

When field resistance is *confirmed* (as defined above), the following steps will be taken by the registrant:

- EPA will receive notification within 30 days of resistance confirmation;
- Affected customers and extension agents will be notified about confirmed resistance within 30 days;
- Monitoring will be increased in the affected area and local target pest populations will be sampled annually to determine the extent and impact of resistance;
- If appropriate (depending on the resistant pest species, the extent of resistance, the timing of resistance, and the nature of resistance, and the availability of suitable alternative control measures), alternative control measures will be employed to reduce or control target pest populations in the affected area. Alternative control measures may include advising customers and extension agents in the affected area to incorporate crop residues into the soil following harvest to minimize the possibility of over-wintering insects, and/or applications of chemical insecticides;
- Unless otherwise agreed with EPA, stop sale and distribution of the relevant lepidopteran-active Bt corn hybrids in the affected area immediately until an effective local mitigation plan approved by EPA has been implemented;
- The registrant will develop a case-specific resistance management action plan within 90 days according to the characteristics of the resistance event and local agronomic needs. The registrant will consult with appropriate stakeholders in the development of the action plan, and the details of such a plan shall be approved by EPA prior to implementation;

CONCURRENCES

SYMBOL								
SURNAME				S				
DATE								

6/12


- Notify affected parties (e.g. growers, consultants, extension agents, seed distributors, university cooperators and state/federal authorities as appropriate) in the region of the resistance situation and approved action plan; and
- In subsequent growing seasons, maintain sales suspension and alternative resistance management strategies in the affected region(s) for the Bt corn hybrids that are affected by the resistant population until an EPA-approved local resistance management plan is in place to mitigate the resistance.

A report on results of resistance monitoring and investigations of damage reports must be submitted to the Agency annually by August 31<sup>st</sup> each year for the duration of the conditional registration.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of Bt11 x MIR604 corn constitutes acceptance of these conditions.

A stamped copy of the FIFRA label is enclosed for your records.

Sincerely,

  
 Sheryl K. Reilly, Ph.D., Chief  
 Microbial Pesticides Branch  
 Biopesticides and Pollution  
 Prevention Division (7511P)

CONCURRENCES

SYMBOL								
SURNAME				6				
DATE								

2  
12

# Bt11 x MIR604 Corn Seed

[Alternate brand name: *Agrisure*<sup>TM</sup> CB/LL/RW]

## Plant-incorporated protectants:

**Cry1Ab and mCry3A proteins for control of corn borers and corn rootworms**

This product is effective in controlling corn leaf, stalk, and ear damage caused by corn borers and root feeding damage caused by corn rootworms.

## Active Ingredients:

- Bacillus thuringiensis* Cry1Ab delta-endotoxin protein and the genetic material necessary for its production (via elements of pZO1502) in corn (SYN-BT011-1).... ≤0.0029%\*
- Modified Cry3A protein and the genetic material necessary for its production (via elements of pZM26) in corn (SYN-IR604-5) ..... ≤0.0069%\*

## Inert Ingredients:

- Phosphinothricin acetyltransferase marker protein and the genetic material necessary for its production (via elements of pZO1502) in corn (SYN-BT011-1)... ≤0.00002%\*
- Phosphomannose isomerase protein and the genetic material necessary for its production (via elements of pZM26) in corn (SYN-IR604-5)..... ≤0.0002%\*

\*Percent (wt/wt) of whole plant on a dry weight basis

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

EPA Registration No. 67979-8  
EPA Establishment No. 66736-NC-01

Syngenta Seeds, Inc. - Field Crops - NAFTA  
P.O. Box 12257  
3054 East Cornwallis Rd.  
Research Triangle Park, NC 27709

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in any manner inconsistent with this labeling. The subject registration will automatically expire at midnight September 30, 2010. All corn seed that contains the plant-incorporated protectant sold or distributed by Syngenta Seeds or its distributors must be accompanied by informational material (e.g., a bag tag) indicating the registration number (67979-8) and the active ingredients, and stipulating that growers read the Grower Guide (or equivalent guidance) prior to planting the seed.

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated**  
**DEC 05, 2008**  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended, for the pesticide  
 registered under EPA Reg. No. 67979-8

**Insects Controlled or Suppressed**

Field corn has been genetically transformed to produce the insecticidal proteins Cry1Ab and mCry3A for control or suppression of the following lepidopteran and coleopteran insects:

- European corn borer (*Ostrinia nubilalis*)
- Southwestern corn borer (*Diatraea grandiosella*)
- Southern cornstalk borer (*Diatraea crambidoides*)
- Corn earworm (*Helicoverpa zea*)
- Fall armyworm (*Spodoptera frugiperda*)
- Sugarcane borer (*Diatraea saccharalis*)
- Common stalk borer (*Papaipema nebris*)
- Western corn rootworm (*Diabrotica virgifera virgifera*)
- Northern corn rootworm (*Diabrotica barberi*)
- Mexican corn rootworm (*Diabrotica virgifera zea*)

**Insect Resistance Management**

The following information regarding commercial production of Bt11 x MIR604 corn must be included in the Grower Guide (or equivalent).


**Corn Belt / Non-Cotton Growing Region Refuge Requirements**

For corn grown in the US Corn Belt two options for deployment of the refuge are available to growers.

The first option is planting a common refuge for both corn borers and corn rootworms. The common refuge must be planted with corn hybrids that do not contain Bt technologies for the control of corn rootworms or corn borers. The refuge area must represent at least 20% of the grower's corn acres (i.e., sum of [Bt11 x MIR604] acres and refuge acres). It can be planted as a block adjacent to the [Bt11 x MIR604] field, perimeter strips, or in-field strips. If perimeter strips are implemented, the strips must be at least 4 consecutive rows wide. If strips within the [Bt11 x MIR604] field are implemented, then at least 4 consecutive rows should be planted. The common refuge can be treated with a soil-applied or seed-applied insecticide to control rootworm larvae and other soil pests. The refuge can also be treated with a non-Bt foliar insecticide for control of late-season pests if pest pressure reaches an economic threshold for damage; however, if rootworm adults are present at the time of foliar applications then the [Bt11 x MIR604] field must be treated in a similar manner.

Economic thresholds will be determined using methods recommended by local or regional professionals (e.g., Extension Service agents, crop consultants, etc.). The following is a schematic of one common refuge-deployment option:

### Common Refuge




The second option is planting separate refuge areas for corn borers and corn rootworms. The corn borer refuge must be planted with a non-Bt/lepidopteran-protected hybrid, must represent at least 20% of the grower's corn acres (i.e., sum of [Bt11 x MIR604] acres and corn borer refuge acres), and must be planted within ½ mile of the [Bt11 x MIR604] field. The corn borer refuge can be treated with a soil-applied or seed-applied insecticide for corn rootworm larval control, or a non-Bt foliar-applied insecticide for corn borer control if pest pressure reaches an economic threshold for damage. The corn rootworm refuge must be planted with a non-Bt/corn rootworm-protected hybrid, but can be planted with Bt corn hybrids that control corn borers. The corn rootworm refuge must represent at least 20% of the grower's corn acres (i.e., sum of [Bt11 x MIR604] acres and corn rootworm refuge acres) and can be planted as an adjacent block, perimeter strips, or in-field strips. The corn rootworm refuge can be treated with a soil-applied or seed-applied insecticide to control rootworm larvae and other soil pests. The refuge can also be treated with a non-Bt foliar insecticide for control of late-season pests; however, if rootworm adults are present at the time of foliar applications then the [Bt11 x MIR604] field must be treated in a similar manner. Growers who fail to comply with the IRM requirements risk losing access to the product.

The following is a schematic of one separate refuge option with the corn rootworm refuge planted as a block within the field and the corn borer refuge planted within a ½ mile of the Bt11 x MIR604 field:

**Separate-Refuge Option**

{Two-Refuge Option, Double-Refuge Option, Paired-Refuge Option}




**Corn/Cotton-Growing Area {Cotton-Growing Area} Refuge Requirements**

For [Bt11 x MIR604] corn grown in cotton-growing areas the common refuge and separate refuge options are also available, however, the refuge area is larger. Cotton-growing areas include the following states: Alabama, Arkansas, Florida, Georgia, Louisiana, North Carolina, Mississippi, South Carolina, Oklahoma (only the counties of Beckham, Caddo, Comanche, Custer, Greer, Harmon, Jackson, Kay, Kiowa, Tillman, and Washita), Tennessee (only the counties of Carroll, Chester, Crockett, Dyer, Fayette, Franklin, Gibson, Hardeman, Hardin, Haywood, Lake, Lauderdale, Lincoln, Madison, Obion, Rutherford, Shelby, and Tipton), Texas (except the counties of Carson, Dallam, Hansford, Hartley, Hutchinson, Lipscomb, Moore, Ochiltree, Roberts, and Sherman) Virginia (only the counties of Dinwiddie, Franklin City, Greensville, Isle of Wight, Northampton, Southampton, Suffolk City, Surrey, and Sussex), and Missouri (only the counties of Dunkin, New Madrid, Pemiscot, Scott, and Stoddard).

The first option is planting a common refuge for both corn borers and corn rootworms. The common refuge must be planted with corn hybrids that do not contain Bt technologies for the control of corn rootworms or corn borers. The refuge area must represent at least 50% of the grower's corn acres (i.e., sum of [Bt11 x MIR604] acres and refuge acres). It can be planted as a block adjacent to the [Bt11 x MIR604] field, perimeter strips, or in-field strips. If perimeter strips are implemented, the strips must be at least 4 consecutive rows wide. If strips within the [Bt11 x MIR604] field are implemented, then at least 4 consecutive rows should be planted. The common refuge can be treated with a soil-applied or seed-applied insecticide to control rootworm larvae and other soil pests. The refuge can also be treated with a non-Bt foliar insecticide for control of late-season pests if pest pressure reaches an economic threshold for damage; however, if rootworm adults are present at the time of foliar applications then the [Bt11 x MIR604] field must be treated in a similar manner.

The following is a schematic of one common refuge-deployment option:


**Common Refuge**




The second option is planting separate refuge areas for corn borers and corn rootworms. The corn borer refuge must be planted with a non-Bt/lepidopteran-protected hybrid, must represent at least 50% of the grower's corn acres (i.e., sum of [Bt11 x MIR604] acres and corn borer refuge acres), and must be planted within 1/2 mile of the [Bt11 x MIR604] field. The corn borer refuge can be treated with a soil-applied or seed-applied insecticide for corn rootworm larval control, or a non-Bt foliar-applied insecticide for corn borer control if pest pressure reaches an economic threshold for damage. The corn rootworm refuge must be planted with a non-Bt corn/rootworm-protected hybrid, but can be planted with Bt corn hybrids that control corn borers. The corn rootworm refuge must represent at least 20% of the grower's corn acres (i.e., sum of [Bt11 x MIR604] acres and corn rootworm refuge acres) and be planted as an adjacent block, perimeter strips, or in-field strips. The corn rootworm refuge can be treated with a soil-applied or seed-applied insecticide to control rootworm larvae and other soil pests. The refuge can also be treated with a non-Bt foliar insecticide for control of late-season pests; however, if rootworm adults are present at the time of foliar applications then the [Bt11 x MIR604] field must be treated in a similar manner. Growers who fail to comply with the IRM requirements risk losing access to the product.

The following are schematics for two separate-refuge options with the corn rootworm refuge planted as a block within the Bt11 x MIR604 field and the corn borer refuge planted as a block within a 1/2 mile of the Bt11 x MIR604 field:

**Separate -Refuge Options**  
{Two-Refuge Options, Double-Refuge Options, Paired Refuge Options}



- or -



Grower agreements will specify that growers must adhere to the refuge requirements that will be described in the Grower Guide {IRM Guide} for Bt11 x MIR604 corn or other applicable product use documents. Growers who fail to comply with the IRM requirements risk losing access to the product.

These refuge requirements do not apply to seed propagation of inbred and hybrid corn seed corn up to a total of 20,000 acres per county and up to a combined U.S. total of 250,000 acres per PIP active ingredient per registrant per year.