

68173-3

07/10/2013

1/17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

July 10, 2013

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Kaken Pharmaceutical Co. Ltd.
C/O Conn & Smith, Inc., Agent
6713 Catskill Road Lorton, VA 22079
Attn: Cynthia Ann. Smith

Subject: Notification of minor label changes per PR Notice 98-10.

1. Add alternate text that excludes a permitted use.
2. Add 0.5 lb. and 33 lb. package sizes.
3. Remove redundancies in the table regarding ornamentals.
4. Add "Not for use in California" statements to two sections of the table.
5. Delete a colon from subsection header.

Product Name: **Endorse Water Dispersible Granules.**
EPA Reg. No: **68173-3.**
Your Submission dated June 14, 2013.

Dear Ms. Smith:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under Pesticide Registration Notice (PRN) 98-10 dated above. A screen of this request has been conducted for its applicability under PRN 98-10, and it has been determined that the action(s) requested falls within the scope of this Notice. Our records have been duly noted, and the label submitted with this application has been stamped "Notification Accepted" and will be placed accordingly in our records.

Three (3) copies of final printed labeling must be submitted to the Agency before your product as modified, may be sold or distributed [PR Notice 82-2 and 40 CFR 156.10(a) (6)].

If you have any questions regarding this action, you may contact Mr. Sylvester George at (703) 603-0688 or via e-mail at george.sylvester@epa.gov.

Sincerely,

Linda A. Hollis

Linda A. Hollis, chief
Biochemical Pesticide Branch
Biopesticides and Pollution
Prevention Division (7511P)



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 68173-3	2. EPA Product Manager Linda Hollis	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) ENDORSE Water Dispersible Granules	PM# 91	
5. Name and Address of Applicant (Include ZIP Code) Kaken Pharmaceutical Co., Ltd. c/o Conn & Smith, Inc., Agent 6713 Catskill Road, Lorton, VA 22079 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Notification Accepted

Date: 7/10/2013

Reviewer: Sylvester George

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of minor label changes:

- (1) add alternate text that excludes a permitted use; (2) add 0.5 lb. and 33 lb. package sizes; (3) remove redundancies; (4) add "Not for use in California" statements; and (5) delete a colon from a subsection header.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 0.5 lb; 1 lb; 2.4 lb; 33 lb		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Cynthia Ann Smith	Title U.S. Agent	Telephone No. (Include Area Code) 703-339-1117 cindy@connsmith.com
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title U.S. Agent Vice President, Conn & Smith, Inc.	
4. Typed Name Cynthia Ann Smith	5. Date June 14, 2013	

June 14, 2013
HAND DELIVERED

Document Processing Desk (NOTIF; E-SUB)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room 5-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Ms. Linda Hollis

Dear Ms. Hollis:

RE: ENDORSE Water Dispersible Granules
EPA Registration No. 68173-3
Notification of Minor Label Change

This letter and the enclosures are submitted on behalf of Kaken Pharmaceutical Co., Ltd.

The most recent label notification was accepted by EPA on September 27, 2012. The purpose of this submission is to update the label by notification to:

1. Add alternate front panel text and General Information text for the AFFIRM WDG alternate brand name that excludes use on field grown food crops;
2. Add two new packages sizes (0.5 pounds and 33 pounds);
3. Remove redundancies in the table regarding ornamentals;
4. Add "Not for use in California" statements to two sections of the table regarding turf grasses; and
5. Delete a colon in the table regarding turf grasses after a sub-section header.

The following are enclosed:

1. Application for amended registration dated June 14, 2013;
2. Label dated June 14, 2013;
 - a. With the changes noted;
 - b. With the changes implemented;
3. Certification of Label Integrity dated June 14, 2013; and
4. Compact Disk (CD) with the June 14, 2013 notification label:
 - a. 068173-00003.20130614.notification_changes_noted.pdf
 - b. 068173-00003.20130614.notification_changes_implemented.pdf

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR §152.46, and no other changes have been made to the labeling of the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR § 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Please feel free to contact me if there are any questions regarding this notification.

Sincerely,



Cynthia Ann Smith, Vice President
Agent for Kaken Pharmaceutical Co., Ltd.

cc: K. Takei, Kaken Pharmaceutical Co., Ltd.

Notification Accepted

068173-00003.20130614.notification_changes_implemented.pdf
ENDORSE Water Dispersible Granules (EPA Reg. No. 68173-3)
June 14, 2013 label NOTIFICATION based upon the September 27, 2012 ACCEPTED notification label
CHANGES IMPLEMENTED
Page 1 of 14

Date: 7/10/2013

Reviewer: *Sylvester George*

Alternate Brand Names for ENDORSE Water Dispersible Granules:

- VERANDA™ Water Dispersible Granules
- AFFIRM™ WDG

**ENDORSE® Water Dispersible Granules
For use on Turf, Ornamentals, and Various Crops**

Alternate text:

For disease control on various crops including:
cucurbit vegetables, fruiting vegetables, pome fruits, strawberries,
and non-bearing fruit vines, and non-bearing fruit and nut trees

Alternate text:

For disease control on Turf, Ornamentals, Christmas Trees,
and various crops including:
cucurbit vegetables, fruiting vegetables, pome fruits, strawberries,
and non-bearing fruit vines, and non-bearing fruit and nut trees

Alternate text:

For use on Turf, Ornamentals, and
Various Food Crops grown only in a Greenhouse, Lath house, or Shade house

GROUP 19 FUNGICIDE

ACTIVE INGREDIENT:

Polyoxin D zinc salt 11.3%

OTHER INGREDIENTS: 88.7%

TOTAL 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE SIDE OR BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS


EPA Reg. No. 68173-3

EPA Est. No. _____

Kaken Pharmaceutical Co., Ltd.
Agrochemicals & Animal Health Products Department
28-8, Honkomagome 2-chome
Bunkyo-ku, Tokyo 113-8650, Japan

For Product Information Call _____

Net Content: 0.5 pounds
1 lb
2 lbs 6.4 oz (2.4 lb)
33 lb (15 kg)



FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBERS	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p> <p style="text-align: center;">FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE</p> <p>Call PROSAR at 1-866-303-6952 or 1-651-632-8946 if calling from outside of the U.S.</p> <p><i>Alternate text:</i> Call CHEMTREC at 1-800-424-9300 or the Poison Control Center at 1-800-222-1222.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through the skin, swallowed or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks
- Shoes
- Chemical-Resistant gloves

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
<ul style="list-style-type: none"> • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

6/17

ENVIRONMENTAL HAZARDS

For terrestrial uses. This pesticide is moderately toxic to aquatic invertebrates and fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not allow runoff into lakes, streams, ponds or public waterways. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Granules exposed on soil surface may be hazardous to fish and aquatic invertebrates. Collect granules spilled during loading.

Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours unless wearing appropriate PPE.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, socks, shoes, chemical resistant gloves.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep children and pets out of treated area until sprays have dried.

GENERAL INFORMATION

ENDORSE Water Dispersible Granules is not for use on turf being grown for commercial seed production.

ENDORSE Water Dispersible Granules is not for homeowner use to treat food crops.

ENDORSE Water Dispersible Granules is for use in controlling or suppressing certain diseases on almonds, cucurbit vegetables, fruiting vegetables, ginseng, grapes, pistachios, pome fruits, potatoes, strawberries, and seedlings or transplants of cucurbit vegetables and fruiting vegetables in a Greenhouse, Lath or Shade house, Ornamentals (Greenhouse, Lath or Shade house, Field, Nursery, Container, and Residential and Commercial Landscapes), and Turf (including golf courses, sod, residential lawns, parks, athletic fields, and commercial and institutional grounds) composed of cool and warm season grasses such as Bluegrass, Bentgrass, Fescue, Ryegrass, Bermudagrass, St. Augustinegrass, Zoysiagrass, or their mixtures.

Alternate text:

ENDORSE Water Dispersible Granules is for use in controlling or suppressing certain diseases on:

- 1) Cucurbit vegetables, fruiting vegetables, ginseng, grapes, pome fruits, strawberries, and seedlings or transplants of cucurbit vegetables and fruiting vegetables in a Greenhouse, Lath house or Shade house,
- 2) Ornamentals (Greenhouse, Lath house or Shade house, Field, Nursery, Container, and Residential and Commercial Landscapes), and
- 3) Turf (including golf courses, sod, residential lawns, parks, athletic fields, and commercial and institutional grounds) composed of cool and warm season grasses such as Bluegrass, Bentgrass, Fescue, Ryegrass, Bermudagrass, St. Augustinegrass, Zoysiagrass, or their mixtures.

Apply ENDORSE Water Dispersible Granules as a preventative or curative treatment in conjunction with good management practices.

RESISTANCE MANAGEMENT RECOMMENDATIONS

ENDORSE Water Dispersible Granules contains a Group 19 fungicide. Fungal isolates with acquired resistance to Group 19 may eventually dominate the fungal population if Group 19 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by ENDORSE Water Dispersible Granules or other Group 19 Fungicides.

To delay fungicide resistance, consider:

- Avoiding the consecutive use of ENDORSE Water Dispersible Granules or other target site of action Group 19 fungicides that have a similar target site of action, on the same pathogens.
- Using tank-mixtures or premixes with fungicides from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Basing fungicide use on a comprehensive IPM program.
- Monitoring treated fungal populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

MIXING AND APPLICATION

When diseases not specified on this label are present or expected, it is appropriate to mix ENDORSE Water Dispersible Granules with appropriately labeled fungicides. When tank mixing ENDORSE Water Dispersible Granules with other products, observe all precautions and limitations on each separate

product label. It is always advisable to conduct a tank mix compatibility test when you plan to mix this product with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. To assess the potential for phytotoxicity, test tank mixtures on a small number of plants prior to more widespread application.

Do not apply this product through any type of irrigation system.

Mixing instructions for ENDORSE Water Dispersible Granules:

- Fill spray tank with water to ½ of the intended final volume.
- Start spray tank agitation.
- Add the appropriate amount of product to the tank according to the rates in the following tables. Follow the directions above for mixing other products in the spray tank.
- Agitate to ensure thorough mixing while adding the remaining required water.
- DO NOT allow the spray mixture to stand without agitation.

Mix only the amount of spray solution needed to treat the desired area.

For optimum control of labeled diseases, apply ENDORSE Water Dispersible Granules with sufficient water to provide a thorough coverage.

ALMONDS		
Disease Control	Rate	Application Notes
Alternaria (<i>Alternaria alternata</i>)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> • Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). • Begin sprays preventatively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/acre) per season. 		

CUCURBIT VEGETABLES †		
[Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra), <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon]		
Disease Control	Rate	Application Notes
Powdery Mildew (<i>Sphaerotheca</i> sp.) Gummy Stem Blight (<i>Didymella bryoniae</i>) Gray Mold (<i>Botrytis</i> sp.) Corynespora leaf spot (<i>Corynespora cassicola</i>) Scab (<i>Cladosporium</i> sp.) Early Blight (<i>Alternaria</i> sp.)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> • Mix in appropriate spray volume appropriate for sufficient coverage. • For all diseases, spray product preventatively and continue as needed on a 10-14 day interval. • 0 day Preharvest Interval.
Notes: † Not for use in California. <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 5 applications of ENDORSE Water Dispersible Granules (3.5 oz ai/acre) per season. 		

FRUITING VEGETABLES (Except Cucurbits) † [Eggplant, Groundcherry, Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato]		
Disease Control	Rate	Application Notes
Powdery Mildew (<i>Leveillula taurica</i> , <i>Oidiopsis sipula</i>)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> Mix in appropriate spray volume appropriate for sufficient coverage. For all diseases, spray product preventatively and continue as needed on a 10-14 day interval. 0 day Preharvest Interval.
Botrytis Rot (<i>Botrytis</i> sp.)		
Early Blight (<i>Alternaria solani</i>)		
Disease Suppression		
Anthracnose (<i>Colletotricum coccodes</i>)		
Notes:		
† Not for use in California.		
• Use in alternation with fungicides that have different modes of action.		
• Do not apply more than 5 applications of ENDORSE Water Dispersible Granules (3.5 oz ai/acre) per season.		

GINSENG †		
Foliar Disease Control	Rate	Application Notes
Alternaria blight (<i>Alternaria panax</i>)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> Apply as a foliar spray every 7–10 days. Apply prior to disease development and when conditions are conducive for disease.¹
Botrytis blight (<i>Botrytis cinerea</i>)		
Root and Crown Disease Control	Rate	Application Notes
Cylindrocarpon root rot (<i>Cylindrocarpon destructans</i>)	6.2 oz/acre (0.7 oz ai/acre)	• Apply as a drench every 14 days.
Rhizoctonia root and crown rot (<i>Rhizoctonia solani</i>)		• Apply as a drench every 14–28 days.
Notes:		
† Not for use in California.		
• Apply within 2 weeks following plant emergence and continue throughout the season.		
¹ Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases.		
• Use in alternation with fungicides that have different modes of action.		
• Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/acre) per season.		

GRAPES		
Disease Control	Rate	Application Notes
Botrytis Bunch Rot (<i>Botrytis cinerea</i>) Powdery Mildew (<i>Uncinula necator</i>)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> Mix in appropriate spray volume appropriate for sufficient coverage. For Botrytis Bunch Rot control, spray product at veraison and 7 days prior to harvest. For Powdery Mildew control, begin sprays preventatively and continue as needed on a 14 day interval. 0 day Preharvest Interval. Do not apply by air.
Disease Suppression	Rate	Application Notes
Powdery Mildew (<i>Uncinula necator</i>)	6.2 oz/acre (0.7oz ai/acre)	<ul style="list-style-type: none"> For Powdery Mildew suppression, begin sprays preventively and continue as needed on a 14-day interval.
Notes: <ul style="list-style-type: none"> Use in alternation with fungicides that have different modes of action. Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/acre) per season. 		

PISTACHIOS		
Disease Control	Rate	Application Notes
Alternaria (<i>Alternaria</i> sp.)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage (avoid every other row applications)). For both diseases, begin sprays preventatively and continue as needed on a 10–14 day interval. 0 day Preharvest Interval. Do not apply by air. Use of an adjuvant may enhance coverage.
Disease Suppression		
Botryosphaeria blight (<i>Botryosphaeria</i> sp.)		
Notes: <ul style="list-style-type: none"> Use in alternation with fungicides that have different modes of action. Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/acre) per season. 		

12/17

POME FRUITS † [Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, Quince]		
Disease Control	Rate	Application Notes
Powdery Mildew <i>(Apple: Podosphaera leucotricha</i> <i>Pear: Phyllactinia mali)</i> Alternaria Leaf Spot <i>(Alternaria mali)</i> Blotch <i>(Diplocarpon mali)</i>	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> Mix in appropriate spray volume appropriate for sufficient coverage. Start sprays for scab control at green tip and continue as needed on a 7-10 day interval. For Powdery Mildew control, begin preventatively and repeat as necessary on a 10-14 day interval. Use of an alternation program with a sterol inhibiting fungicide is recommended. 0 day Preharvest Interval.
Disease Suppression		
Scab <i>(Venturia sp.)</i>		
Notes: † Not for use in California. <ul style="list-style-type: none"> Use in alternation with fungicides that have different modes of action. Do not apply more than 6 applications of ENDORSE Water Dispersible Granules (4.2 oz ai/acre) per season. 		

POTATOES †		
Disease Control	Rate	Application Notes
Early Blight <i>(Alternaria solani)</i>	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none"> Mix in appropriate spray volume appropriate for sufficient coverage. For all diseases, spray product preventatively and continue as needed on a 10-14 day interval. 0 day Preharvest Interval.
Notes: † Not for use in California. <ul style="list-style-type: none"> Use in alternation with fungicides that have different modes of action. Do not apply more than 5 applications of ENDORSE Water Dispersible Granules (3.5 oz ai/acre) per season. 		

13/17

068173-00003.20130614.notification_changes_implemented.pdf
ENDORSE Water Dispersible Granules (EPA Reg. No. 68173-3)
June 14, 2013 label NOTIFICATION based upon the September 27, 2012 ACCEPTED notification label
CHANGES IMPLEMENTED
Page 10 of 14

STRAWBERRIES †		
Disease Control	Rate	Application Notes
Anthracnose Leaf and Fruit Rot (<i>Colletotricum</i> sp.)	6.2 oz/acre (0.7 oz ai/acre)	<ul style="list-style-type: none">Mix in appropriate spray volume appropriate for sufficient coverage.Begin applications preventatively and continue as needed on a 7–10 day interval.0 day Preharvest Interval.Do not apply by air.Use of adjuvant may enhance coverage.
Botrytis Fruit Rot (<i>Botrytis cinerea</i>)		
Powdery Mildew (<i>Sphaerotheca macularis</i>)		
Notes: † Not for use in California. <ul style="list-style-type: none">Use in alternation with fungicides that have different modes of action.Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/acre) per season.		

ORNAMENTALS		
Foliar Disease Control	Rate	Application Notes
Anthracnose (<i>Colletotrichum</i>) Botrytis blight (<i>Botrytis cinerea</i>) Curvularia Downy Mildew (<i>Pernospora sp.</i>) (<i>Plasmopara sp.</i>) Powdery Mildew (<i>Oidium sp.</i>) (<i>Erysiphe sp.</i>) (<i>Sphaerotheca sp.</i>) Rhizoctonia aerial blight (<i>Rhizoctonia solani</i>)	0.25 – 0.5 lb/100 gal/acre (0.03 – 0.06 lb ai/100 gal/acre)	<ul style="list-style-type: none"> • Apply as a full coverage foliar spray every 7 – 10 days with sufficient water for plant type and maturity (50-300 gallons/acre). • Apply prior to disease development and when conditions are conducive to disease development. Consult your local extension agent or local recommendations for recommendations for information relating to proper timing for control of these diseases.**
Alternaria blight (<i>Alternaria panax</i>) Petiole Rot (<i>Myrothecium</i>) Apple scab (<i>Venturia inaequalis</i>)	0.5 lb/100 gal/acre (0.06 lb ai/100 gal/acre)	
Root and Crown Disease Control	Rate	Application Notes
*** Rhizoctonia root and crown rot (<i>Rhizoctonia solani</i>)	0.25 – 0.5 lb/100 gal/acre (0.03 – 0.06 lb ai/ 100 gal/acre)	<ul style="list-style-type: none"> • Apply as a soil drench every 14 – 28 days.
Black Root Rot (<i>Thielaviopsis</i>)	0.5 lb (0.06 lb ai/100 gal/acre)	
Notes: <ul style="list-style-type: none"> • ** Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases. • *** Includes Christmas tree (nursery and plantation) and non-bearing vines fruit and fruit and nut trees. • Use in alternation with fungicides that have different modes of action. 		

15/17

TURF GRASSES		
Disease	Rate	Application Notes
Control	2.4 lbs /acre (0.27 lb ai/acre)	<ul style="list-style-type: none"> Repeat treatment on a 7-14 day schedule where environmental conditions are conducive to development of diseases. Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases. When symptoms are present, best control will be achieved by using a shorter interval. For best results apply after mowing. Apply to obtain thorough plant coverage in a minimum of 66 gallons water per acre.
Brown Patch (<i>Rhizoctonia solani</i>)		
Large Patch (<i>Rhizoctonia solani</i>)		
Aids in Control †		
Cool Season Brown Patch or Yellow Patch (<i>Rhizoctonia cerealis</i>)		
Foliar and Basal Anthracnose (<i>Colletotrichum cereale</i>)		
Gray Snow Mold (<i>Typhula ishikariensis</i> and <i>Typhula incarnate</i>)		
Leaf Spot/Melting Out (<i>Dreschlera poae</i>)		
Pink Snow Mold (<i>Microdochium nivale</i>)		
Red Thread (<i>Laetisaria fuciformis</i>)		
Rhizoctonia Damping Off (<i>Rhizoctonia solani</i>)		
Rhizactonia Leaf and Sheath Blight (<i>Rhizoctonia zeae</i>)		
Waitea Patch (Brown Ring Patch) (<i>Waitea circinata</i>)		
Zoysia Patch (<i>Rhizoctonia solani</i>)		

16/17

TURF GRASSES		
Disease	Rate	Application Notes
Aids in Suppression †		
Gray Leaf Spot (<i>Pyricularia grisea</i>)		
Suppression and Short Term Control	1 oz/1000 sq ft in minimum of 1 gallon of water (0.27 lb ai/acre in 66 gallons of water)	<ul style="list-style-type: none"> • Make two-three applications on a 7-day interval schedule. • Use a penetrating wetting agent. • Immediately following application, water in the treatment with sufficient irrigation (1/8-1/4") to wet the active root zone.
Fairy Ring (including <i>Marasmius</i> spp., <i>Lepiota</i> , spp. and <i>Agarius</i> spp.)		
Notes: † Not for use in California. <ul style="list-style-type: none"> • With the exception for treatment of Fairy Ring, irrigation or rainfall soon after treatment with ENDORSE Water Dispersible Granules will decrease disease control. For best control, do not irrigate for 12 hours after treatment. • Use in alternation with fungicides that have different modes of action. 		

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store in dry place away from food or feed.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed, by State and local authorities, by burning. If burned, stay out of smoke.

**FOR 24-HOUR CHEMICAL EMERGENCY (Spill, Leak, Fire or Accident) ASSISTANCE:
 CALL CHEMTREC 1-800-424-9300**

17/17

Warranty and Disclaimer Statement

1. The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Kaken Pharmaceutical Co. Ltd. ("Kaken"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.
2. Kaken warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.
3. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Kaken, and is subject to the inherent risks described above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, KAKEN DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, KAKEN, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF KAKEN IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.**
4. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF KAKEN, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT KAKEN'S ELECTION, THE REPLACEMENT OF THE PRODUCT.**

ENDORSE® is a registered trademark of Arysta LifeScience North America, LLC.
VERANDA™ is a trademark of Arysta LifeScience North America, LLC.
AFFIRM™ is a trademark of Cleary Chemical, LLC.