


68186-1

03/23/2000

1/12



PURE PRODUCT FROM THE JOJOBA BEAN

For Powdery Mildew Control

ACCEPTED
3/23/00

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 68186-1

ACTIVE INGREDIENT:

Jojoba Oil	97.50%
Other Ingredients	2.50%
Total	100.00%

EPA Reg. No. 6S186-1
Patent Pending

EPA Est. No. 4849S-CA-001

**-KEEP OUT OF REACH OF CHILDREN-
CAUTION**

FIRST AID	
If Product:	
Gets in eyes then:	
<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 5 minutes. ▪ Remove contact lenses, if present. ▪ Continue to rinse eyes for 10-15 minutes. ▪ Call a poison control center or doctor for treatment advice. 	
Gets on skin then:	
<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice. 	
Is swallowed then:	
<ul style="list-style-type: none"> ▪ Have person sip a glass of water if able to swallow. 	

- NOTES:**
- Have the product container or label with you when calling a poison control center or doctor or going for treatment.
 - In the event of a medical emergency, you may also contact the National Pesticide Telecommunications Network at 1-800-858-7378.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents:
2.5 Gallons

Manufactured For:
IJO PRODUCTS, LLC
Fresno, CA 93755

**PRECAUTIONARY STATEMENTS:
HAZARDS TO
HUMANS AND DOMESTIC ANIMALS:**


CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear long-sleeved shirts, pants, waterproof gloves, shoes and socks. Follow manufacturer instructions for cleaning and maintaining personal protective equipment. If no instructions for washables available; use soap and water. Keep personal protective equipment separate from other laundry.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply directly to lakes, streams or ponds. Do not apply directly to water, or areas where surface water is present, or to intertidal areas below mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash water. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Apply this product only as specified on this label.




STORAGE AND DISPOSAL:

**DO NOT CONTAMINATE WATER, FOOD
OR FEED BY STORAGE OR DISPOSAL:**

STORAGE: Store product in original container in a cool, (not in direct sunlight), dry, locked place out of reach of children. Do not store below 50°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facilities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State or local authorities, by burning.



AGRICULTURAL USE REQUIREMENTS:

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It also contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short-sleeved shirt and short pants; waterproof gloves; shoes plus socks; and protective eyewear.

**DIRECTIONS FOR USE:
READ ENTIRE LABEL BEFORE USING
GROUND APPLICATION ONLY**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or by drift. Only protected handlers may be in area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulations.

Grapes – For control of powdery mildew make the first application after powdery mildew infection is observed on new shoots. Make a full coverage spray of the growing shoots using 2 quarts E-RASE in 50-100 gallons water per acre according to the extent of foliar growth. E-RASE is a contact control agent that requires thorough spray coverage to achieve commercial control of powdery mildew. Make a second application when mildew infections are again observed. Increase the spray gallonage, as increased growth occurs, to 100-200 gallons of water per acre and include 2 quarts E-RASE per acre, depending on the extent of growth and the extent of disease pressure. [Foliar coverage on pubescent (hairy leafed) foliage will be more difficult to achieve. To achieve full foliar wetting, it may be necessary to increase the surfactant level, but not to exceed 0.5% (v/v)] To avoid foliar injury, do not continue to make applications if chlorotic and/or brown leaf margins, or other injury symptoms occur.

If flower development is advancing close to bloom do not make a third application until after bloom has passed and when disease is again observed. Use the rate of E-RASE and spray gallonage required for the second spray application. On wine and raisin grapes, additional mildew sprays can continue when damaging mildew infections are observed. If loss of natural finish at harvest is critical as with fruit destined for fresh-market end use, application of E-RASE should not be made closer than 6-8 weeks prior to harvest and the concentration shall not exceed 2 quarts E-RASE per 100 gallons water per acre. If fruit finish at harvest is not critical, E-RASE can be used to control mildew with repeated application using prescribed rates and spray volume through harvest as economic levels of mildew occur. For applications made later than 2-4 weeks preharvest, do not exceed 2 quarts E-RASE in 100-200 gallons water per acre per application or injury may occur in the form of berry desiccation. Do not spray when temperatures drop below 50°F.

If E-RASE is combined in the spray tank with other pesticides, the most restrictive label requirements apply.

Shake well before using and agitate spray solution. Be sure tank mix combinations, if used, are physically compatible before using them on large areas. If they produce crop injury or do not mix well, do not use the products in combination.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.

4/12

DETURTM

PURE PRODUCT FROM THE JOJOBA BEAN
For Control of Whitefly & Powdery Mildew

ACCEPTED
3/23/00
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 68186-1

ACTIVE INGREDIENT

Jojoba Oil	97.50%
Other Ingredients	2.50%
Total	100.00%

EPA Reg. No. 58186-1

EPA Est. No. 68186-CA-002 or
48498-CA-001

JOJOBA OIL is extracted from JOJOBA SEEDS

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID
If Product:
Gets in eyes then:
<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 5 minutes. Remove contact lenses, if present. Continue to rinse eyes for 10-15 minutes. Call a poison control center or doctor for treatment advice.
Gets on skin then:
<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Is swallowed then:
<ul style="list-style-type: none"> Have person sip a glass of water if able to swallow.

NOTES:

- Have the product container or label with you when calling a poison control center or doctor or going for treatment.
- In the event of a medical emergency, you may also contact the National Pesticide Telecommunications Network at 1-800-858-7378.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Net Contents:
2.5 Gallons

LJO PRODUCTS, LLC
Fresno, CA 93755
(3/2000)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS:**

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear long-sleeved shirts, pants, waterproof gloves, shoes and socks. Follow manufacturers instructions for cleaning and maintaining personal protective equipment. If no instructions for washables available; use soap and water. Keep personal protective equipment separate from other laundry.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply directly to lakes, streams or ponds. Do not apply directly to water, or areas where surface water is present, or to intertidal areas below mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash water. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Apply this product only as specified on this label.

**STORAGE AND DISPOSAL
DO NOT CONTAMINATE WATER, FOOD OR
FEED BY STORAGE OR DISPOSAL:**

STORAGE: Store product in original container in a cool, (not in direct sunlight), dry, locked place out of reach of children. Do not store below 50°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facilities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State or local authorities, by burning.

AGRICULTURAL USE REQUIREMENTS:

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It also contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short-sleeved shirt and short pants; waterproof gloves; shoes plus socks; and protective eyewear.

DIRECTIONS FOR USE

READ ENTIRE LABEL BEFORE USING

GROUND APPLICATION ONLY


It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or by drift. Only protected handlers may be in area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulations.

All Crops – For control of white fly use a rate of ¼% to 1% (1 quart to 1 gallon per 100 gallons water) DETUR at any time during the growing season, including the day of harvest. Thorough and uniform crop coverage is required. Repeat applications may be needed, particularly with heavy population pressure. Adults are only controlled as they become entangled in the drying deposit. Eggs are smothered by the waxy coating. Gaps in spray coverage may allow some eggs to survive, requiring subsequent application.

Spray coverage is important. Continuous agitation of the spray mixture is required to assure proper deposit of DETUR. Spray gallonage may vary according to crop size. Use ¼% to 1% v/v DETUR in 10-15 gallons of total spray per acre with small crops, such as early lettuce. Increase spray volume as crop size and foliage density increase. One hundred (100) or more gallons per acre of spray may be required to adequately cover infested trees. Sprays should be made to wet without excessive runoff. Reapply following significant rain storms.

DETUR can be applied with most conventional spray equipment, i.e., dilute airblast, or boom-type sprays may be made. Electrostatic sprays can also be used. Do not exceed 1% DETUR concentration and do not make 1% applications closer together than seven days; stop application if chlorotic and/or brown leaf margins, or other injury symptoms occur. Too frequent applications of the high rate may injure some foliage. Lower rates, i.e. 1 to 2 quarts DETUR per acre in enough water for appropriate spray coverage as described above can be used in a sustained weekly program during periods of excessive growth dilution, i.e. when rapid growth of the crop to be protected is occurring.

If DETUR is combined in the spray tank with other pesticides the most restrictive label requirements apply.




All Plants In and Around the Home – [Foliar coverage on pubescent (hairy leafed) plants will be more difficult to achieve. To achieve full foliar wetting, it may be necessary to increase the surfactant level, but not to exceed 0.5% (v/v).] For control of powdery mildew make the first application after powdery mildew infection is observed. Make a full coverage spray of the growing plants using 1-2 quarts DETUR in 100 gallons water per acre (.75 fl. oz. to 1.5 fl. oz. in 2.5 gallons water per 1000 ft²) according to the extent of foliar growth and extent of disease pressure. Detur is a contact control agent that requires thorough spray coverage to achieve commercial control of powdery mildew. Make follow-on applications when mildew infections are again observed. As increased growth occurs, increase the spray gallonage to 100-200 gallons of water containing 2 quarts DETUR per acre (1.5 fl. oz. in 2.5 to 5.0 gallons of water per 1000 ft²), depending on the extent of growth and the extent of disease pressure. To avoid foliar injury, do not continue to make applications if chlorotic, and/or brown leaf margins, or other injury symptoms occur.

Continuous agitation of the spray mixture is required to assure proper deposit of DETUR. DETUR can be used to control mildew with repeated application using prescribed rates and spray volume through harvest as economic levels of mildew occur.

Grapes - For control of powdery mildew make the first application after powdery mildew infection is observed on new shoots. Make a full coverage spray of the growing shoots using 2 quarts DETUR in 50-100 gallons water per acre according to the extent of foliar growth. Detur is a contact control agent that requires thorough spray coverage to achieve commercial control of powdery mildew. Make a second application when mildew infections are again observed. Increase the spray gallonage, as increased growth occurs, to 100-200 gallons of water per acre and include 2 quarts DETUR per acre, depending on the extent of growth and the extent of disease pressure. [Foliar coverage on pubescent (hairy leafed) foliage will be more difficult to achieve. To achieve full foliar wetting, it may be necessary to increase the surfactant level, but not to exceed 0.5% (v/v).] To avoid foliar injury, do not continue to make applications if chlorotic, and/or brown leaf margins, or other injury symptoms occur.

If flower development is advancing close to bloom do not make a third application until after bloom has passed and when disease is again observed. Use the rate of DETUR and spray gallonage required for the second spray application. On wine and raisin grapes, additional mildew sprays can continue when damaging mildew infections are observed. If loss of natural finish at harvest is critical as with fruit destined for fresh-market end use, application of DETUR should not be made closer than 6-8 weeks prior to harvest and the concentration shall not exceed 2 quarts DETUR per 100 gallons water per acre. If fruit finish at harvest is not critical, DETUR can be used to control mildew with repeated application using prescribed rates and spray volume through harvest as economic levels of mildew occur. For applications made later than 2-4 weeks preharvest, do not exceed 2 quarts DETUR in 100-200 gallons water per acre per application or injury may occur in the form of berry desiccation. Do not spray when temperatures drop below 50°F.

Ornamentals (Outdoor, Greenhouse, Shadehouse, Nurseries) – For control of powdery mildew make the first application after powdery mildew infection is observed. Make a full coverage spray of the growing plants using 1-2 quarts DETUR in 100 gallons water per acre (.75 fl. oz. to 1.5 fl. oz. in 2.5 gallons water per 1000 ft²) according to the extent of foliar growth and extent of disease pressure. Detur is a contact control agent that requires thorough spray coverage to achieve commercial control of powdery mildew. [Foliar coverage on pubescent (hairy leafed) plants will be more difficult to achieve. To achieve full foliar wetting, it may be necessary to increase the surfactant level, but not to exceed 0.5% (v/v).] Make follow-on applications when mildew infections are again observed. As increased growth occurs, increase the spray gallonage to 100-200 gallons of water containing 2 quarts DETUR per acre (1.5 fl. oz. in 2.5 to 5.0 gallons of water per 1000 ft²), depending on the extent of growth and the extent of disease pressure. To avoid foliar injury, do not continue to make applications if chlorotic, and/or brown leaf margins, or other injury symptoms occur.



Spray coverage is important. Continuous agitation of the spray mixture is required to assure proper deposit of DETUR. If time of flowering is critical, as with cutflowers destined for fresh-market, the concentration shall not exceed 2 quarts DETUR per 100 gallons water per acre, (1.5 fl. oz. in 2.5 gallons water per 1000 ft²) per application. If flowering delay is not critical, DETUR can be used to control mildew with repeated application using prescribed rates and spray volume through harvest as economic levels of mildew occur.


8/12

NOTE: Greenhouse and Outdoor Ornamentals: Environmental factors significantly affect phytotoxicity. While DETUR has been tested on numerous ornamental plants without causing serious phytotoxicity, because of the numerous species and varieties grown, it is recommended that a small group of plants be treated at the recommended rate under the anticipated growing conditions and observed for phytotoxic symptoms before treating a large number of plants.

If DETUR is combined in the spray tank with other pesticides, the most restrictive label requirements apply.

Shake well before using and agitate spray solution. Be sure tank mix combinations, if used, are physically compatible before using them on large areas. If they produce crop injury or do not mix well, do not use the products in combination.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.





ERFANSE™

PURE PRODUCT FROM THE JOJOBA BEAN
For Powdery Mildew Control

ACCEPTED
3/23/00
Under the Federal Insecticides, Fungicides, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 68186-1

ACTIVE INGREDIENT:

Jojoba Oil	97.50%
Other Ingredients	<u>2.50%</u>
Total	100.00%

EPA Reg. No. 68186-1
Patent Pending

EPA Est. No. 48498-CA-001

**-KEEP OUT OF REACH OF CHILDREN-
CAUTION**

FIRST AID	
If Product:	
Gets in eyes then:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 5 minutes. ▪ Remove contact lenses, if present. ▪ Continue to rinse eyes for 10-15 minutes. ▪ Call a poison control center or doctor for treatment advice.
Gets on skin then:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
Is swallowed then:	<ul style="list-style-type: none"> ▪ Have person sip a glass of water if able to swallow.

- NOTES:**
- Have the product container or label with you when calling a poison control center or doctor or going for treatment.
 - In the event of a medical emergency, you may also contact the National Pesticide Telecommunications Network at 1-800-858-7378.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents:
2.5 Gallons

Manufactured For:
IJO PRODUCTS, LLC
Fresno, CA 93755

**PRECAUTIONARY STATEMENTS:
HAZARDS TO
HUMANS AND DOMESTIC ANIMALS:**

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear long-sleeved shirts, pants, waterproof gloves, shoes and socks. Follow manufacturer instructions for cleaning and maintaining personal protective equipment. If no instructions for washables available; use soap and water. Keep personal protective equipment separate from other laundry.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply directly to lakes, streams or ponds. Do not apply directly to water, or areas where surface water is present, or to intertidal areas below mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash water. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Apply this product only as specified on this label.


STORAGE AND DISPOSAL:

**DO NOT CONTAMINATE WATER, FOOD
OR FEED BY STORAGE OR DISPOSAL:**

STORAGE: Store product in original container in a cool, (not in direct sunlight), dry, locked place out of reach of children. Do not store below 50°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facilities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State or local authorities, by burning.



AGRICULTURAL USE REQUIREMENTS:

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It also contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short-sleeved shirt and short pants; waterproof gloves; shoes plus socks; and protective eyewear.

**DIRECTIONS FOR USE
READ ENTIRE LABEL BEFORE USING
GROUND APPLICATION ONLY**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or by drift. Only protected handlers may be in area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulations.

Ornamentals (Outdoor, Greenhouse, Shadehouse, Nurseries) – For control of powdery mildew make the first application after powdery mildew infection is observed. Make a full coverage spray of the growing plants using 1-2 quarts E-RASE in 100 gallons water per acre (.75 fl. oz. to 1.5 fl. oz. in 2.5 gallons water per 1000 ft²) according to the extent of foliar growth and extent of disease pressure. E-RASE is a contact control agent that requires thorough spray coverage to achieve commercial control of powdery mildew. [Foliar coverage on pubescent (hairy leafed) plants will be more difficult to achieve. To achieve full foliar wetting, it may be necessary to increase the surfactant level, but not to exceed 0.5% (v/v).] Make follow-on applications when mildew infections are again observed. As increased growth occurs, increase the spray gallonage to 100-200 gallons of water containing 2 quarts E-RASE per acre (1.5 fl. oz. in 2.5 to 5.0 gallons of water per 1000 ft²), depending on the extent of growth and the extent of disease pressure. To avoid foliar injury, do not continue to make applications if chlorotic, and/or brown leaf margins, or other injury symptoms occur.

Spray coverage is important. Continuous agitation of the spray mixture is required to assure proper deposit of E-RASE. If time of flowering is critical, as with cutflowers destined for fresh-market, the concentration shall not exceed 2 quarts E-RASE per 100 gallons water per acre, (1.5 fl. oz. in 2.5 gallons water per 1000 ft²) per application. If flowering delay is critical, E-RASE can be used to control mildew with repeated application using prescribed rates and spray volume through harvest as economic levels of mildew occur.

NOTE: Greenhouse and Outdoor Ornamentals: Environmental factors significantly affect phytotoxicity. While E-RASE has been tested on numerous ornamental plants without causing serious phytotoxicity, because of the numerous species and varieties grown, it is recommended that a small group of plants be treated at the recommended rate under the anticipated growing conditions and observed for phytotoxic symptoms before treating a large number of plants.

If E-RASE is combined in the spray tank with other pesticides, the most restrictive label requirements apply.

Shake well before using and agitate spray solution. Be sure tank mix combinations, if used, are physically compatible before using them on large areas. If they produce crop injury or do not mix well, do not use the products in combination.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.

NEW E-RASE™

PURE PRODUCT FROM THE JOJOBA BEAN
For Powdery Mildew Control

ACCEPTED
3/23/00
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 68186-1

ACTIVE INGREDIENT:

Jojoba Oil	97.50%
Other Ingredients	2.50%
Total	100.00%

EPA Reg. No. 68186-1
Patent Pending

EPA Est. No. 48498-CA-001

**-KEEP OUT OF REACH OF CHILDREN-
CAUTION**

FIRST AID	
If Product:	
Gets in eyes then:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 5 minutes. Remove contact lenses, if present. Continue to rinse eyes for 10-15 minutes. Call a poison control center or doctor for treatment advice.
Gets on skin then:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Is swallowed then:	<ul style="list-style-type: none"> Have person sip a glass of water if able to swallow.

NOTES:

- Have the product container or label with you when calling a poison control center or doctor or going for treatment.
- In the event of a medical emergency, you may also contact the National Pesticide Telecommunications Network at 1-800-858-7378.

**PRECAUTIONARY STATEMENTS:
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS:**

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

**READ ENTIRE SUPPLEMENTAL LABELING
AND USE ONLY AS DIRECTED**

AGRICULTURAL USE REQUIREMENTS:

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements: in the Directions for Use section for information about this standard.

BATCH NO.

Net Contents: 2.5 Gallons

Manufactured For:
IJO PRODUCTS, LLC
Fresno, CA 93755

for outside container

2/11

PAGE 5

FAX: 5592750342

SIEMER&ASSOC

MAR-14-00 TUE 12:32 PM

P. 05

95%

5592750842

MAR-14-2000 15:46