# 10/17/97

# **DELTAMETHRIN 4% COLLAR**

**FLEA & TICK COLLAR FOR DOGS** 

70

61 NN

P4:03

M1/3

ACTIVE INGREDIENT:	Percentage by Weight
Deltamethrin [(s)-alpha-cyano-3-phenoxybenzyl-(1R,3R)-3-	4.0%
(2,2-dibromovinyl)-2,2-dimethyl-cyclopropanecarboxylate]	
INERT INGREDIENTS	<u>96.0%</u>
Total	100.0%

## CAUTION: Do Not Let Children Play With This Collar

### See Side Panel for additional Precautionary Statements

1

EPA Reg. No. 68451-1 EPA Est. No. 68451-FRA-1

NET CONTENTS: 1 Collar NET WT. 1.1 oz

ACCEPTEI	2
OCT 17 1997	
Under the Federal Insecticide. Fungicide, and Rodenticilie Au as amonded, for the posticide registered sader EPA ling. No. 68451-1	ci.

page 1 of 3 13 June 1997

Hoechst Roussel Vet SA 102 route de Noisy 93235 Romainville, France

### **READ ENTIRE LABEL BEFORE EACH USE**

### **USE ONLY ON DOGS**

## **DELTAMETHRIN 4% COLLAR**

### FLEA & TICK COLLAR FOR DOGS

Deltamethrin 4% Collar provides full season, up to 6 months protection against fleas and ticks.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION:

Do not open protective pouch until ready to use. Do not let children play with this collar.

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Do not use on puppies under 12 weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated, or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterianarian immediately.

### FIRST AID

**IF SWALLOWED:** Call a physician or poison control center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention. IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

#### **EMERGENCY PHONE NUMBERS:**

FOR HUMAN, FIRE, ENVIRONMENTAL: 1-800-228-5635, ext. 132 FOR ANIMALS: 1-800-345-4735, EXT. 104

Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Applying other pesticides on the dog may not be necessary while the collar is being\_worn.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Deltamethrin 4% Collar, containing deltamethrin insecticide, has been specially formulated using patented insecticiderelease technology. Maximum effectiveness may not occur for 2 –3 weeks after collar placement. Fleas (*Ctenocephalides sp.*) on the dog will be killed and ones which are present in the dog's environment that may appear on your pet will be killed. Collar will kill ticks including Brown dog tick (*Rhipicephalus sanguineus*) and American dog tick (*Dermacentor variabilis*). This collar should be worn continuously. Reapply a new collar every 6 months.

Place the collar around dog's neck, buckle and adjust for proper fit. Cut off approximately 2 inches from the buckle and dispose of excess length by wrapping in newspaper and placing in trash. The collar must be worn loosely so that two fingers may be placed between collar and dog's neck. Living and rest areas of pet must also be treated with appropriate pest control measures to ensure control of pests. Wetting will not impair the collar's effectiveness or the pet's protection. If the dog goes swimming or is out in the rain, it is not necessary to remove the collar . Deltamethrin 4% Collar may be used in addition to a lead or constraint collar . Use only one Deltamethrin collar at a time.

> page 2 of 3 13 June 1997

STORAGE AND DISPOSAL: Store in original, unopened container, away from children. Do not reuse container or used collar. Wrap in newspaper and put in trash.

#### **IMPORTANT NOTICE: DISCLAIMER**

Read "IMPORTANT NOTICE: DISCLAIMER" before buying or using. If terms are not acceptable, return at once unopened. HOECHST ROUSSEL VET S.A. warrants only that the product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to HOECHST ROUSSEL VET S.A., and user assumes the risk of any such use. HOECHST ROUSSEL VET S. A MAKES NO OTHER WARRANTY, EXPRESSED OR IMPLIED, INCLUDING ANY IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR OF MERCHANTABILITY. In no case shall HOECHST ROUSSEL VET S.A. be liable for consequential, special, indirect or incidental damages resulting from the use or handling of this product. The foregoing conditions of sale and warranty can be varied only by an agreement in writing signed by a duly authorized representative of HOECHST ROUSSEL VET S.A.

The Hoechst Name and logo are registered trademarks of Hoechst AG.

© 1997 HRV S.A.

Part No. XXXXX

Universal Product Code Number (Bar Code)

page-3 of 3 13 June 1997 97 06 13a