

68451-1

08/20/2003

1/2

DELTAMETHRIN 4% COLLAR

FLEA & TICK COLLAR FOR DOGS

Kills Fleas and Ticks for up to 6 Months
LONG LASTING PROTECTION FOR UP TO 6 MONTHS
Kills Ticks (including deer ticks which may carry Lyme disease)
ALSO KILLS FLEAS

ACTIVE INGREDIENT:	Percentage by Weight
Deltamethrin	4.0%
INERT INGREDIENTS	<u>96.0%</u>
Total	100.0%

CAUTION: Do Not Let Children Play With This Collar


See Insert or Side Panel for additional Precautionary Statements
See Back for additional Precautionary Statements

Adjustable – One Size Fits All

EPA Reg. No. 68451-1 EPA Est. No. 68451-FRA-1 NET CONTENTS: 1 Collar
NET WT. 0.9 oz

Distributed by:
Intervet Inc.
405 State Street
Millsboro, DE 19966

Product of France



READ ENTIRE LABEL BEFORE EACH USE

USE ONLY ON DOGS

DELTAMETHRIN 4% COLLAR

FLEA & TICK COLLAR FOR DOGS

Deltamethrin 4% Collar provides full season, up to 6 months protection against fleas and ticks.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Do not open protective pouch until ready to use. Do not let children play with this collar.

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Do not use on puppies under 12 weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated, or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING::	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p style="text-align: center;">For incidents involving HUMANS, call 1-800-680-9206 For incidents involving ANIMALS, call 1-800-548-2423 Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Applying other pesticides on the dog may not be necessary while the collar is being worn.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Deltamethrin 4% Collar, containing deltamethrin insecticide, has been specially formulated using patented insecticide-release technology. Maximum effectiveness may not occur for 2-3 weeks after collar placement. Fleas (*Ctenocephalides sp.*) on the dog will be killed and ones which are present in the dog's environment that may appear on your pet will be killed. Collar will also kill ticks for up to 6 months. Collar will kill ticks including Brown dog tick (*Rhipicephalus sanguineus*), American dog tick (*Dermacentor variabilis*) and deer ticks (*Ixodes scapularis* and *Ixodes pacificus*) which may carry the Lyme disease. This collar should be worn continuously. Reapply a new collar every 6 months.

Place the collar around dog's neck, buckle and adjust for proper fit. Cut off approximately 2 inches from the buckle and dispose of excess length by wrapping in newspaper and placing in trash. The collar must be worn loosely so that two fingers may be placed between collar and dog's neck. Living and rest areas of pet must also be treated with appropriate pest control measures to ensure control of pests. Wetting will not impair the collar's effectiveness or the pet's protection. If the dog goes swimming or is out in the rain, it is not necessary to remove the collar. Deltamethrin 4% Collar may be used in addition to a lead or constraint collar. Use only one Deltamethrin 4% Collar at a time.

STORAGE AND DISPOSAL: Store in original, unopened container, away from children. Do not reuse container or used collar. Dispose of in trash.

IMPORTANT NOTICE: DISCLAIMER

Read "IMPORTANT NOTICE: DISCLAIMER" before buying or using. If terms are not acceptable, return at once unopened. INTERVET INC. warrants only that the product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to INTERVET INC., and user assumes the risk of any such use. INTERVET INC. MAKES NO OTHER WARRANTY, EXPRESSED OR IMPLIED, INCLUDING ANY IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR OF MERCHANTABILITY. In no case shall INTERVET INC. be liable for consequential, special, indirect or incidental damages resulting from the use or handling of this product. The foregoing conditions of sale and warranty can be varied only by an agreement in writing signed by a duly authorized representative of INTERVET INC.