

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

MAR 16 2004

JH Biotech, Inc.  
4951 Olivas Park Drive  
Ventura, CA 93006  
Attn: Tom Dullam  
Head, Plant Nutrition Department

Subject: Label Amendment Application Dated 1/23/04  
Product Name: Fosphite Fungicide  
EPA Registration No:68573-2

Dear Mr. Dullam:

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

Sheryl K. Reilly, Ph.D., Chief  
Biochemical Pesticides Branch  
Biopesticides and Pollution  
Prevention Division (7511C)

Enclosure

CONCURRENCES								
SYMBOL	7511C							
SURNAME	Reilly	Dullam						
DATE	3/15/04	3/16/04						

**Master Label**

**FOSPHITE FUNGICIDE**

**A FUNGICIDE FOR THE CONTROL OF DOWNY MILDEW, PHYTOPHTHORA, PYTHIUM AND VARIOUS OTHER DISEASES ON AGRICULTURAL AND GREENHOUSE CROPS, INDOOR AND OUTDOOR ORNAMENTALS, BEDDING PLANTS AND FORESTREY APPLICATIONS AND DOMESTIC TREES INCLUDING CONIFERS, PINES, OAKS AND TURF**

**ACTIVE INGREDIENT:**  
 Mono- and dipotassium salts of phosphorous acid.....53%

**OTHER INGREDIENTS:**.....47%

**TOTAL**.....100%


**A: SUBLABEL - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS SUBJECT TO WORKER PROTECTION STANDARD**

**B: SUBLABEL - HOUSEHOLD PRODUCTS: FOR NON-COMMERCIAL APPLICATORS (NOT SUBJECT TO WORKER PROTECTION STANDARD)**

**C: SUBLABEL - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS SUBJECT TO WORKER PROTECTION STANDARD (TRUNK INJECTION ONLY)**

**MANUFACTURED BY:**  
 JH Biotech, Inc.  
 P.O. Box 3538  
 Ventura, CA 93006

EPA Reg. No. 68573-2



**SUBLABEL - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS  
SUBJECT TO WORKER PROTECTION STANDARD**

**FOSPHITE FUNGICIDE**

A FUNGICIDE FOR THE CONTROL OF DOWNY MILDEW, PHYTOPHTHORA, PYTHIUM AND VARIOUS OTHER DISEASES ON AGRICULTURAL AND GREENHOUSE CROPS, INDOOR AND OUTDOOR ORNAMENTALS, BEDDING PLANTS AND FORESTREY APPLICATIONS AND DOMESTIC TREES INCLUDING CONIFERS, PINES, OAKS AND TURF

This package can treat up to \_\_\_\_\_ sq. ft. (acres) at the maximum treatment rate.

**ACTIVE INGREDIENTS:**

Mono- and dipotassium salts of Phosphorous Acid\* .....53%

OTHER INGREDIENTS:.....47%

TOTAL.....100%

\*Contains 6.22 lbs./gallon of active ingredients, mono- and dipotassium salts of Phosphorous Acid. Equivalent to 3.90 lbs. Phosphorous Acid/gallon.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side (or back) panel for First Aid and additional precautionary statements.

**NET CONTENTS:**

- 1 pint (0.473 liters)
- 1 quart ( 0.946 liters)
- 1 gallon (3.785 liters)
- 2 1/2 gallons (9.463 liters)
- 5 gallons (18.925 liters)
- Bulk Containers (100 – 4000 gallons)

**MANUFACTURED BY:**

JH Biotech, Inc.  
P.O. Box 3538  
Ventura, CA 93006

EPA Reg. No. 68573-2  
EPA Est. No. 68573-CA001

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after using.

<b>FIRST AID STATEMENTS</b>	
If on skin or clothing	<ul style="list-style-type: none"> <li>·Take off contaminated clothing.</li> <li>·Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>·Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>·Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>·Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>·Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency medical treatment information.</p>	

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves, shoes plus socks and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS BOX**

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Read entire label before using this product.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS:**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, socks, shoes and chemical resistant gloves.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

**GENERAL APPLICATION INSTRUCTIONS**

FOSPHITE is applied by various application methods, including foliar sprays, trunk injection, soil drench, soil incorporation, and bare root dip. For foliar sprays, apply FOSPHITE with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

**MIXING INSTRUCTIONS**

1. Fill the spray tank with 1/2 of the required volume of water.
2. Add FOSPHITE slowly to the tank and agitate by hydraulic or mechanical means.
3. Continue to fill the tank with water to the desired volume while agitating.
4. Continue agitation when applying.

**COMPATIBILITY**

FOSPHITE is compatible with most products used in agriculture; however, crop sensitivity to these mixtures may vary. In particular, mixtures of FOSPHITE with foliar fertilizers and copper products may not be compatible or may cause phytotoxicity to certain plants. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. If these combinations or others have not been previously used, do not tank mix without first testing the mixture's compatibility nor apply it without assessing its safety to the crop (phytotoxicity).

Test the compatibility of spray adjuvants (i.e. stickers, spreaders, wetting agents) with FOSPHITE before use. Do not use acidifying type compatibility agents.

To determine the compatibility of FOSPHITE with other products, use a jar compatibility test. Add the correct proportions of each product and the appropriate quantity of water to clean container, thoroughly mix, then let stand for 3-5 minutes. If the mixture remains in solution or can be remixed readily, the products are considered compatible. To determine if a combination is phytotoxic to a specific crop, spray a few plants/trees, and then evaluate 3-7 days later for visual symptoms.

**APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS**

Apply this product through microjet, drip, solid set and center pivot sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

**PREPARATION OF INJECTION EQUIPMENT:** Remove scale, pesticide residues and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

**APPLICATION INSTRUCTIONS:** Fill tank with 1/2 to 3/4 of desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of Fosphite, then the remaining volume of water. Set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the solution of FOSPHITE into the irrigation water line so as to deliver the desired rate per acre. Inject the FOSPHITE solution with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. For other questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

**NOTE:** For Microjet and Drip Irrigation Systems: Avoid further irrigation of the treated area for 24 to 48 hours after treatment with FOSPHITE has been completed. For Solid Set and Center Pivot Irrigation Systems: Avoid further irrigation of the FOSPHITE treated area until after the foliage has completely dried. This will help to prevent washing the chemical off the crop.

**GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS**

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Maintain continuous agitation in mix tank during mixing and application to assure uniformity solution.** Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution, or when lines containing the product must be dismantled or drained. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of chemigation water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut down and adjust the system as needed.

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

GENERAL INSTRUCTIONS FOR APPLICATIONS

FOOD CROPS

FOSPHITE fungicide is used for control of Downy Mildew, Erwinia amylovora, Pythium, Fusarium, Rhizoctonia and Phytophthora disease (such as purple blotch, late blight, blue mold, soft rot, root rot, brown rot, foot rot, crown/collar rot, canker, fruit/foliar rot, fire blight, heart rot, pink rot, pythium leak, red stele, leather rot, etc.) of agronomic crops. FOSPHITE is applied to plants including:

Berry Crop Grouping: Blackberry ( including bingberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Grapes (all types), Himalayaberry, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these), Blueberry, Cane Berries, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Raspberry (black and red) and Strawberry.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./ gal. (5 L/1000 L of water )	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

02/23/04

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Brassica Crop Grouping:** Broccoli (all types), Brussels sprouts, Cabbage (all types), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach and Rape greens.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	1 qt./100 gal. of water, 1/3 fl. oz./ gal. (2.5 L/1000 L of water)	Apply as pre-plant dip immediately prior to planting. ( 100 gallons is intended to treat the number of slips required to plant one acre.)
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.



**Cereal Grains, Forage and Fiber Crop Grouping:** Alfalfa, Barley, Buckwheat, Canola, Clover (all types), Corn (all types), Cotton, Hay, Kudzu, Lespedeza, Lupin, Millet, Oats, Popcorn, Rice, Rye, Safflower, Sainfoin, Sorghum, Sunflower, Teosinte, Trefoil, Triticale, Vetch, Wheat and Wild rice.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Citrus Crop Grouping (Bearing and Non-Bearing) including Avocado:** Avocado, Calamondin, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange (all types), Pummelo and Satsuma madarin.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./ gal. (5 L/1000 L of water)	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Cucurbit Crop Grouping:** Chayote, Chinese waxgourd, Citron melon, Cucumber (all types), Gherkin, Gourd, Muskmelon, Pumpkin, Squash (summer and winter) and Watermelon.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Herbs and Spices Crop Grouping:** Allspice, Angelica, Anise, Annatto, Balm, Basil, Borage, Bumet, Camomile, Caper buds, Caraway, Cardamom, Cassia bark and buds, Catnip, Celery seed, Chervil, Chive, Cinnamon, Clary, Clove, Cocoa, Coriander, Costmary, Culantro, Cumin, Curry, Dill, Fennel, Fenugreek, Grains of paradise, Hops, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage, Mace, Marigold, Marjoram, Mint, Mustard, Nasturtium, Nutmeg, Parsley, Pennyroyal, Pepper, Poppy seed, Rosemary, Rue, Saffron, Sage, Savory, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff and Wormwood.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Legume Vegetables Crop Grouping:** Bean (all types), broad bean (fava bean), Chick pea (garbanzo bean), Guar, Jackbean, Lablab bean, Lentil, Pea (all types), Peanut, Pigeon pea, Soybean and Sword bean.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year. season.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Pome Fruits Crop Grouping (Bearing and Non-Bearing) including Papaya: Apple, Crabapple, Loquat, Mayhaw, Papaya, Pear, and Quince.**

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> )	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./ gal. (5 L/1000 L of water )	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Stone Fruits Crop Grouping including Olives: Apricot, Cherry, Nectarine, Olive, Peach, Plum, Plumcot and Prune.**

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./ gal. (5 L/1000 L of water )	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more 6 times per crop cycle or year.
	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Tree Nuts Crop Grouping including Vines:** Almond, Beech nut Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Grapes (all types), Hickory nut, Kiwi, Macadamia nut, Pecan, Pistachio and Walnut.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	2 qts./100 gal. of water, 2/3 fl. oz./ gal. (5 L/1000 L of water )	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.



**Vegetables Crop Grouping Including Miscellaneous Food Crops:** (Leafy, Root and Tuber) Amaranth (Chinese spinach, tampala, etc.) Arracacha, Arugula, Arrowroot, Artichoke (all types), Asparagus, Beet (all types), Burdock, Canna, Cardoon, Carrot, Cassava, Celeriac, Celery, Chayote, Chervil (turnip-rooted), Chicory, Chilies, Chrysanthemum (edible), Chufa, Cinchoca, Coconut, Coffee, Corn (all types), Cress (all types), Dandelion, Dasheen, Dock, Eggplant, Endive, Fennel, Garlic, Ginger, Ginseng, Horseradish, Leren, Lettuce (all types), Okra, Onion, Orach, Parsley, Parsnip, Peppers (all types), Pineapple, Purslane, Potato (all types), Radicchio, Radish (all types), Rhubarb, Rutabaga, Salisify (all types), Skirret, Spinach (all types), Sugarcane, Sweet Potato, Swiss chard, Tanier, Tobacco, Tomato, Turmeric, Turnip, Yam (all types).

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.
	Root Dip	1 qt./100 gal. of water, 1/3 fl. oz./ gal. (2.5 L/1000 L of water)	Apply as pre-plant dip immediately prior to planting.( 100 gallons is intended to treat the number of slips required to plant one acre.)
	Irrigation	Overhead: 2-4 qts. in at least 100 gal. of water. (7-9 L in at least 1000 L of water) Drip: 2-3 qts. in at least 100 gal. of water. (5-7 L in at least 1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
Downy Mildew	Foliar	1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

ORNAMENTALS

FOSPHITE fungicide is used for control of Pythium, Downy Mildew, Phytophthora and various other diseases of ORNAMENTALS grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. FOSPHITE may be applied to all ornamental and landscape plants including (but not limited to): Aglaonema, Aphelandra, Arborvitae, Ash, Azaleas, Bougainvilles, Boxwood, Bushes, Cattleya skinneri, Cedars, Ceanothus, Cotoneaster, Cissus, Diffenbachia, Dogwood, Ferns, Ficus, Flowering Trees, Hibiscus, Hard Wood Trees, Holly, Home Lawns, Juniper, Magnolia Trees, Maple, Pines, Oaks, Palms, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Red woods, Rhododendron, Roses, Schefflera, Sedum, Sempervivum, Shrubs, Spathiphyllum, Sycamore, Taxus media, Zygocactus.

Apply FOSPHITE prior to disease development in conjunction with good cultural practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or under heat or moisture stress. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Fusarium Pythium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> ) Bacterial Blight ( <i>Xanthomonas campestris</i> )	Foliar Spray	1 to 2 qts./100 gal. of water or 1/3 fl. oz. to 2/3 fl. oz. per gal. of water. (2.5 to 5 L/1000 L of water)	<p><b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year.</p> <p><b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.</p> <p>Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year.</p>
	Soil Drench	3 qts./100 gal. of water or 1 fl. oz. per gal. of water (7 L/1000 L of water)	Apply 1 gal. of diluted solution per sq. yard and irrigate. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year.
	Soil Incorporation	1 to 2 pts./cubic yard of soil mix (300-600 ml/ cubic meter)	Mix 1 to 2 pints of Fosphite with enough water to wet one cubic yard of soil; mix immediately before potting. If conditions are favorable for disease development, it is recommended to make additional applications as a foliar spray or soil drench.
	Irrigation	Overhead: 1-2 qts./ 100 gal. of water. (2.5 - 5 L/1000 L of water) Drip: 1-2 qts./100 gal. of water. (2.5 - 5 L/1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle.

	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
	Root Dip	2 qt./100 gal. of water or 2/3 fl. oz./gal. of water. (5 L/1000 L of water)	Dip transplants into diluted solution of Fosphite. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
Downy Mildew	Foliar Spray	1 to 2 qts./100 gal. of water or 1/3 fl. oz. to 2/3 fl. oz. per gal. of water (2.5 to 5 L/1000 L of water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

\* Check with your local Extension Agent if you are unsure these conditions exist.

### BEDDING PLANTS

FOSPHITE fungicide is used for control of Downy Mildew, Pythium, Phytophthora and various other diseases of bedding plants grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. FOSPHITE may be applied to all bedding plants including: Ageratum, Algerian Ivy, Alyssum, Anthurium, Artemisia, Aster, Baby's Breath, Begonia, Caladium, Carnation, Chrysanthemum, Coleus, Columbine, Daisy, Delphinium, Easter Lily, English Ivy, Foxglove, Gaillardia, Geranium, Gloxinia, Impatiens, Marigold, Pansy, Petunia, Phlox, Pinks, Poinsettia, Primrose, Prostrate Rosemary, Salvia, Snapdragon, Verbena, Vinca, Zinnia, etc.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> )	Foliar Spray	1 to 2 qts./100 gal. water or 1/3 fl. oz. to 2/3 fl. oz. per gal. (2.5 to 5 L/1000 L water)	<p><b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year.</p> <p><b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply more than 6 times per crop cycle or year.</p> <p>Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year.</p>
	Root Dip	2 qt./100 gal. of water or 2/3 fl. oz./gal. of water. (5 L/1000 L of water)	Dip transplants into diluted solution of Fosphite. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
	Soil Drench	1 to 2 qts./100 gal. water (1/3 fl. oz. to 2/3 fl. oz. per gal.) (2.5 to 5 L/1000 L water)	Apply 1 gal. of solution per sq. yard and irrigate. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year.
	Irrigation	Overhead: 1-2 qts./ 100 gal. of water. (2.5 - 5 L/1000 L of water) Drip: 1-2 qts./100 gal. of water. (2.5 - 5 L/1000 L of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.

Downy Mildew	Foliar Spray	1 to 2 qts./100 gal. water (1/3 fl. oz. to 2/3 fl. oz. per gal.) (2.5 to 5 L/1000 L water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply more than 6 times per crop cycle or year.
--------------	--------------	-----------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

\*Check with your local Extension Agent if you are unsure these conditions exist.

**CONIFERS AND TREES IN COMMERCIAL NURSERIES, PLANTATIONS AND FORESTS,  
LANDSCAPES AND PARKS  
(INCLUDING CHRISTMAS TREES, OAKS and PINES)**

FOSPHITE applied in conjunction with good cultural management practices will control Phytophthora root rot of conifers including (but not limited to): Douglas Fir, spruce and pines also other Phytophthora disease of Oaks (Sudden Oak Death) and forest trees. Use the higher rates when disease pressure is severe. Do not apply to trees that are under heat or moisture stress.

Disease	Application Method	Rate	Application Program
Phytophthora	Foliar Spray	1 to 2 qts./100 gal. water /acre (2.5 to 5 L/1000 L water/Ha)  or  1/3 fl. oz. to 2/3 fl. oz. per gal. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply more than 6 times per crop cycle or year.  <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Do not apply more than 6 times per crop cycle or year.
	Soil Drench	1 to 2 qts./100 gal. water (1/3 fl. oz. to 2/3 fl. oz. per gallon) 2.5 to 5 L/1000 L water	Apply 1 gal. of diluted solution per sq. yard and irrigate. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year.
	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications.

(Phytophthora)	Root Dip	2 qt./100 gal. water (2/3 fl. oz./gal.) 5 L/1000 L water	Apply as a pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
----------------	----------	----------------------------------------------------------	-----------------------------------------------------------------------------------------------

Do not graze livestock in treated conifer nurseries and/or plantations. Do not feed forage to livestock from treated nurseries and/or plantations.

**TURF**

FOSPHITE is a systematic fungicide, which is used for the control of Pythium blight, Summer Complex (Bermuda and others), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose and damping-off diseases to turf grasses on sod farms, commercial turf production, golf courses, parks and commercial landscapes. Begin preventative applications when conditions first favor disease and repeat at recommended intervals. Use higher rates under severe disease conditions.

Disease	Application Method	Rate	Application Program
Pythium (Yellow Turf)	Foliar Spray	3-6 fl. oz./1,000 sq. ft.	Apply recommended amount in 4 to 10 gal. of water per 1,000 sq. ft. Apply at 10 to 21 day intervals as needed. Do not mow or irrigate treated areas until sprays have completely dried.
Rhizoctonia Sclerotinia Helminthosporium Alternaria Anthracnose (Colletotrichum graminicola)		or	
		1 to 3 qts./100 gal. of water/acre. (5 to 7 L/1000 L of water / Ha)	For sod farms and large areas: Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.
	Irrigation	Overhead: 2-3 qts./acre. (5 - 7 L/ Ha) in at least 100 gallons (1000 L) of water.	Apply with normal irrigation schedule. Apply at 10 to 21 day intervals and repeat as needed.

\*Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, keep out of smoke.

**NOTICE TO BUYER**

To the extent permitted by law, all conditions and warranties, and statutory or other rights of action which buyer or any other user may have against JH Biotech, Inc. or seller are hereby excluded. JH Biotech, Inc. Hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by JH Biotech, Inc. or on its behalf unless it is established that such information or advice was provided negligently and that the product was been used strictly as directed, JH Biotech, Inc.'s liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price thereof.

24/33

**SUBLABEL - HOUSEHOLD PRODUCTS: FOR NON-COMMERCIAL APPLICATORS  
(NOT SUBJECT TO WORKER PROTECTION STANDARD)**

**FOSPHITE FUNGICIDE**

Fosphite is a systemic fungicide for the control of Downy mildew, Phytophthora, Pythium and various other diseases in garden plants, ornamental trees and plants, Phytophthora in conifers, pines and oaks and Pythium in turf.

This package can treat up to \_\_\_\_\_ sq. ft. (acres) at the maximum treatment rate.

**ACTIVE INGREDIENTS:**

Mono- and dipotassium salts of Phosphorous Acid\* .....53%

OTHER INGREDIENTS:.....47%

TOTAL.....100%

\*Contains 6.22 lbs./gallon of active ingredients, mono- and dipotassium salts of Phosphorous Acid. Equivalent to 3.90 lbs. Phosphorous Acid/gallon.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side (or back) panel for First Aid and additional precautionary statements.

**NET CONTENTS:**

- 1 pint (0.473 liters)
- 1 quart ( 0.946 liters)
- 1 gallon (3.785 liters)
- 2 1/2 gallons (9.463 liters)
- 5 gallons (18.925 liters)

**MANUFACTURED BY:**

JH Biotech, Inc.  
P.O. Box 3538  
Ventura, CA 93006

EPA Reg. No. 68573-2  
EPA Est. No. 68573-CA001



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

<b>FIRST AID STATEMENTS</b>	
If on skin or clothing	<ul style="list-style-type: none"> <li>·Take off contaminated clothing.</li> <li>·Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>·Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>·Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>·Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>·Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency medical treatment information.</p>	

**ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of rinsate or equipment wash waters.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Read entire label before using this product.

**STORAGE AND DISPOSAL**

**STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry place.

**PESTICIDE AND CONTAINER DISPOSAL:** **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused products down any indoor or outdoor drain.

### GENERAL APPLICATION INSTRUCTIONS

FOSPHITE is applied by various application methods, including foliar sprays, soil drench, soil incorporation, and bare root dip. For foliar sprays, apply FOSPHITE with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

### MIXING INSTRUCTIONS

1. Fill the spray tank with 1/2 of the required volume of water.
2. Add the amount of FOSPHITE as indicated under the Rate portion of the table below.
3. Continue to fill the tank with water to the desired volume while agitating.
4. Continue agitation when applying.

### INSTRUCTIONS FOR APPLICATIONS

#### ORNAMENTALS, BEDDING PLANTS AND TREES (CONIFERS, OAKS, ETC.)

FOSPHITE fungicide is used for control of Pythium, Downy Mildew, Phytophthora and various other diseases (see list below) of ORNAMENTAL PLANTS AND TREES. Plants, which can be treated with FOSPHITE, include (but not limited to):

#### ORNAMENTALS

Ageratum, Algerian Ivy, Aglaonema, Alyssum, Anthurium, Aphelandra, Arborvitae, Artemisia, Aster, Azaleas, Baby's Breath, Begonia, Bougainvillea, Boxwood, Caladium, Carnation, *Cattleya skinneri*, Ceanothus, Chrysanthemum, Coleus, Columbine, Cotoneaster, Cissus, Daisy, Delphinium, Dieffenbachia, Dogwood, Easter Lily, English Ivy, Ficus, Foxglove, Gaillardia, Geranium, Gloxinia, Hibiscus, Impatiens, Japanese Holly, Juniper, Leather-leaf Fern, Marigold, Monterey Pine, Pansy, Petunia, Peperomia, Philodendron, Phlox, Photinia, Pinks, Pittosporum, Pieris, Poinsettia, Pothos, Primrose, Prostrate Rosemary, Rhododendron, Roses, Salvia, Schefflera, Sedum, Sempervivum, Snapdragon, Spathiphyllum, Taxus media, Verbena, Vinca, White Cedar, White Pine, Zinnia, Zygocactus

**TREES:** Ash, Firs, Magnolia, Maple, Oaks, Spruces, Sycamore, Pines, etc.

Make applications prior to disease development. Use the higher rate when disease pressure is severe.

Apply FOSPHITE prior to disease development in conjunction with good cultural practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or under heat or moisture stress. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant.

Disease	Application Method	Rate	Application Program
Phytophthora Fusarium Pythium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> )	Foliar Spray	1/3 fl. oz. to 2/3 fl. oz. per gal. of water. (2.5 - 5 ml/L water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per year.

	Soil Drench	1/3 fl. oz. to 2/3 fl.oz. per gal. of water. (2.5 - 5 ml/L water)	Apply 1 gal. of solution per sq. yard and irrigate. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per year.
	Soil Incorporation	1 to 2 pts./cubic yard. (300-600 ml/cubic meter)	Mix 1 to 2 pints of Fosphite with enough water to wet one cubic yard of soil; mix immediately before potting. If conditions are favorable for disease development, it is recommended to make additional applications as a foliar spray or soil drench.
	Root Dip	2/3 fl. oz./gal. of water (5 ml/L water)	Dip transplants into diluted solution of Fosphite to thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
Downy Mildew	Foliar Spray	1/3 fl. oz. to 2/3 fl. oz. per gal. of water (2.5 - 5 ml/L water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per year.

**TURF**

FOSPHITE is a systematic fungicide, which is used for the control of Pythium blight, Summer Complex (Bermuda and others), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose and damping-off diseases to turf grasses. Begin preventative applications when conditions first favor disease and repeat at recommended intervals. Use higher rates under severe disease conditions.

Disease	Application Method	Rate	Application Program
Pythium (Yellow Turf) Rhizoctonia Sclerotinia Helminthosporium Alternaria Anthracnose (Colletotrichum graminicola)	Foliar Spray	3-6 fl. oz./1,000 sq. ft.	Apply recommended amount in 4-10 gal. of water per 1,000 sq. ft. Apply at 7 to 21 day intervals as needed. Do not mow or irrigate treated areas until sprays have completely dried.

### GARDEN CROPS

FOSPHITE fungicide is used for control of Downy Mildew, Pythium, Phytophthora and various other diseases (such as purple blotch, late blight, blue mold, soft rot, root rot, brown rot, foot rot, crown/collar rot, canker, fruit/foliar rot, fire blight, heart rot, pink rot, pythium leak, red stele, leather rot, etc.) of garden crops. FOSPHITE may be applied to plants including (but not limited to): Almond, Apple, Apricot, Artichoke, Asparagus, Avocado, Baby Lettuce, Banana, Beet, Blackberry, Blueberry, Broadbean, Cane Berries, Carrot, Cherry, Chestnut, Chilies, Cinchona, Cinnamon, Citrus, Cole Crops, Corn, Cranberry, Crucifers, Cucurbits, Dates, Dry Beans, Eggplant, Fresh beans, Garlic, Grapes, Kiwi, Lettuce, Lima Beans, Macadamia, Melons, Mint, Nectarine, Nut Crops, Nutmeg, Okra, Olive, Onion, Papaya, Peach, Peanut, Pear, Peas, Peppers, Pineapple, Plum, Pome Fruit, Potato, Prune, Raspberry, Rhubarb, Root crops, Rosemary, Spinach, Squash, Strawberry, String Beans, Stone Fruit, Sugarbeet, Sunflower, Tomato, Walnut.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply FOSPHITE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Rhizoctonia Pythium Fusarium Fire Blight ( <i>Erwinia amylovora</i> )	Foliar	1 to 2 qts./ 100 gal. of water or 1/3 fl. oz. to 2/3 fl. oz. per gal. (2.5 to 5 L/1000/L)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per year.
Downy Mildew	Foliar	1 to 2 qts. (1/3 fl. oz. to 2/3 fl. oz. per gal.) 2.5 to 5 L/1000 L	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per year.

### NOTICE TO BUYER

To the extent permitted by law, all conditions and warranties, and statutory or other rights of action which buyer or any other user may have against JH Biotech, Inc. or seller are hereby excluded. JH Biotech, Inc. Hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by JH Biotech, Inc. or on its behalf unless it is established that such information or advice was provided negligently and that the product was been used strictly as directed, JH Biotech, Inc.'s liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price thereof.

29/33

**SUBLABEL - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS  
SUBJECT TO WORKER PROTECTION STANDARD  
(TRUNK INJECTION ONLY)**

**FOSPHITE FUNGICIDE**

**A FUNGICIDE FOR THE CONTROL OF DOWNY MILDEW, PHYTOPHTHORA, PYTHIUM  
AND VARIOUS OTHER DISEASE ON AGRICULTURAL TREES, INDOOR AND OUTDOOR  
ORNAMENTALS, FORESTREY APPLICATIONS INCLUDING DOMESTIC TREES,  
CONIFERS, PINES, AND OAKS**

This package can treat up to \_\_\_\_\_ sq. ft. (acres) at the maximum treatment rate.

**ACTIVE INGREDIENTS:**

Mono- and dipotassium salts of Phosphorous Acid\* ..... 53%

**OTHER INGREDIENTS:**..... 47%

**TOTAL**..... 100%

\*Contains 6.22 lbs./gallon of active ingredients, mono- and dipotassium salts of Phosphorous Acid. Equivalent to 3.90 lbs. Phosphorous Acid/gallon.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side (or back) panel for First Aid and additional precautionary statements.

**NET CONTENTS:**

- 1 pint (0.473 liters)
- 1 quart ( 0.946 liters)
- 1 gallon (3.785 liters)
- 2 1/2 gallons (9.463 liters)
- 5 gallons (18.925 liters)

**MANUFACTURED BY:**  
JH Biotech, Inc.  
P.O. Box 3538  
Ventura, CA 93006

EPA Reg. No. 68573-2  
EPA Est. No. 68573-CA001

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after using.

<b>FIRST AID STATEMENTS</b>	
If on skin or clothing	<ul style="list-style-type: none"> <li>·Take off contaminated clothing.</li> <li>·Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>·Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>·Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>·Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>·Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency medical treatment information.</p>	

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves, shoes plus socks and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS BOX**

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Read entire label before using this product.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS:**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, socks, shoes, and chemical resistant gloves.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

**GENERAL APPLICATION INSTRUCTIONS**

FOSPHITE is a systemic fungicide for the prevention and treatment of downy mildew, Sudden Oak Death (*Phytophthora ramorum*), root rots (*Phytophthora spp.*), pythium and various other disease of agricultural trees, indoor and outdoor ornamentals, forestry applications including domestic trees, conifers, pines and oaks.

**COMPATIBILITY**

Use only FOSPHITE alone when used as a trunk injection. To avoid undesirable phytotoxicity, do not apply FOSPHITE to trees treated with other pesticides or copper based compounds at intervals less than 20 days, unless instructed to do so by your consultant.

**GENERAL INSTRUCTIONS FOR APPLICATIONS**

**FOOD CROPS**

FOSPHITE fungicide is used for control of Downy Mildew, *Erwinia amylovora*, Pythium, Fusarium, Rhizoctonia and Phytophthora diseases (such as purple blotch, late blight, blue mold, soft rot, root rot, brown rot, foot rot, crown/collar rot, canker, fruit/foliar rot, fire blight, heart rot, pink rot, pythium leak, red stele, leather rot, etc.) of agronomic crops. FOSPHITE may be applied to plants including:

**Citrus Crop Grouping (Bearing and Non-Bearing) including Avocado:** Avocado, Calamondin, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange (all types), Pummelo and Satsuma mandarin.

**Pome Fruits Crop Grouping: (Bearing and Non-Bearing)** Apple, Crabapple, Loquat, Mayhaw, Pear, Quince.

**Stone Fruits Crop Grouping:** Apricot, Cherry, Nectarine, Peach, Plum, Plumcot and Prune.

**Tree Nuts and Vines Crop Grouping:** Almond, Beech nut Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Grapes, Hickory nut, Macadamia nut, Pecan, Pistachio and Walnut.

**ORNAMENTALS**

FOSPHITE fungicide is used for control of Pythium, Downy Mildew, Phytophthora and various other disease diseases of ORNAMENTALS grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. FOSPHITE may be applied to ornamental and landscape plants including: Boxwood, Cedars, Ceanothus, Dogwood, Ficus, Flowering Trees, Hard Wood Trees, Juniper, Magnolia Trees, Pines, Oaks, Palms, Photinia, Pieris, Red woods and Shrubs,

**CONIFERS AND TREES IN COMMERCIAL NURSERIES, PLANTATIONS AND FORESTS,  
LANDSCAPES AND PARKS  
(INCLUDING CHRISTMAS TREES, OAKS and PINES)**

FOSPHITE applied in conjunction with good cultural management practices will control Phytophthora root rot of conifers including (but not limited to): Douglas Fir, spruce and pines also other Phytophthora disease of Oaks (Sudden Oak Death) and forest trees. Use the higher rates when disease pressure is severe. Do not apply to conifers that are under heat or moisture stress.

Disease	Application Method	Rate	Application Program
Phytophthora Fusarium Pythium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> ) Bacterial Blight ( <i>Xanthomonas campestris</i> )	Trunk Injection	3 teaspoons / linear yard of canopy width or 2 in. of trunk diameter at breast height (DBH) (15 ml / linear meter of canopy width or 5 cm DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications.



**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage	Store in a cool, dry place.
Pesticide Disposal	Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal	Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

**NOTICE TO BUYER**

To the extent permitted by law, all conditions and warranties, and statutory or other rights of action which buyer or any other user may have against JH Biotech, Inc. or seller are hereby excluded. JH Biotech, Inc. Hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by JH Biotech, Inc. or on its behalf unless it is established that such information or advice was provided negligently and that the product was been used strictly as directed, JH Biotech, Inc.'s liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price thereof.