# RSR MITE AND LICE SPRAY

# A WATER-BASED BIRD SPRAY With Feather Conditioner & Cologne

#### KILLS LICE AND MITES ON PET BIRDS

ACTIVE INGREDIENTS:	
Pyrethrins	0.03%
*Piperonyl Butoxide, Technical	
INERT INGREDIENTS:	
	100.00%

<sup>\*</sup>Equivalent to 0.24% (butylcarbityl) (6-propylpiperonyl) ether and 0.06% related compounds.

#### KEEP OUT OF REACH OF CHILDREN

# **CAUTION**

See Back Panel for Precautionary Statements

MANUFACTURED BY: SPEER PRODUCTS, INC. P. O. BOX 18993 MEMPHIS, TENNESSEE 38181-0993

NET CONTENTS\_\_\_\_OZ

EPA Est. 11715-TN-1

EPA Reg. No. 68688-46

## PRECAUTIONARY STATEMENTS


# CAUTION -

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing vapors. Wash hands with soap and warm water after use. In the home, all food processing surfaces, exposed food and utensils should be covered during treatment, or thoroughly washed before use.

<ul> <li>Call poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> </ul>
<ul> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor fot treatment advice.</li> </ul>

Have the producct container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-757-4943 for emergency medical treatment information.



# PHYSICAL HAZARDS

Do not use or store near heat or open flame.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

# SHAKE WELL BEFORE USING HOLD CONTAINER UPRIGHT

TO KILL LICE AND MITES ON BIRDS: For light infestation of lice or mites, mist birds twice a week until lightly moist, especially at tip of tail. Cover cage with cloth until bird is dry. For severe mite infestation, remove bird from cage and spray perches thoroughly. Clean cages thoroughly at regular intervals.

Keep birds out of cold drafty areas. Do not spray sick birds or birds with open lesions. Avoid spraying in eyes and nares.

#### STORAGE AND DISPOSAL

STORAGE: Store in a cool area away from heat or open flame.

**DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

c\mspub\rsr.46.pub(2/95)(11/26/02 NOT1F. 2001 1&6)