PO MERIMATINE CATE TO CO

For Systemic Insect Control on Landscape Ornamentals and Interior Plantscapes

[Tree Injection]
[Convenient, Easy-to-Use]
[Premeasured Doses]

Controls [{Choose from Listed Pests}]

{Listed Pests:} Adelgids, Armored Scales (suppression), Aphids, Elm Leaf Beetles,
Black Vine Weevil Larvae, Eucalyptus Longhorned Borer, Flathead Borers
(including Bronze Birch Borer and Alder-Birch Borer), Japanese Beetles, Lacebugs, Leafhoppers,
Leafminers, Mealybugs, Sawfly Larvae, Pine Tip Moth Larvae, Psyllids, Royal Palm Bugs, Soft Scale
Insects, Thrips (Suppression), White Grub Larvae (such as Japanese Beetle Larvae, Chafers,
Phyllophaga spp., Asiatic Garden Beetle and Oriental Beetle) and Whiteflies
On All Types of Ornamental Trees

Active Ingredient:	
Imidacloprid (CAS No. 138261-41-3)	5%
Other Ingredients	<u>95%</u>
Total	
Contains 6 grams (0.2 oz.) active ingredient per 120 ml (4 oz.) bottle.

Keep Out of Reach of Children WARNING

See below [outsert] [back panel] for additional precautionary statements.

Net Wt. 4 oz. (120 ml) [ampules, Net Wt. 0.36 oz. each, oz. total]		6 L 0 1 6 C C
EPA Reg. No. 69117-1	EPA Est	0

ACCEPTED
JUN 1 7 2002
Under the Reduct Inspeticide,
Pengicide, and Rodenficide Act, as emended, for the penticide
Chair the Pedend Insecticite, President, and Rodenticite Act, as assended, for the posticite Registered tracky IDA Reg. No. 69/1/7-/

[Sold by]
ArborSystems
Omaha, NE 68134

69117-1 **New Label**: Page 2 of 4

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

WARNING: Harmful if swallowed, inhaled or absorbed through skin. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear safety glasses. Thoroughly wash with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

First Aid

If Swallowed	 Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
if inhaled	Move person to fresh air.
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
İ	Call a poison control center or doctor for further treatment advice.
if on Skin	Take off contaminated clothing.
or Clothing	Immediately rinse skin with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If in Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
	ct container or label with you when calling a poison control center or doctor, or going for treatment. You ct 1-800-698-4641 for emergency medical treatment information.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Environmental Hazards

This product is highly toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming plants or weeds. Do not apply this product or allow it to drift to blooming plants or weeds if bees are visiting the treatment area.

For assistance, contact ArborSystems at 402-339-4459

(Professional Use Label)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For Use By Professional Aborists/Applicators

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store above 65°F in a cool, dry place and in such a manner as to prevent cross contamination with fertilizers, food, feed and other pesticides. Store in original container out of reach of children, preferably in a locked storage area. Disposal: Do not reuse empty container; wrap and put in trash.

GENERAL INFORMATION Do not Apply to Food Bearing Plants

ArborSystems Tree Injection System using the Patented WedgleTM Tip and WedgeChekTM
The ArborSystems Tree Injection System injects pesticides or micronutrients into a tree with minimal wounding. This system injects chemicals into the area of bark separation where the chemical can actively be transferred to other parts of the tree. Injections made during the spring or summer will be more successful than waiting until the fall. Specifically, each tree needs to be inspected. The bark needs to be pliable and that depends upon local weather conditions.

Note: When using Pointer Insecticide, the ArborSystems Tree Injection System must be cleaned periodically to prevent

clogging.

How to Use the ArborSystems Tree Injection System

- 1. Use protective eyewear and rubber or neoprene gloves when handling or installing the ArborSystems Tree Injection System.
- 2. Attach the chemical bottle to the SW100 Injector Gun.

3. Attach the Wedgle Tip to the SW100 Injector Gun. Make sure the small hole in the tip is pointing straight up. Set the delivery volume on the syringe to 1 ml or 1/2 ml, as appropriate.

4. Determine the thickness of the bark. Insert the Wedgle Tip into the fissure of the tree. If the tip penetrates more than the length of a WedgeChek, the injection will be made in the fissure. If the tip penetrates less than the length of a WedgeChek, the injection will be made through the tight bark of the tree. On thin-barked trees, you may find it necessary to insert the WedgeChek in only one-half as far as normal.

Insert the WedgeChek™ Punch into the tree. Remove the punch and squeeze the trigger to remove the bark core. Examine the core to make sure the core that has been removed is approximately the same length as the WedgeChek. Repeat this step while moving around the flare of the tree. Be sure to follow the label directions on this outsert to determine how many injection sites are needed on any given tree. Usually this is approximately 1 injection for every 4" to 6" of trunk circumference measured at 4.5' above the ground. Make WedgeChek insertions around the flare of the tree spacing evenly.

Place the WedgeChek on the protruding pin on the bottom of the grip then insert the WedgeChek directly into the

punched hole. The barbs on the WedgeChek will secure it in the tree.

7. Insert the Wedgle Tip with a straight motion through the WedgeChek. Make sure the hole in the tip passes completely through the WedgeChek. As the tip is pushed through the WedgeChek and the bark the blunt-edged tip automatically stops when it reaches the wood.

- 8. With a smooth motion, firmly squeeze both handles at the same time. This allows the bark to separate and the chemical to be deposited into the area. The WedgeChek plugs the hole, allowing the chemical to remain in the tree.
- 9. Slowly withdraw the tip and repeat this step while moving around the flare of the tree.
- 10. Move laterally and make the next injection until the entire tree trunk has been treated. After the tree has taken up the chemical, each WedgeChek can either be removed or left in the tree.

Pests Controlled Adelgids, Armored Scales (suppression), Aphids, Elm Leaf Beetles, Black Vine Weevil Larvae, Eucalyptus Longhorned Borer, Flatheaded Borers (including bronze birch borer and alder-birch borer), Japanese Beetles, Lacebugs, Leafhoppers, Leafminers, Mealybugs, Sawfly Larvae, Pine Tip Moth Larvae, Psyllids, Royal Palm Bugs, Soft Scale Insects, Thrips (suppression), White Grub Larvae (such as Japanese beetle larvae, Chafers, Phyllophaga spp., Asiatic Garden Beetle and

Dosage

1 ml per 4" to 6" of trunk circumference measured at 4.5' above the ground.

Note: Depending on type of bark, the dosage per injection site may have to be reduced to ½ ml per site and the number of injection sites doubled.

Cross-section of tree showing Wedgle™ Tip and WedgeChek™

{Domestic/Homeowner Use Label -- Ampules}

Oriental Beetle) and Whiteflies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. .

Storage and Disposal

Storage: Store above 65°F in a cool, dry place and in such a manner as to prevent cross contamination with fertilizers, food, feed and other pesticides. Store in original container out of reach of children, preferably in a locked storage area. Disposal: If empty: Do not reuse ampules. Place in trash or offer for recycling if available. If partly filled: Call your local solid or household waste agency [or 1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION Tree Insecticide injectors Do not Apply to Food Bearing Plants

This product contains an insecticide that is injected into holes drilled into the tree trunk at the flare of the tree. The insecticide controls Adelgids, Armored Scales (suppression), Aphids, Elm Leaf Beetles, Black Vine Weevil Larvae, Eucalyptus Longhorned Borer, Flatheaded Borers (including bronze birch borer and alder-birch borer), Japanese Beetles, Lacebugs, Leafhoppers, Leafminers, Mealybugs, Pine Tip Moth Larvae, Psyllids, Royal Palm Bugs, Soft Scale Insects, Sawfly Larvae, Thrips (suppression), White Grub Larvae (such as Japanese beetle larvae, Chafers, *Phyllophaga* spp., Asiatic Garden Beetle and Oriental Beetle) and Whiteflies on all types of ornamental trees. The pre-measured doses are convenient and easy to use.

{Illustration of measuring tree}	{Illustration of tree and drilling hole}	{Illustration of product application}
Tools: Tape measure and 3/8" drill. Dosage: Measure the circumference of the tree trunk at 4 feet above the ground. You will need one ampule (0.36 oz.) for every four to six inches of circumference.	instructions: Drill 1/2" deep holes into the tree at the flare of the tree. Drill the holes at a downward slant (about 45°). Evenly space the holes around the tree. Use only a clean, sharp drill bit. Brad point drill bits are best.	Use rubber or neoprene gloves when handling this product. Cut off the tip and squeeze the chemical into the holes. Use one ampule for each hole.

Treatments are more effective early in the growing season. Injections to conifers may be either slow in uptake or have no uptake at all.

Note: Holes drilled in a tree are wounds. Do not drill holes (three-eighths (3/8) inch and larger at 3-inch spacing) in a tree more than once a year or more than two consecutive years.

{Professional Use and Domestic/Homeowner Use Labels}

Notice of Warranty

ArborSystems warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. ArborSystems does not make or authorize any agent or representative to make any other warranty, guarantee or representation, express or implied, concerning this product.

For assistance contact

ArborSystems

P.O. Box 34645 Omaha, NE 68134

Toll-Free: 1-800-698-4641 Phone: (402) 339-4459 FAX: (402) 339-5011

www.arborsystems.com

Pointer™ Insecticide, Wedgle™ Tip and WedgeChek™ are registered trademarks of ArborSystems

Wedgle™ Tip is protected by U.S. Patent #5,239,773

WedgeChek™ is protected by U.S. Patent #5,797,215

l	1	Denotes alternate/optional language
1	1	Denotes language that does not appear on the market labeling