69117-		•	28-2002	-			ų – J
Please read instructions on reverse before completing form. United States Environmental Protection Agency 1200 Pennsylvania Avenue, N.W. Washington, DC 20460-0001			>y	<u> </u>	Registration Amendment Other	OPP Ider	MB No. 2070-0060.
		pplication fo	or Pesticide –				
1. Company/Product Number EPA Reg. No. 69117-2			2. EPA Prod	uct Manag nas C. H		Proposed Cl	assification
4. Company/ Product (Nam Grey	^{ie)} nound™ Insectici	ide	PM #			None	Restricted
5. Name and Address of Applicant (Include ZIP Code) ArborSystems, LLC c/o RegWest Company 30856 Rocky Road Greeley, CO 80631-9375			3(c)(3)(b)(i) labeling to:	 6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to: EPA Reg. No 			
	eck if this is a new add	ress	Product Na	me			
		S	ection – II				
Amendment – Explain I	pelow.		Final prir	nted labels	in response to Ag	ency letter da	ted
Resubmission in respon	nse to Agency letter da	ted	🛄 "Me Too	" Applicatio	on.		
Notification - Explain be				Explain be	low.	NOTIFIC	ATION
Explanation: Use add	litional page(s) if neces	ssary. (For Section	I and Section II.)			NUTIFIC	AUON
penalties under sections	12 and 14 of FIFRA		ection – III				
1 Material This Product V	Vill Be Packaged In:	3	ection – III				
❑ Yes* ❑ Yes ❑ Y ❑ No ❑ No ❑ N		Water Soluble Packa Yes No	es Q Metal Q Plastic Q Class				
*Certification must be submitted	If "Yes" Unit Packaging Wgt.	No. Per Container	lf "Yes" Packaging Wgt.	No. Pe Contair	ner 🖸 Pape		
3. Location of Net Contents	Information	4. Size(s) Retail				ocation of Label Directions	
Label Container				On Label On Labeling Accompanying Product		ing Product	
6. Manner in Which Label is Affixed to Product Lithog		ograph					
			ection - IV				
1. Contact Point (Comp	lete items directly be			e contact	ed, if necessary,	to process th	nis application).
Name Title		,, ,, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Telephone No. (Include Area Cod		Code)		
Kim Davis, CC, APC			Consultant/Agent		((970) 353-0611	
I certify that the state accurate and complete punishable by fine or in	e. I acknowledge :	that any knowin th under applic	rm and all attach ngly false or misle cable law.			e, Rece	Application lived Stamped)
2. Signature		3. Title	3. Title			·	
4. Typed Name		5. Dat	Consultant/Agent 5. Date			\neg .	
Kim Davis, CC, APC May 20, 2002							
EPA Form 8570-1 (Rev. 8-9	Previous editions ar	e obsolete.					

.

69117-2: Page 1 of 4 04/06/01 SAL + 05/20/02 NOTIF XXXXX = addition; XXXXX = deletion

{Box and 4-ounce Container Label}

GREYHOUND™ INSECTICIDE

For Use by Professional Arborists/Applicators

ACTIVE INGREDIENT:

> Keep Out of Reach of Children WARNING See <u>insert</u> <u>outsert</u> for additional [complete] precautionary statements.

NOTIFICATION

MAY 2.8 2002

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with fertilizers, food, feed and other pesticides. Store in original container out of reach of children, preferably in a locked storage area. **Disposal:** Do not reuse empty container; wrap and put in trash.

Refer to insert Outsert for complete Directions for Use.

Net Contents 4 oz. (120 ml)

EPA Reg. No. 69117-2 EPA Est. _____

[Sold by] ArborSystems, LLC Omaha, NE 68134 www.arborsystems.com

{Insert Outsert Label }

GREYHOUND INSECTICIDE

For Systemic Insect Control on Landscape Ornamentals and Interior Plantscapes ' For Use by Professional Arborists/Applicators

ACTIVE INGREDIENT: Abamectin B ₁ *	
OTHER INGREDIENTS	
Total	100%
*A mixture of avermectins containing \ge 80% avermectin avermectin A _{1a}) and \le 20% avermectin B _{1b} (5-0-demet propyI)-25-(1-methylethyl) avermectin A _{1a})	
Contains 0.0047 pound Abamectin per 4 oz. bo	ottle

69117-2: Page 2 of 04/06/01 SAL + 05/20/02 NOTIF XXXXX = addition; XXXXX = deletion

Keep Out of Reach of Children WARNING See inside below for additional [complete]

precautionary statements.

Net Contents 4 oz. (120 ml)

EPA Reg. No. 69117-2 EPA Est.

[Sold by]

ArborSystems, LLC Omaha, NE 68134

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals. Harmful if inhaled or absorbed through the skin. Avoid contact with skin. May be fatal if swallowed. Use protective eyewear and rubber or neoprene gloves. Remove

contaminated clothing and wash before reuse. Thoroughly wash with soap and water after handling and before eating, drinking or using tobacco.

entra a la ba

 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Call a poison control center or doctor for treatment advice. Immediately give one or two glasses of water and induce vomiting by gently touching back of the throat with finger. Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled 		First Aid
 Call a poison control center or doctor for treatment advice. f on Skin or Clothing Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Call a poison control center or doctor for treatment advice. Immediately call a poison control center or doctor for treatment advice. Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicater by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines,	If in Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Immediately give one or two glasses of water and induce vomiting by gently touching back of the throat with finger. Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. A doctor for the treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor for further treatment advice. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicater(try clinica		Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Immediately give one or two glasses of water and induce vomiting by gently touching back of the throat with finger. Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Abve person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. 		 Call a poison control center or doctor for treatment advice.
 Call a poison control center or doctor for treatment advice. f Swallowed Immediately give one or two glasses of water and induce vomiting by gently touching back of the throat with finger. Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.	If on Skin	Take off contaminated clothing.
 Immediately give one or two glasses of water and induce vomiting by gently touching back of the throat with finger. Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.	or Clothing	 Immediately rinse skin with plenty of water for 15-20 minutes.
 Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting or give anything by mouth to an unconscious person. f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicater by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.		 Call a poison control center or doctor for treatment advice.
 Do not induce vomiting or give anything by mouth to an unconscious person. finhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.	If Swallowed	I minimute latery give one of the glabbee of match and induce formany by gently todorning back of the
 f Inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure. 		 Immediately call a poison control center or doctor for treatment advice.
 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.		 Do not induce vomiting or give anything by mouth to an unconscious person.
mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half nour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.	If Inhaled	Move person to fresh air.
Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.		• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle tremors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.		mouth-to-mouth, if possible.
may also contact 1-800-698-4641 for emergency medical treatment information. Note to Physician: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.		
remors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.		
wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.	tremors. Toxic hour of exposur electrolyte imba along with other	ity following accidental ingestion of this product can be minimized by inducing vomiting within one half e. If toxicity from exposure has progressed to cause severe vomiting the extent of resultant fluid and lance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical
	clinical condition wise to avoid	n is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Environmental Hazards

This product is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Do not apply this product to blooming crops if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

69117-2: Page 3 of 4 04/06/01 SAL + 05/20/02 NOTIF XXXXX = addition; XXXXX = deletion

For assistance, contact ArborSystems at 402-339-4459.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT USE ON CROP-BEARING TREES

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with fertilizers, food, feed and other pesticides. Store in original container out of reach of children, preferably in a locked storage area. **Disposal:** Do not reuse empty container; wrap and put in trash.

GENERAL INFORMATION

ArborSystems Injection System using the Patented Wedgle™ Tip and WedgeChek™


The ArborSystems injection system injects pesticides or micronutrients into a tree with minimal wounding. The system will inject chemicals into the area of bark separation where the chemical can actively be transferred to other parts of the tree. Spring through late summer is the ideal time for injection. Injections made during the spring or summer will be more successful than waiting until the fall. Specifically, each tree needs to be inspected. The bark needs to be pliable and that depends upon local weather conditions.

Injections can be made either in anticipation of pest presence or after pests are observed. Pest control is anticipated to last a typical growing season, although it may be recommended to inject the plants one or two times per year or more often under severe and repeated pest pressure. This product has been evaluated for phytotoxicity on a wide range of ornamental plants. However, since all ornamental plants have not been tested, it is recommended that one plant be treated first to make certain that no phytotoxicity occurs. Do not use on sick plants or plants weakened by extreme environmental conditions such as heat, flooding, cold, etc.

How to Use the ArborSystems Injection System

- 1. Use protective eyewear and rubber or neoprene gloves when handling or installing the ArborSystems injection system.
- 2. Remove the small tab on top of the bottle and attach the bottle of chemical to the injector gun.
- 3. Attach the Wedgle[™] Tip to the injector gun. Make sure the small hole in the tip is pointing straight up. Set the delivery volume on the syringe to 1 ml or ½ ml, as appropriate.
- 4. Determine the thickness of the bark. Insert the Wedgle[™] Tip into the fissure or valley of the tree. If the tip penetrates more than the length of a WedgeChek[™], the injection will be made in the fissure. If the tip penetrates less than the length of a WedgeChek[™], the injection will be made through the tight outer bark of the tree.
- 5. Insert the WedgeChek[™] Punch into the tree. Rotate the punch to separate the core from the tree bark and remove the punch from the tree. Squeeze the trigger to remove the bark core and make sure the removed core is as long as the WedgeChek[™]. Repeat this step while moving around the circumference of the tree. Be sure to follow the label directions on this insert the chemical bottle outsert to determine how many injection sites are needed on any given tree. Usually this is approximately 1 injection for every 4 to 6 inches of circumference.
- 6. Put the WedgeChek[™] on the Wedgle[™] Tip, prior to inserting the tip into the tree.
- 7. Using a straight motion insert the Wedgle[™] Tip into the tree until it stops against the wood. This will usually be about ½" on Elms, shorter on crabapple, oak and sycamore trees. As the tip is pushed into the tree, the Wedgle[™] Tip cuts through the WedgeChek[™] and the blunt-edged tip automatically stops when it reaches the wood.
- Using a smooth motion, firmly squeeze both handles at the same time. This motion allows the bark to separate and the chemical to be deposited into the area. The WedgeChek[™] plugs the hole, allowing the chemical to remain in the tree.
- 9. Slowly withdraw the tip and repeat this step while moving around the circumference of the tree.
- 10. Move laterally and make the next injection until the entire tree trunk has been treated. The WedgeCheks™ may be left in the tree and used to inject the tree the following year.

Indications	Dosage
Indicated for the treatment of woody trees and shrubs for the control of Elm Leaf Beetles and Lace Bugs	1 mt per 4" to 6" of trunk circumference. Note: Depending on type of bark, the dosage per injection site may have to be reduced to ½ ml per site and the number of injection sites doubled.



Injection Into the Fissures (Not Bark) of the Tree

Notice of Warranty

ArborSystems, **LLC** warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. ArborSystems, **LLC** does not make or authorize any agent or representative to make any other warranty, guarantee or representation, express or implied, concerning this product.

For assistance contact

ArborSystems, LLC

P.O. Box 34645 Omaha, NE 68134 Toll-Free 1-800-698-4641 Phone (402) 339-4459 FAX (402) 339-5011 www.arborsystemslc.com Greyhound[™] Insecticide, Wedgle[™] Tip and WedgeChek[™] are registered trademarks of ArborSystems, LLC Wedgle[™] Tip is protected by U.S. Patent #5,239,773 WedgeChek[™] is protected by U.S. Patent #5,797,215

•




.

3 1 1

BREYHOUND™ Insecticide BREYHOUND™ Insecticide To Use by Professional Arborists/Applicators ADD BY ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD BY ADD ADD BY ADD BY ADD BY ADD BY ADD BY ADD BY ADD ADD BY AD

•

.



,

81

9 10 10

.

•

e

.

*

GREYHOUND[™] INSECTICIDE

or Systemic Insect Control on Landscap Ornamentals and Interior Plantscapes For Use by Professional Arborists/Applicators

ACTIVE INGREDIENT:

.

.

Abamectin B ₁ *	
OTHER INGREDIENTS	5
	Total: 100%

*A mixture of avermectins containing \geq 80% avermectin 8₁₀ (5-0-demethyl avermectin A₁₀) and \leq 20% avermectin B_{1b} (5-0-demethyl avermectin B_{1b} (5-0-demethyl-25-del(1-methylproph))-25-(1-methylethyl) avermectin A₁₀.

Contains 0.0047 pound Abamectin per 4 oz. bottle.

Keep Out of Reach of Children WARNING

See inside for additional precautionary statements Net Contents 4 oz. (120 ml)

EPA Reg. No. 69117-2 EPA Est. 69117-NE-1

Sold by

ARDOR SYSTEMS Ömaha, NE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND

HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing Prolonged or frequently repeated exposure may cause allergic skin reactions in some indi-viduals. Harmful if inholed or absorbed through the skin. Avoid contact with skin. May be fatal if swallowed. Use protective eyewcar and rubber on reoprene gloves. Remove con-tominated clothing and wash before reuse. Wash thoroughly with soop and water diter handling and before enting, dimking or using tobacco.

FIRST AID

FIRST AID If in Eyes: Hold eye open and rinse skowly and gently with water for 15-20 minutes. Remove control tenses, if present, after the first 5 min-utes, then continue insing eye. Call a poison control center or doctor for treatment advice. If on Skin or Orthing: Take off contaminated (afthing. Immediately nakes skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If Swallowed: Immediately give one or two glasses of water and induce vorning by gently touching back of the throat with finger. Immediately call a poison control center or doc-tor for treatment advice. Do not induce vorning or give anything by mouth to an unconscious person. If Inhelad: Move person to fresh air. If person is not breathing, call 911 or an ambu-fance, then give artificial respiration, preferably by mouth-to-mouth, if possible, Call a poison control center or doctor for further treatment advice. odvice

Have the product container or label with you when calling a polson control center or doctor, or going for treatment. You may also contact 1-800-699-4641 for emergency medical treatment information.

I-bou-bye-doal to emergency medical integriment information.
Note to Physician: Early signs of intoacation include mydriasis (dikated pupils), ataxia (unsteadiness) and muscle tremors. Toxicity following accidental ingestion of this product can be minimized by inducing vomiting within one holf hour of exposure. It toxicity form exposure hose progressed to cause serve vomiting the extent of resultant fluid and electrolyte imbalance should be guede. Appropriate supportive parenteral fluid replacement therapy should be given along with other required supportive measures (such other required supportive cases, observations as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until the clinical condition is stable and normal. Since ubomechin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity in botentiaty toxic aborectin exposure.
ENVIROMENTAL HAZARDS

ENVIROMENTAL HAZARDS

ENVIRONMENTAL HAZARDS This product is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rin-sorte. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Do not apply this product to blooming crops if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS Do not use or store near heat or open flame DIRECTIONS FOR USE

It is a violation of Federal law to use this prod-uct in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE AND DISPOSAL Do Not Use on Crop-Bearing Trees. Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with fertilizers, food, feed and other pesticides. Store in origi-nal container out of reach of children, prefer-ably in a locked storage area. Disposal: Do not reuse empty container, wrap and put in trash.

continued on back

GENERAL INFORMATION

GENERAL INFORMATION AborSystems Injection System using the patented Wedgle" Tip and WedgeChek" The AborSystems Injection System injects pest-ades or mKoroutherts into a tree with minimal wounding. The system will inject chemicals into the area of bark separation where the chemical can actively be transferred to other parts of the ree. Injectors made during the spring or summer will be more successful than waiting until the fall. Specifically, each tree needs to be inspected. The bark needs to be plakele and that depends upon local weather conditions.

.

local weather conditions. Injections can be made either in amicipation of pest presence or after pests in an observed. Pest control is anticipated to last a typical growing seo-son, although it may be recommended to inject the plants one or two times per year or more often under server and repeated pest pressure. This product has been evaluated for phytotoxicity on a wide range of omamental plants. However, since all ornamental plants have not been tested, it is recommended that one plant be treated first to make certain that no phytotoxicity occurs. Do not use on side plants or plants weakened by extreme environmental conditions such as heat, flooding, cold, etc. HOW TO LISE THE ADBODEVETENCE

HOW TO USE THE ARBORSYSTEMS INJECTION SYSTEM:

- Use protective evenear and rubber or neo-prene gloves when handling or installing the Arbor Systems Injection System.
- Remove the small tab on top of the bottle and attach the bottle of chemical to the injector gun.
- mector gun.
 3. Attach the Wedgle^{III} Tip to the injector gun.
 Make sure the small hole in the tip is point-ing straight up. Set the delivery volume on the syringe to 1 ml or 1/2 ml, as appropriate
- the symple to I mi or 1/2 mi, as appropriate 4. Determine the thickness of the bark. Insert the Wedgle¹⁰. Tip into the fissure or valley of the tree. If the tip penetrates more than the length of a WedgeChek¹⁰, the night-on will be made in the fissure. If the tip penetrates less than the length of a WedgeChek¹⁰, the injection will be made through the tight outer bark of the tree.
- bark of the tree. 5. Insert the WedgeChek¹⁴ Punch into the tree. Rottle the punch to separate the core from the tree. Sulleast the trigger to remove the bark core and make sure the removed core is as long as the WedgeChek¹⁴. Repeat this step while moving around the circumference of the tree. Be sure to follow the bloed direc-tions on this insert to datermine how many injection sites a opprovincity Lingeton for every 4 to 6 inches of circumference.
- 6. Put the WedgeChek'" on the Wedgle'" Tip, prior to inserting the tip into the tree.
- prior to inserting the tip into the tree. 7. Using a straight motion insert the Wedgle²⁶. Tip into the tree until it stops against the wood. This will usually be about 1/2° on Elms, shorter on crabaple, ook and sycamore trees. As the tip is pushed into the tree, the Wedgle²⁶. Tip cuts through the WedgeChek⁸⁶ and the butth-edged tip auto-matically stops when it reaches the wood.
- Manucliny study when a reaches he wood. 8. Using a smooth motion, finnly squeeze both handles at the same time. This motion allows the bark to separate and the chemical to be deposited into the area. The WedgeChek[®] plags the hole, allowing the chemical to remain in the tree.
- Slowly withdraw the tip and repeat this step while moving around the circumference of the tree.
- 10. Move laterally and make the next injection until the entire tree trunk has been treated. The WedgeCheks'" may be left in the tree and used to inject the tree the following year

INDICATIONS


Indicated for the treatment of woody trees and shrubs for the control of Elm Leaf Beetles and Lace Bugs

DOSAGE 1 ml per 4* to 6* of trunk circumference. Note: Depending on type of bark, the dosage per injection site may have to be reduced to 1/2 ml per site and the number of injection sites doubled.

NOTICE OF WARRANTY

ArborSystems warrants that this product con-forms to the chemical description on the label and is reasonably fit for use under average and is reasonably the for use under average conditions when used strictly in accordance with the directions on the labeling. ArborSystems does not make or authorize any agent or representative to make any other warranty, guarantee or representation, express or implied, concerning this product.

For assistance contact



P.O. Box 34645 • Ornaha, NE 68134 Toll-Free: 1-800-698-4641 Phone: (402) 339-4459 FAX: (402) 339-5011 www.arborsystems.com

GREYHOUND^{***} Insecticide, Wedgle^{***} Tip and WedgeChek^{***} are trademarks of ArborSystems Wedgie " Tip is protected by U.S. Potern #5,239,773

WadowChek " is protected by U.S. Potent #5,797,215

A5_GH/IN 02-103 2 5M

.

,

1 . .

3

• · · · .

.

7

•.

Lэ

۵ 4,

ŧ

Federal Express May 20, 2002

Mr. Thomas C. Harris Document Processing Desk - *NOTIF* Office of Pesticide Programs - *7504C* **U.S. Environmental Protection Agency** 1921 Jefferson Davis Highway Crystal Mall 2, Room 266A Arlington, VA 22202-5601



Dear Mr. Harris:

Subject: ArborSystems, LLC Greyhound™ Insecticide EPA Reg. No. 69117-2

On behalf of its client **ArborSystems**, **LLC**, RegWest Company is submitting the attached materials as notification of a label revision for subject product. RegWest Company will act as sole agent in this endeavor.

Attached are the following:

- 1. Application for Pesticide: Other
- 2. One draft Reference Label
- 3. One copy of market labeling

The *Reference Label* illustrates the revisions as described below. The revisions are primarily being made due to the new packaging concept. Previously, for the professional market the product was sold in a 4- ounce container with an outsert (in a plastic sleeve) attached to the bottle. We will now be selling the product with the following components:

- 1. A "generic" box with the vertical self-adhesive label attached to the outside of the box.
- 2. The 4-ounce bottle with the horizontal self-adhesive label attached to it. The bottle is placed in the box.
- 3. The package insert that will be placed in the box with the product.

The following changes were made to the label (pages refer to the Reference Label):

- 1. Changed "outsert" to "insert" as a descriptor for where to find additional precautionary statements (page 1).
- 2. Changed "outsert" to "insert" as a descriptor for where to find complete Directions for Use (page 1).
- 3. Dropped ",LLC" from the company name. ArborSystems is still a limited liability corporation, however, for consistency they no longer desire the "LLC" to be a part of the logo and name on their marketing materials (pages 1, 2 and 4).
- 4. Changed "below" to "inside" as a descriptor for where to find additional precautionary statements (page 2)
- 5. Deleted redundant verbiage:
 - a. Omaha, NE 68134 added website (page 1);
 - b. "For assistance, contact ArborSystems at 402-339-4459" (page 3);
 - c. "Spring through late summer is the ideal time for injection" (page 3).
- 6. Changed "the chemical bottle outsert" to "this insert" (point number 5, page 3).
- 7. To the outer box added the verbiage about, and the graphic of, the injector gun.

Please contact me at (970) 353-0611 if you require additional information or have any questions.

Regards, RegWest Company

Kim Davis, CC, APC Consultant/Agent

30856 Rocky Road Greeley.CO 80631-9375 970-353-0611 970-353-0613 Fax regwestcc@aol.com

Attachments pc: ASI - Doolittle, C.