

69361-3

5/6/2014

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

May 6, 2014

N. Bhushan Mandava, Ph.D., Agent
Repar Corporation
c/o Mandava Associates, LLC
1050 Connecticut Ave., NW, Suite 1000
Washington, DC 20036

Subject: Label Amendments (change ingredients statement, create sub-labels for
consumer and professional uses)
Repar Broad Spectrum Weed and Feed
EPA Reg. No. 69361-3
Resubmission Dated March 14, 2014

Dear Dr. Mandava:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

A stamped copy of the label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", enclosed within a large, hand-drawn oval.

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs



Master Label *typo No. 5/2/14*

Net Wt. 17.5 Lbs. (7.93 kg)
Treats Up To 5,000 Sq. Ft.

REPAR BROAD SPECTRUM WEED & FEED

This label contains two separate sub-labels for, Residential Label (Sub-Label A) and Commercial Label (Sub-Label B) product use.

- For use on warm and cool season turf types as listed.
- Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds.
- For a thick green weed free lawn.

ACTIVE INGREDIENTS:

Chlorflurenol -Methyl 0.17%
 Dicamba 0.13%

OTHER INGREDIENTS:

..... 99.70%
 Total..... 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

ACCEPTED
05/06/2014
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 69361-3

Manufactured for:
 Repar Corporation
 P.O. Box 4321
 Silver Spring, MD 20914

EPA Reg. No. 69361-3
EPA Est. No.-

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.



Net Wt. 17.5 Lbs. (7.93 kg)
Treats Up To 5,000 Sq. Ft.

SUB-LABEL A: RESIDENTIAL LABEL

REPAR BROAD SPECTRUM WEED & FEED

- For use on warm and cool season turf types as listed.
- Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds (as listed).
- For a thick green weed free lawn.

ACTIVE INGREDIENTS:

Chlorfuretol-Methyl	0.17%
Dicamba	0.13%
OTHER INGREDIENTS:	99.70%
Total	100.00%

EPA Reg. No. 69361-3

EPA Est. No.-

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide of the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, except Desirable Clovers, Bent, Dichondra and Carpet Grass. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purpose.

WEED CONTROL

Repar Broad Spectrum Weed and Feed controls a wide range of perennial and annual weeds including Dandelion, Buckhorn Plaintain, Common Chickweed, White and Red Clover, Hop Clover, Marechal, Heal All, Henbit, Ground Ivy, Knotweed, Black Medic, Purslane, Yellow Rocket, Thistles, & other listed Broadleaf Weeds

WEED LIST

Annuals

Amaranth, Spiny (Spiny Pigweed)	Goosefoot, Nettleleaf	Pennygrass, Field (Fanweed)	Senna, Coffee
Aster, Slender	Henbit	Frenchweed	Sesbania, Hemp
Bedstraw	Jimsonweed	Stinkweed	Shephardspurse
Beggarweed, Florida	Knotweed	Pepperweed, Virginia (Peppergrass)	Sicklepod
Broomweed, Common	Kochia	Pigweed, Prostrate	Sida, Prickly (Teaweed)
Buckwheat, Wild	Kochia (triazine resistant)	P weed, Redroot	Smartweed, Green
Buffalobur	Ladyshumb	(Carelessweed)	Smartweed,
Burcucumber	Lambsquarters, Common	Pigweed, Rough (Carelessweed)	Pennsylvania
Buttercup, Roughseed	Lambsquarters, (triazine resistant)	Pigweed, Rough (Carelessweed)	Sneezeweed, Bitter
Carpetweed	Lettuce, Prickly	Pigweed, Tumble	Sowthistle, Annual
Catchfly, Nightflowering	Mallow, Venice	Poorjoe	Sowthistle, Spiny
Chamomile, Corn	Maresail	Puncturevine	Spanishneedles
Chickweed, Common	(Horseweed)	Purslane, Common	Spikeweed, Common
Clovers, Annual	Mayweed	Pursley, Florida	Spurge
Cockle, Corn	Medic Black	Radish, Wild	Spur, Corn
Cockle, Cow	Morningglory, Ivy/leaf	Ragweed, Common	Starbur, Bristly
Cocklebur, Common	Morningglory, Tall	Ragweed, Giant	Stumpweed, Rough
Croton, Tropic	Mustard, Tansy	(Buffaloweed)	Sunflower, Common
Croton, Woolly	Mustard, Wild	Rubberweed, Bitter	(Wild)
Daisy, English	Mustard (Yellowtops)	Nightshade, Black	Sunflower, Volunteer
Eveningprimrose,	Cutleaf		Thistle, Russian
Filebane, Annual			Velvetleaf
			Waterhemp
			Waterprimrose, Winged
			Wormwood, Annual

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear goggles, face shield or safety glasses and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) call CHEMTREC at 1-800-424-9300.

SEE REVERSE SIDE FOR ADDITIONAL INFORMATION

Biennials

Burdock, Common
 Carrot, Wild
 (Queen Anne's Lace)
 Cockle, White
 Eveningprimrose,
 Common

Plantain, Bracted
 Ragwort, Tansy
 Starthistle, Yellow
 Sweetclover
 Tassel

Thistle, Bull
 Thistle, Milk
 Thistle, Musk
 Thistle, Plumelless

Perennials

Alfalfa*
 Anticoke, Jerusalem
 Aster, Spiny
 Aster, Whitehead
 Bedstraw, Smooth
 Bindweed, Field
 Bindweed, Hedge
 Bindweed, Texas
 Bursage*
 (Bur Ragweed,
 Lakeweed,
 Goldenrod, Canada
 Goldenrod, Missouri
 Goldenweed, Common
 Hawkweed
 Henbane, Black
 Hosenettle, Carolina
 Ironweed
 Ivy, Ground
 Knapweed, Black
 Knapweed, Russian
 Chicory

Milkweed, Climbing
 Milkweed, Common
 Milkweed, Honeyvine
 Milkweed, Western
 Whorled
 (Eveningprimrose)
 Night, Singing
 Nightshade, Silverleaf
 (White Hosenettle)
 Onion, Wild
 Pennywort
 Vetch
 Violet, Wild
 Waterhemlock
 Waterprimrose
 Creeping
 Woodborer, Creeping*
 (Common Yellow)
 Wormwood, Common
 Wormwood, Louisiana
 Yankeweed*

Sowthistle
 Sowthistle, Perennial
 Spurge, Leafy
 Sundrop, Halfshrub
 (Eveningprimrose)
 Thistle, Canada
 Toadflax Dalmation
 Tropical Soda Apple
 Trumpet creeper
 (Buckvine)

* Noted perennials may be controlled using Repair Broad Spectrum Weed and Feed at rates lower than those directed for other listed perennial weeds.

Woody

Ailanthus
 Sassafras
 Oak, Poison
 Honeysuckle
 Huisache

Kudzu
 Serviceberry
 Brazilian Pepper
 Witchhazel

Sagebrush, Fringe
 Wild Grape
 Tatbush
 Huckleberry

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, except Desirable Clovers, Bent, Dichondra and Carpet Grass.

EFFECTIVE APPLICATION OF REPAIR BROAD SPECTRUM WEED AND FEED FOR MOST WEED LEAVES SHOULD BE DAMP FROM RAIN, DEW OR SPRINKLING TO ALLOW THE PRODUCT TO STICK TO THE LEAF. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL.

WHAT TO LOOK FOR:

Repair Broad Spectrum Weed and Feed, properly applied, first reduces or eliminates the formation of most flower heads and the seeds they produce and then attacks and removes the mature weed itself. Look for an absence or stunting of seed heads in flowering weeds such as clover or dandelions. Some weeds may also show twisting stems and dwarfed appearance with curling of the leaves.

Repair Broad Spectrum Weed and Feed is designed with an excellent turf nutrient diet to allow grass plants to gradually fill in the areas around dying weeds. This allows for a more natural recovery of the turf and minimizes reinfestations of bare areas by wind-borne weed seeds. Applied as directed, the rich green color will not be affected.

UNDER NORMAL GROWING CONDITIONS, WEED KILL SHOULD BE COMPLETE WITHIN 21 TO 35 DAYS

SPREADER SETTINGS

Spreader	Rate: 3.5#/M	Pattern Setting	Width
Earthway Broadcast	11	-	8
Lely	4 1/2	III - B	42
Lesco Broadcast	D 1/2	0.0	11
Pritzlawn Broadcast	4 1/4	A	7
Scots Broadcast	1	5	10
Spyker	3	-	9
Vicon	29	-	4
Cyclone Broadcast	3 3/4	-	5

Important: These are approximate settings. Age and condition of your spreader can cause wide variation. Be sure to calibrate your spreader with each application.

WHEN TO APPLY:

Apply any time weeds are actively growing and temperature is above 60°F. Do not apply when turf is under stress from heat, excessive moisture or drought. Control is quicker and more complete if applied in Spring prior to weed flower formation. Control is slower if applied when weather is cool. On newly seeded areas, wait until grass has been cut two or three times before using. Delay planting new grass for at least three weeks as the active ingredients may temporarily inhibit seed germination.

HOW TO APPLY:

EFFECTIVE APPLICATION OF REPAIR BROAD SPECTRUM WEED AND FEED REQUIRES THE WEED LEAVES TO BE DAMP FROM RAIN, DEW OR SPRINKLING PRIOR TO APPLICATION. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL. Apply evenly over the entire lawn at the rate of 3.5 lbs/1,000 sq ft. Rain or heavy sprinkling with 24 hours of application will temporarily delay weed control. Mowing within 24 hours may reduce effectiveness. Wait at least 45 days before making a second application. Do not make more than two applications per year.

Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over application and burning of grass. **SPECIAL NOTE:** Repair Broad Spectrum Weed And Feed is designed to be absorbed by plant leaves. Accidental application of this product to the foliage of ornamentals such as roses may result in the loss of flowers. For this reason, extra care should be taken when using a broadcast or rotary type spreader.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store in original container in a locked storage area. Keep container closed and away from food, feedstuffs and domestic water supplies.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION OF LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to the extent of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS. YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.
 Manufactured for:
 Repair Corporation, P.O. Box 4321
 Silver Spring, Maryland 20914



Net Wt. 17.5 Lbs. (7.93 kg)
Treats Up To 5,000 Sq. Ft.

SUB-LABEL B: COMMERCIAL LABEL

REPAR BROAD SPECTRUM WEED & FEED

- For use on warm and cool season turf types as listed.
- Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds (as listed).
- For a thick green weed free lawn.

ACTIVE INGREDIENTS:

Chlorfuretol-Methyl	0.17%
Dicamba	0.13%
OTHER INGREDIENTS:	99.70%
Total	100.00%

EPA Reg. No. 69361-3

EPA Est. No. -

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt and long pants, socks, shoes and chemical resistant gloves made of any waterproof material.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) call CHEMTREC at 1-800-424-9300.

FOR PRODUCT USE information call (202) 223-1424.

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any irrigation system.
Do not apply this product in a way that will contact any person, or animals, either directly or through drift. Only protected handlers may be in the area during application.
Do not allow people or animals to enter the treated area until dusts have settled.

WEED CONTROL

Repar Broad Spectrum Weed and Feed controls a wide range of perennial and annual weeds including Dandelion, Buckhorn Plantain, Common Chickweed, White and Red Clover, Hop Clover, Marechal, Heal All, Henbit, Ground Ivy, Knotweed, Black Medic, Purslane, Yellow Rocket, Thistles, & other listed Broadleaf Weeds

WEED LIST

Annuals

Amaranth, Spiny (Spiny Pigweed)	Goosefoot, Nettleleaf	Henbit	Penycress, Field (Fanweed)	Senna, Coffee
Aster, Slender	Jimsonweed	Knotweed	Frenchweed	Sesbania, Hemp
Bedstraw	Kochia	Kochia	Stinkweed	Shepherdspurse
Beggarweed, Florida	Kochia	Ladythumb	Pepperweed, Virginia (Peppergrass)	Sicklepod
Broomweed, Common	(triazine resistant)	Lambquarters, Common	Pigweed, Prostrate	Sida, Prickly (Teaweed)
Buckwheat, Wild	Lambquarters, Common	Lambquarters, Common	P weed, Redroot (Carelessweed)	Smartweed, Green
Burclover, California	Lambquarters, Common	Lambquarters, Common	Pigweed, Rough (Carelessweed)	Smartweed, Pennsylvania
Burcuumber	Lambquarters, Common	Lettuce, Prickly	Pigweed, Rough (Carelessweed)	Sneezeweed, Bitter
Buttercup, Roughseed	Lettuce, Prickly	Mallow, Common	Pigweed, Rough (Carelessweed)	Sowthistle, Annual
Carpetweed	Mallow, Venice	Mallow, Venice (Horseweed)	Pigweed, Tumble (triazine resistant)	Spikeweed, Common
Catchfly, Nightflowering	Mareslail	Mayweed	Poorjoe	Spurge
Chamomile, Corn	Mayweed	Medic, Black	Puncturevine	Spurr, Corn
Chickweed, Common	Medic, Black	Morningglory, Ivyleaf	Purslane, Common	Starbur, Brisly
Clovers, Annual	Morningglory, Ivyleaf	Morningglory, Tall	Pusley, Florida	Stumpweed, Rough
Cockle, Corn	Morningglory, Tall	Mustard, Tansy	Radish, Wild	Sunflower, Common (Wild)
Cockle, Cow	Mustard, Tansy	Mustard, Wild	Ragweed, Giant	Sunflower, Volunteer
Cocklebur, Common	Mustard, Wild	Mustard (Yellowjaps)	Ragweed, Common	Thistle, Russian
Croton, Tropic	Nightshade, Black	Nightshade, Black	Ragweed, Common	Thistle, Russian
Croton, Woolly			Ragweed, Common	Velvetleaf
Daisy, English			Ragweed, Common	Waterhemp
Eveningprimrose,			Ragweed, Common	Waterprimrose, Winged
Cutleaf			Ragweed, Common	Wormwood, Annual
Fleabane, Annual			Ragweed, Common	

SEE REVERSE SIDE FOR ADDITIONAL INFORMATION

5/6

Biennials

Burdock, Common
Carrot, Wild
(Queen Anne's Lace)
Cockle, White
Eveningprimrose,
Common

Perennials

Alfalfa*
Atrichoke, Jerusalem
Aster, Spiny
Bedstraw, Smooth
Birdweed, Field
Bindweed, Hedge
Bindweed, Texas
Bursage*
(Bar Ragweed,
Lakeweed,
Povertyweed)
Bursage, Woollyleaf
(Lakeweed)
Buttercup, tall
Campion, Bladder
Chickweed, Field
Chickweed, Mouseear
(Canada)
Chicory

Clover, Hop*
Dandelion, Common*
Dock, Broadleaf*
(Bitterdock)
Dock, Curly*
Dogbane, Hemp
Dogfennel*
(Cypressweed)
Fern, Bracken
Garlic, Wild
Goldenrod, Canada
Goldenrod, Missouri
Goldenweed, Common
Hawkweed
Henbane, Black
Horsenettle, Carolina
Ironweed
Ivy, Ground
Knapweed, Black
Knapweed, Russian

Plantain, Bracted
Ragwort, Tansy
Spearhead, Yellow
Sweetclover
Teasel

Milkweed, Climbing
Milkweed, Common
Milkweed, Honeyvine
Milkweed, Western
Whorled
Nettle, Stinging
Nighthshade, Silverleaf
(White Horsenettle)
Onion, Wild
Pennywort
(Dollarweed)
Plantain, Broadleaf*
Plantain, Buckhorn
Pokeweed
Ragweed, Western
Sericia Lespedeza
Smartweed, Swamp
Snakeweed, Broom
Sorrel, Red*
(Sheep Sorrel)

Sowthistle
Sowthistle, Perennial
Spurge, Leafy
Sundrop, Halfshrub
(Eveningprimrose)
Thistle, Canada
Toadflax, Dalmation
Tropical Soda Apple
Trumpetcraper
(Buckvine)
Vetch
Violet, Wild
Waterhemp
Waterprimrose
Creeping
Woodsorrel, Creeping*
(Common Yellow)
Wormwood, Common
Wormwood, Louisiana
Yankee weed*
Yarrow, Common

* Noted perennials may be controlled using Repair Broad Spectrum Weed and Feed at rates lower than those directed for other listed perennial weeds.

Woody

Allianthus
Sassfras
Oak, Poison
Honeysuckle
Huisache

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, **except** Desirable Clovers, Bent, Dichondra and Carpet Grass.

EFFECTIVE APPLICATION OF REPAIR BROAD SPECTRUM WEED AND FEED FOR MOST WEED LEAVES SHOULD BE DAMP FROM RAIN, DEW OR SPRINKLING TO ALLOW THE PRODUCT TO STICK TO THE LEAF. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL.

WHAT TO LOOK FOR:

Repair Broad Spectrum Weed and Feed, properly applied, first reduces or eliminates the formation of most flower heads and the seeds they produce and then attacks and removes the mature weed itself. Look for an absence or stunting of seed heads in flowering weeds such as clover or dandelions. Some weeds may also show twisting stems and dwarfed appearance with curling of the leaves.

Repair Broad Spectrum Weed and Feed is designed with an excellent turf nutrient diet to allow grass plants to gradually fill in the areas around dying weeds. This allows for a more natural recovery of the turf and minimizes reinfestations of bare areas by wind-borne weed seeds. Applied as directed, the rich green color will not be affected.

UNDER NORMAL GROWING CONDITIONS, WEED KILL SHOULD BE COMPLETE WITHIN 21 TO 35 DAYS

SPREADER SETTINGS

Spreader	Rate: 3.5#/M	Pattern Setting	Width
Earthway Broadcast	11	-	8
Lely	4 1/2	III - B	42
Lesco Broadcast	D 1/2	0.0	11
Prizelawn Broadcast	4 1/4	A	7
Scotts Broadcast	1	5	10
Spyker	3	-	9
Vicon	29	-	4
Cyclone Broadcast	3 3/4	-	5

Important: These are approximate settings. Age and condition of your spreader can cause wide variation. Be sure to calibrate your spreader with each application.

WHEN TO APPLY:

Apply any time weeds are actively growing and temperature is above 60°F. Do not apply when turf is under stress from heat, excessive moisture or drought. Control is quicker and more complete if applied in Spring prior to weed flower formation. Control is slower if applied when weather is cool. On newly seeded areas, wait until grass has been cut two or three times before using. Delay planting new grass for at least three weeks as the active ingredients may temporarily inhibit seed germination.

HOW TO APPLY:

EFFECTIVE APPLICATION OF REPAIR BROAD SPECTRUM WEED AND FEED REQUIRES THE WEED LEAVES TO BE DAMP FROM RAIN, DEW OR SPRINKLING PRIOR TO APPLICATION. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL. Apply evenly over the entire lawn at the rate of 3.5 lbs./1,000 sq. ft. Rain or heavy sprinkling with 24 hours of application will temporarily delay weed control. Mowing within 24 hours may reduce effectiveness. Do not make more than one application per year. Not for use on sod farms.

Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over application and burning of grass. SPECIAL NOTE: Repair Broad Spectrum Weed And Feed is designed to be absorbed by plant leaves. Accidental application of this product to the foliage of ornamentals such as roses may result in the loss of flowers. For this reason, extra care should be taken when using a broadcast or rotary type spreader.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store in original container in a locked storage area. Keep container closed and away from food, feedstuffs and domestic water supplies.
PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION OF LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.
LEGAL RIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Manufactured for:

Repair Corporation, P.O. Box 4321
Silver Spring, Maryland 20914