



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**  
WASHINGTON, DC 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Repar Corporation  
% Bushan Mandava  
Mandava Associates, LLC  
1730 M Street, N.W., Suite 806  
Washington, DC 20036-4510

JUN 26 2008

Subject: Tebucon 3.6F Fungicide  
EPA Reg. No. 69361-11  
Your amendment dated May 6, 2008  
EPA Decision Number 393399

Dear Dr. Mandava:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

1. Qualify registrant name and address by adding "Manufactured for:"
2. Add the heading **DIRECTIONS FOR USE** immediately above the misuse statement.
3. Add an "ENVIRONMENTAL HAZARDS" section:

**"ENVIRONMENTAL HAZARDS"**

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

**Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours."

Repar Corporation  
P.O. Box 4321  
Silver Spring, MD 20914  
(202) 223 - 1424

Tebucon 3.6F Fungicide

EPA Reg. No. 69361-11

For Use on Wheat and Barley

## Supplemental Label

# TEBUCON 3.6F FUNGICIDE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Tebucon 3.6F Fungicide.

### USE DIRECTIONS FOR SPECIFIC CROPS

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF TEBUCON 3.6 F
Barley	Rusts ( <i>Puccinia spp.</i> )	4 fl oz per acre
	Head blight ( <i>Fusarium spp.</i> ) - Suppression	
	Notes: Apply TEBUCON 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of TEBUCON 3.6F. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.	
Application timing directions: Rusts: Apply TEBUCON 3.6F at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of TEBUCON 3.6F for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.		
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with TEBUCON 3.6F. TEBUCON 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCON 3.6F will be resistant to weathering. TEBUCON 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).		

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**

6/26/2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.

69361-11

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF TEBUCON 3.6 F
Wheat	Rusts leaf, stem, and stripe. ( <i>Puccinia spp.</i> )	4 fl oz per acre
	Head blight or scab ( <i>Fusarium spp.</i> ) — Suppression	
	Notes: Wheat fields should be observed for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of TEBUCON 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with TEBUCON 3.6F. Apply TEBUCON 3.6F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.	
	Application timing directions: Rusts: Apply TEBUCON 3.6F at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of TEBUCON 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with TEBUCON 3.6F. TEBUCON 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCON 3.6F will be resistant to weathering. TEBUCON 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).		

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.  
For PRODUCT USE information Call (202) 223 - 1424

As with any crop-protection product, always read and follow label directions.  
For additional information call (202) 223 - 1424