



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

69361-46

Date of Issuance:

10/07/2014

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Repar Permethrin H&G

Name and Address of Registrant (include ZIP Code):

**Madhu Mandava
Agent for Repar Corporation
c/o Mandava Associates, LLC
6860 N. Dallas Parkway, Suite 200
Plano, TX 75024**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continued on page 2

Signature of Approving Official:

Richard Gebken, Product Manager 10, Insecticide Branch
Registration Division (7504P)

Date:

10/07/2014

2. You are required to comply with the DCI Order identified below:

- a. Permethrin GDCI-109701-1281, issued on 02/26/2013
- b. Permethrin GDCI-109701-1113, issued on 02/01/2012

You must comply with all of the data requirements within the deadlines established by the order. If you have questions about the Generic DCI orders listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

http://www.epa.gov/oppsrrd1/contacts_prd.htm

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.
4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 69361-46.”
5. Submit one copy of the final printed label for the record before you release the product for shipment.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 11/19/2013

If you have any questions, please contact Mr. Driss Benmhend, at (703) 308-9525 or via email at benmhend.driss@epa.gov.

Sincerely,



Richard Gebken
Product Manager 10
Insecticide Branch
Office of Pesticide Programs
Date: October 7, 2014


Enclosure: Label Stamped “accepted”

Restricted Use Pesticide

Due to acute primary skin irritation.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

REPAR PERMETHRIN H & G
Home and Garden Uses
Outdoor Use Only



ACTIVE INGREDIENT:

Permethrin* 38.0%

OTHER INGREDIENTS:** 62.0%

TOTAL: 100.0%

* cis/trans ratio: Max. 55% (+/-) cis and min. 45% (+/-) trans

** Contains Petroleum Distillates.

KEEP OUT OF REACH OF CHILDREN
WARNING - AVISO

See Back/Side Panel for Additional Precautionary Statements

FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 or 1-800-900-4044 for emergency medical treatment information.	
NOTE TO PHYSICIAN: This product contains petroleum distillates. May pose an aspiration pneumonia hazard. Consideration should be given to gastric lavage with an endotracheal tube in place. Treatment is controlled removal of exposure followed by symptomatic and supportive care.	

Repar Corporation
P. O. Box 4321
Silver Spring, MD 20914

EPA Reg. No.: 69361-
Net Contents:

EPA Est. No.:

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Do not get on skin or clothing. Avoid contact with eyes. Avoid breathing spray mist or vapor. Wear coveralls over short sleeved shirt and short pants, chemical resistant footwear and socks, chemical resistant gloves (barrier laminate or Viton, selection category G). When mixing and loading wear a chemical resistant apron. For overhead exposure wear chemical resistant headgear. When cleaning equipment add a chemical resistant apron.

(people/pet icon) – (optional) People and pets may enter treated area after application has dried.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

Applicators using UL V cold foggers or fog/mist generators in indoor spaces must wear:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves,
- Chemical-resistant footwear plus socks, and
- Chemical-resistant headgear, if overhead exposure

All other mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks
- Chemical-resistant gloves for all handlers except for applicators using motorized ground equipment, pilots and flaggers,
- Chemical-resistant apron for mixers/loaders, persons cleaning equipment, and persons exposed to the concentrate.

See Engineering Controls for additional requirements.

USER SAFETY RECOMMENDATIONS:

- User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Failure to follow the directions for use and precautions on this label may result in poor pest control, plant injury or illegal residues.

PRODUCT INFORMATION

This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding, in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

This product may be used to control insect pests outdoors, on ornamentals, lawns, trees, shrubs, and vines in landscape areas, as a perimeter treatment around residential buildings. This product may also be used to control home-invading insects as a residual spray on outside surfaces, such as walks, garbage areas, window frames, doorways, porches, and patios. Not for use on plants being grown for commercial sale, or plants grown for seed production or sod farms and golf course turf, or in greenhouses or nurseries.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- 1) Treatment to soil or vegetation around structures;
- 2) Applications to lawns, turf, and other vegetation;
- 3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

Not for use in outdoor residential misting systems.

Application Instructions: This product mixes readily with water and is specifically packaged for residential use as a diluted spray. Use a clean sprayer, such as a tank type or hose-end sprayer to mix and apply the proper amount of concentrate with the desired volume of water. For perimeter and lawn applications measure the area to be treated to determine the application volume. For trees, shrubs and flowers apply as a thorough cover spray until foliage is wet. Refer to **Specific Application Instructions** in the application rate Tables for additional instructions.

Entry restriction for products applied as a spray except for sprays applied directly to domestic animals: Do not allow adults, children, or pets to enter the treated area until sprays have dried.

HOME-INVADING PESTS (OUTDOOR USE ONLY)

MIXING INSTRUCTIONS

Mix 1.6 fl. oz. (3 ¼ Tablespoons) of this product per 1 gallon of spray solution to make a 0.5% spray emulsion (1 fl. oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)). Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Spot Treatment

- Apply a 0.5% emulsion as a pin stream low-pressure spray or as a paint-on application with a paintbrush.
- Treat where pests are found or entry points of the structure such as window frames, doorways, porches, patios, and foundation.
- Spot-treatment may also be made to garages, garbage cans, and garbage areas.
- DO NOT apply indoors (including crawl spaces).

Perimeter Application

- Spray a 6 to 8 foot wide band of soil around the perimeter of the house. Treat the foundation wall to a height of 2 to 3 feet. Use a spray volume of 2 to 10 gallons of 0.5% emulsion per 1,000 square feet. If mulch or debris is present, a higher volume application rate may be needed to ensure adequate coverage.
- Note: residual control will vary depending on exposure of the treated area to weathering conditions (temperature, sunlight, moisture). Retreat on a monthly schedule while insects are likely to be active.

Wood Destroying Pests

- Localized Areas Outdoors: For the control of insects such as termites, bees, carpenter ants and wood infesting beetles in localized above-ground areas of infested wood around structures, apply a 0.5% emulsion to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Apply either as a spray or paint-on application. For termites the purpose of application is to destroy workers or winged reproductive forms, which may be present at the time of treatment. This product is not designed for preventative use and should only be applied when pests appear. This application is not a substitute for soil or foundation treatment. For active infestations inside a structure, consult a licensed Pest Control Service.
- Use a hose-end sprayer or sprinkling can to deliver a coarse drenching spray to treat stored lumber or woodpiles. Stored wood can be used for lumber one month after treatment. Do not apply inside structures.
- Repeat treatment as necessary, but not more frequently than once a month. For active termite infestation consult a professional Pest Control Applicator.

HOME-INVADING PESTS (OUTDOOR USE ONLY)	
PEST	SPECIFIC APPLICATION INSTRUCTIONS
Ants (including mounds) Armyworms Bark Beetles Bees Beetles Borers Boxelder Bugs Carpenter Ants Carpenter Bees Centipedes Chinchbugs Cockroaches (Asian) Crickets Earwigs Elm Leaf Fire Ants (Including mounds) Fleas Flies Firebrats Ground Beetles Gypsy Moths (adults & Caterpillars) Millipedes Mole Crickets Pill Bugs Scorpions Silverfish Sod Webworms Sowbugs Spiders Ticks (including Deer Tick, Western Black legged Tick which may carry Lyme disease And Rocky Spotted Fever) Wasps	Boxelder Bugs, Elm Leaf Beetles, and Gypsy Moth Caterpillars: Spray tree trunks, building siding or wherever pests congregate. Avoid excess application. Dampen surface thoroughly, but do not let runoff occur. Borers and Bark Beetles: To prevent infestation of ornamental trees and woody shrubs, spray the bark to the point of saturation, but do not let runoff occur. Repeat treatment every 21 days during early spring until mid-summer. Carpenter Bees: Apply as a pin stream, as a fine/coarse low-pressure spray (20 psi or less) as a spot treatment. Fire Ants (mound drench): Apply 1 to 2 gallons of emulsion to each mound area by sprinkling the mound until it is wet and treat a 4-foot diameter circle around the mound. Use higher volume for mounds larger than 12 inches. For best results, apply in cool weather such as in early morning or late evening hours, but not in the heat of the day. Scorpions: Remove accumulations of lumber, firewood, and other materials which serve as harborage sites and treat beneath and adjacent to infested sites.

LAWN PESTS

This product can be applied to lawns adjacent to or around private homes, duplexes, townhouses, condominiums, house trailers, carports, garages, fence lines, storage sheds, barns, and other residential areas where pests congregate.

MIXING INSTRUCTIONS

Prepare an emulsion by mixing 0.4 to 0.8 fl. oz. (2½ to 5 teaspoons) of this product in sufficient volume of water to cover 1,000 sq. ft. (approximately 4 to 25 gallons of water). (1 fl. oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)). Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Lawn Broadcast Application,

- Apply using a hose-end or pressure type sprayer to deliver a coarse spray of between 4 to 25 gallons of emulsion per 1,000 sq. ft. of lawn.
- Use the higher application rate for heavy infestations.
- Use sufficient volume to uniformly cover plant surface.

Note: Higher volume of water should be used when situations of thick thatch or dense sod exist or in and or drought conditions.

WHEN TO APPLY

- For best results treat in early morning or late evening hours, but not in the heat of the day.
- Spray evenly over the lawn when insects first appear.
- Repeat applications as necessary but not more frequently than every 10-14 days. Do not apply more than 1.6 fl. oz./1,000 sq. ft. per year.

MOUND DRENCH TREATMENT (Ant and Fire Ant Mounds in Lawns)

Use a sprinkling can. Mix 1.6 fl. oz. (3 1/4 Tablespoons) of this product per 1 gallon of spray solution to make a 0.5% spray emulsion {1 fl. oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)}. For large ant mounds, such as Fire Ant mounds, apply 1-2 gallons of a 0.5% emulsion to each mound area including a 4-foot diameter circle around the mound. Apply to smaller mounds until the mound is completely saturated. Use higher water volume for mounds larger than 12". For best results, treat in early morning or late evening hours, but not in the heat of the day.

APPLICATION RATES FOR LAWN			
SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
Lawns, residential including, but not limited to, St. Augustine, Bentgrass, Fescues, Buffalograss, Bahiagrass, Bluegrass, Perennial Ryegrass, and Centipedegrass.	Ants Armyworms Billbugs Chiggers Chinchbugs Crickets Cutworms Earwigs Fruit Flies Fleas Leafhoppers Millipedes Mole Crickets Sod Webworm Sowbugs Ticks (including Brown Dog Tick, Deer Tick, Western Blacklegged Ticks which may carry Lyme disease and Rocky	0.4 to 0.8 fl. ozs. (2 1/2 to 5 tsp.) Per 1,000 sq. ft.	Apply using 4-25 gallons of spray volume/1,000 sq. ft. of lawn. Use higher volumes on St. Augustine, Buffalograss, Bermuda and Bahiagrass lawns. Use the higher rate for fast knockdown and increased residual control. Dense and Excessive (greater than 3 inches) lawn height and arid conditions may require higher volume application rates. Repeat applications may be necessary (10 to 14 day interval) to control heavy infestations during peak periods of insect feeding. Do not exceed a total of 1.6 fl. ozs./1,000 sq. ft. per year (i.e. four (4) applications at the lower rate or two (2) applications at the

	<p>Mountain Spotted Fever)</p> <p><i>For additional pests consult the list of pests in the table "Home Invading Pests"</i></p> <p><i>(Outdoor Use Only)</i></p>		<p>higher rate).</p>
--	---	--	----------------------

ORNAMENTAL PESTS

MIXING INSTRUCTIONS

Mix 0.04 to 0.08 fl. oz. (= 1/8 to 1/4 teaspoons = 1-2 ml.) of this product in 1 gallon of water for general pest control. Refer to the Ornamental Application Rate Table for additional mixing instructions and specific directions for stem-infesting beetles. (1 fl. oz. = 2 Tablespoons (T) = 6 teaspoons (tsp) = 29.6 ml.). Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Ornamentals (Foliage and Flowering Plants)

- Apply using a hose-end or pressure type sprayer to deliver a coarse spray. Apply thoroughly to stem and foliage.
- Use the higher application rate (1/4 tsp. per gal. of water) for heavy infestations. Note: Higher volumes of water should be used when situations of dense foliage exists, or in and or drought conditions.

WHEN TO APPLY

- For best results treat in early morning or late evening hours, but not in the heat of the day.
- Spray evenly over the lawn when insects first appear.
- To prevent insect reinfestation repeat treatments as necessary but not more frequently than once every 14 days.

APPLICATION RATE FOR USE ON ORNAMENTALS			
SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
<p>Ornamental Foliage and Flowering plants (including Fern conifers, woody and herbaceous non-edible Ornamentals)</p>	<p>Ants Aphids Bagworm Beet Armyworm Cabbage Looper Cankerworms Citrus Thirps Coneworm (including Web) and seed bugs Fungus Gnat Gypsy Moth Caterpillars Heliothis spp. Japanese Beetles</p>	<p>0.04 to 0.08 fl. ozs. (1/8 to ¼ tsp.) (1 to 2 ml.) Per gallon</p>	<p>Note: Direct application to blooms may cause browning of petals. Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris</p> <p>Mosquitoes coming to rest on treated plants will be killed for varying periods of time after treatment, depending on exposure of treated areas to weather conditions.</p>

	Lace Bugs Leaf Feeding Caterpillars Leaf-footed bugs Leafminers Leafrollers Lygus Bugs Mealy Bugs Mosquitoes Nantucket Pine Tip Moth Oblique Leafroller Pine Sawflies Plant Bugs Root Weevils (Adult) Rosy Aphid Rose Chafer Spotted Leafminer Stink Bugs Tarnished Plant Bug Tent Caterpillars Webworms Whiteflies Zimmerman Pine Moth		To control Webbing Coneworm make the first application just prior to peak pollen flight. To control other Coneworms and Seed Bugs, make application 30 days prior to flower closure.
--	--	--	---

APPLICATION RATE FOR USE ON ORNAMENTALS			
SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
Ornamental Foliage and Flowering plants (including Fern conifers, woody and herbaceous non-edible Ornamentals)	Bark Beetles and Boring insects (including, but not limited to, Ash Borer, Bronze Birch Borer, Rhododendron Borer, Elm Bark Beetles and Turpentine Beetles	0.3 to 0.6 fl. ozs. (2-4 tsp.) Per gallon	Stem-Infesting Beetles: Treat lower branches and trunk prior to adult emergence. Complete, uniform coverage is needed for best control. A repeat application may be necessary on a 3 to 4 week interval to prevent severe tree stem damage.

TERMITE CONTROL (SUBTERRANEAN)

This product may be used to establish an exterior soil barrier against termites around the structure. The chemical concentrate is mixed as an emulsion and adequately dispersed in the soil to provide a continuous exterior barrier between wood in the structure and the termite colonies in the soil.

Note: A complete termite barrier may require treatment with other EPA registered products to the inside perimeter of the foundation and to other interior critical areas. For these areas, and if termites are found inside of the home, consult a Licensed Professional Pest Control Service for treatment. For use only around existing structures. Do not use inside dwelling. Not for Preconstruction use.

PRECAUTION: When treating adjacent to an existing structure check the area to be treated; and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. If there are any signs of leakage into the structure during application have all people and pets leave the structure. After application, check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the cleanup is completed.

MIXING INSTRUCTIONS

Using a sprinkler can add 3.2 fl. oz. (6.5 Tablespoons) to 2 gallons of water (to make a 0.5% emulsion) for each 5 linear feet of treatment to be provided. {1 fl. oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)}. Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Trench Treatment

- Remove all wood debris and wood forms. Around piers, chimney bases and along foundations of poured concrete, dig a trench 6 inches wide and 4 inches deep. If footing is more than 12 inches deep drill holes with a steel rod or pipe one foot apart that extend from the bottom of the trench to the top of the footing. Make holes closer in hard packed clay soils. **Never dig below the top of the footing.**
- Apply a 0.5% emulsion at a rate of 4 gallons per 10 linear feet per foot of depth. Apply half of the emulsion when backfilling the trench. (One pint of this product will make 10 gallons of 0.5% emulsion and treat 25 linear feet).

Precaution: DO NOT treat structures that contain cisterns or wells. Soil in the vicinity of wells, cisterns or ponds or structures with sump pumps or French drains should not be treated. For those situations, contact your County Cooperative Extension Agent or consult a commercial pest control service.

Fence Posts

- Pre-installation: Using a sprinkler can, pour 1/2 gallon of 0.5% spray emulsion into the fence post hole and add another 1/2 gallon in mixture with soil when filling the hole so that all the soil is treated.
- Post-Installation: Installed poles and posts may be treated by first digging a trench around the post and applying 1 gallon of 0.5% emulsion per one foot depth of trench for posts less than 6 inches in diameter. For larger diameter posts increase the application rate to 4 gallons for each 10 linear feet of trench and per foot of depth.

WHEN TO RE-APPLY

Retreatment for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide barrier in the soil. These vulnerable or reinfested areas maybe retreated in accordance with application techniques described in the product's labeling. The timing and type of these

retreatments will vary, depending on factors such as termite pressure, soil types, soil conditions and other factors, which may reduce the effectiveness of the barrier.

Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.

IMPORTANT

DO NOT use on vegetables or food crops. DO NOT treat pets with this product. DO NOT spray inside dwellings. DO NOT treat soil that is water saturated or frozen or in any conditions where run-off or movement from the treatment area (site) is likely to occur. DO NOT apply when wind velocity favors drift.

STORAGE AND DISPOSAL

Pesticide Storage: Keep out of the reach of children or animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. Carefully open containers. After partial use replace lid and close tightly. Do not put concentrate or dilute material into food or drink containers.

Pesticide Disposal:

If empty: Non-refillable container. Do not reuse container. Do not rinse unless required for recycling. Place in trash.

If partly filled: Call your local solid waste agency for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of REPAR CORPORATION or the SELLER. All such risks shall be assumed by the Buyer.

REPAR CORPORATION warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks referred to above.

REPAR CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, REPAR CORPORATION AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. REPAR CORPORATION and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR CORPORATION.

List of (Optional) User Friendly Statements to be added to the Front and /or Back/Side Final Printed Label:

Use Statements

Insecticide
Insecticide/Termiticide
Termiticide
Lawn & Ornamental
Soil Insect & Termite Killer
Fire Ant Killer
Ant & Roach Killer
Outdoor Perimeter
For Outdoor Residential Use Only

Control Statements

Kill Ticks that may carry Lyme disease
Kills Home Invading Pests (Outdoor)
Kills Named Insects
An easy to apply product to control wood invading insects
Termiticide Formulation for Homeowner Use (Outdoor Use Only)
Termite Insecticide (Outdoor Use Only)
Homeowner Use (Outdoor Use Only)
Controls Wood-Destroying Insects

Icons/Pictures

(Icon of People)
(Icon of Sprinkling Can)
(Icon of Hand Held Sprayer)
(Tree Icon)
(Plant Icon)
UPC Symbology
Picture of a lawn
Pictures of one or more of the following insects: Termites,
Ants (including Fire ant mounds and Carpenter Ants),
Cockroach, spider, scorpion, fleas, tick, mosquito

Other

Concentrate
Treats up to (#will depend on application) Square Feet
Made in the USA
Formulated and Packaged in the USA
How to Apply
When to Apply
Amount to Use