

69407-1

11/07/2000

1/2

69407-1 New Label: Page 1 of 2
10/17/00**FLEA, TICK AND EAR MITE TREATMENT****For Use on Cats and Dogs to Control Fleas, Ticks and Ear Mites****For Use on Premises to Control Fleas and Ticks****An Aqueous, Stable Microemulsion****Prevents Dry Skin****ACTIVE INGREDIENTS:**

Pyrethrins (CAS No. 8003-34-7) 0.15%

Piperonyl Butoxide, Technical (CAS No. 51-03-6) 1.50%

OTHER INGREDIENTS 98.35%

Total 100.00%

Keep Out of Reach of Children**CAUTION**

See back panel for additional precautionary statements.

ACCEPTED

NOV 7 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 69407-1

Net Contents: ____ fl. oz.

EPA Reg. No. 69407-1

EPA Est. 69407-CA-1

ANIMAL DERMATOLOGY LABORATORIESP.O. Box 19097
Irvine, CA 92713

READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS AND CATS [DOGS, CATS AND LIVESTOCK]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage: Store in original container in a cool, dry (preferably locked) place inaccessible to children and pets. **Disposal:** Do not reuse empty container; wrap and put in trash.

SHAKE WELL BEFORE USE

For use on pets to control fleas and ticks: Start spraying at the tail, rapidly moving the dispenser and making sure that the animal's body is covered, including the legs and underbody. While spraying, fluff the hair so that spray will penetrate to the skin. Make sure spray thoroughly wets ticks. Do not spray into eyes, face or genitals. Do not repeat treatment for three days. **For use on pets to control ear mites:** Place ten drops in each ear and massage. Use twice a week for three weeks.

Do not use on puppies or kittens under 12 weeks. Consult a veterinarian before using this product on debilitated, medicated, aged, pregnant or nursing animals.

For use on premises: To control *Fleas* and *Brown Dog Ticks* thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

For use on livestock: To protect cattle, horses and hogs from *Horn Flies*, *House Flies*, *Mosquitoes* and *Gnats*, apply a light mist sufficient to wet the surface of the hair. To control *Stable Flies*, *Horse Flies* and *Deer Flies* apply at the rate of 2 ounces per adult animal, sufficient to thoroughly wet the hair. Repeat treatment daily or at intervals necessary to give continued protection. To control *Blood Sucking Lice* apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To control *Poultry Lice* thoroughly spray roosts, walls and nests or cages, then spray a fine mist over the birds. For control of *Bedbugs* and *Mites* in poultry houses, spray crevices of roost poles, cracks in walls and cracks in nests where these insects hide.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Humans: Harmful if swallowed. Avoid inhalation of vapors. Thoroughly wash with soap and water after use. Avoid spraying food or foodstuffs. When used as premise spray indoors, remove pets, birds and cover fish aquariums before spraying. **Animals:** Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, immediately consult a veterinarian.

First Aid

If Swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-521-2525 for emergency medical treatment information.	

To obtain more information or in case of an emergency call (800) 521-2525.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}